


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

BC

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07D 209/42, A61K 31/40, C07D 403/12, 401/12, 417/12, 401/14, 409/12, 413/12	A1	(11) International Publication Number: WO 96/39384 (43) International Publication Date: 12 December 1996 (12.12.96)
(21) International Application Number: PCT/IB95/00442 (22) International Filing Date: 6 June 1995 (06.06.95) (71) Applicant (for all designated States except US): PFIZER, INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HULIN, Bernard [FR/US]; Pfizer Inc., Eastern Point Road, Groton, CT 06340 (US). HOOVER, Dennis, J. [US/US]; Pfizer Inc., Eastern Point Road, Groton, CT 06340 (US). TREADWAY, Judith, L. [US/US]; Pfizer Inc., Eastern Point Road, Groton, CT 06340 (US). MARTIN, William, H. [US/US]; Pfizer Inc., Eastern Point Road, Groton, CT 06340 (US). PHILLIPS, Douglas [US/US]; Pfizer Inc., Eastern Point Road, Groton, CT 06340 (US). (74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., Patent Dept., 235 East 42nd Street, New York, NY 10017 (US).		(81) Designated States: CA, FI, JP, MX, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: SUBSTITUTED N-(INDOLE-2-CARBONYL)-GLYCINAMIDES AND DERIVATIVES AS GLYCOGEN PHOSPHORYLASE INHIBITORS <div style="text-align: center;">  </div>		
(57) Abstract Compounds of Formula (1) wherein R ₆ is carboxy, (C ₁ -C ₈)alkoxycarbonyl, benzyloxycarbonyl, C(O)NR ₈ R ₉ or C(O)R ₁₂ as glucogen phosphorylase inhibitors, pharmaceutical compositions containing such inhibitors and the use of such inhibitors to treat diabetes, hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemia, hyperlipidemia, atherosclerosis and myocardial ischemia in mammals.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

SUBSTITUTED N-(INDOLE-2-CARBONYL)-GLYCINAMIDES AND DERIVATIVES AS GLYCOGEN PHOSPHORYLASE INHIBITORS

Background of the Invention

This invention relates to glycogen phosphorylase inhibitors, pharmaceutical
5 compositions containing such inhibitors and the use of such inhibitors to treat diabetes, hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemia, hyperlipidemia, atherosclerosis and myocardial ischemia in mammals.

In spite of the early discovery of insulin and its subsequent widespread use in the treatment of diabetes, and the later discovery of and use of sulfonylureas (e.g.
10 Chlorpropamide™ (Pfizer), Tolbutamide™ (Upjohn), Acetohexamide™ (E.I. Lilly), Tolazamide™ (Upjohn)) and biguanides (e.g. Phenformin™ (Ciba Geigy), Metformin™ (G. D. Searle)) as oral hypoglycemic agents, the treatment of diabetes remains less than satisfactory. The use of insulin, necessary in about 10 % of diabetic patients in which synthetic hypoglycemic agents are not effective (Type I diabetes, insulin dependent
15 diabetes mellitus), requires multiple daily doses, usually by self injection. Determination of the proper dosage of insulin requires frequent estimations of the sugar in urine or blood. The administration of an excess dose of insulin causes hypoglycemia, with effects ranging from mild abnormalities in blood glucose to coma, or even death. Treatment of non-insulin dependent diabetes mellitus (Type II diabetes, NIDDM) usually
20 consists of a combination of diet, exercise, oral agents, e.g. sulfonylureas, and in more severe cases, insulin. However, the clinically available hypoglycemics can have other side effects which limit their use. In any event, where one of these agents may fail in an individual case, another may succeed. A continuing need for hypoglycemic agents, which may have fewer side effects or succeed where others fail, is clearly evident.

25 Atherosclerosis, a disease of the arteries, is recognized to be the leading cause of death in the United States and Western Europe. The pathological sequence leading to atherosclerosis and occlusive heart disease is well known. The earliest stage in this sequence is the formation of "fatty streaks" in the carotid, coronary and cerebral arteries and in the aorta. These lesions are yellow in color due to the presence of lipid deposits
30 found principally within smooth-muscle cells and in macrophages of the intima layer of the arteries and aorta. Further, it is postulated that most of the cholesterol found within the fatty streaks, in turn, give rise to development of the "fibrous plaque", which consists of accumulated intimal smooth muscle cells laden with lipid and surrounded by extra-

-2-

cellular lipid, collagen, elastin and proteoglycans. The cells plus matrix form a fibrous cap that covers a deeper deposit of cell debris and more extra cellular lipid. The lipid is primarily free and esterified cholesterol. The fibrous plaque forms slowly, and is likely in time to become calcified and necrotic, advancing to the "complicated lesion" which
5 accounts for the arterial occlusion and tendency toward mural thrombosis and arterial muscle spasm that characterize advanced atherosclerosis.

Epidemiological evidence has firmly established hyperlipidemia as a primary risk factor in causing cardiovascular disease (CVD) due to atherosclerosis. In recent years, leaders of the medical profession have placed renewed emphasis on lowering plasma
10 cholesterol levels, and low density lipoprotein cholesterol in particular, as an essential step in prevention of CVD. The upper limits of "normal" are now known to be significantly lower than heretofore appreciated. As a result, large segments of Western populations are now realized to be at particular high risk. Such independent risk factors include glucose intolerance, left ventricular hypertrophy, hypertension, and being of the
15 male sex. Cardiovascular disease is especially prevalent among diabetic subjects, at least in part because of the existence of multiple independent risk factors in this population. Successful treatment of hyperlipidemia in the general population, and in diabetic subjects in particular, is therefore of exceptional medical importance.

Hypertension (or high blood pressure) is a condition which occurs in the human
20 population as a secondary symptom to various other disorders such as renal artery stenosis, pheochromocytoma or endocrine disorders. However, hypertension is also evidenced in many patients in whom the causative agent or disorder is unknown. While such "essential" hypertension is often associated with disorders such as obesity, diabetes and hypertriglyceridemia, the relationship between these disorders has not
25 been elucidated. Additionally, many patients display the symptoms of high blood pressure in the complete absence of any other signs of disease or disorder.

It is known that hypertension can directly lead to heart failure, renal failure and stroke (brain hemorrhaging). These conditions are capable of causing short-term death in a patient. Hypertension can also contribute to the development of atherosclerosis
30 and coronary disease. These conditions gradually weaken a patient and can led to long-term death.

The exact cause of essential hypertension is unknown, though a number of factors are believed to contribute to the onset of the disease. Among such factors are

stress, uncontrolled emotions, unregulated hormone release (the renin, angiotensin, aldosterone system), excessive salt and water due to kidney malfunction, wall thickening and hypertrophy of the vasculature resulting in constricted blood vessels and genetic factors.

5 The treatment of essential hypertension has been undertaken bearing the foregoing factors in mind. Thus a broad range of beta-blockers, vasoconstrictors, angiotensin converting enzyme inhibitors and the like have been developed and marketed as antihypertensives. The treatment of hypertension utilizing these compounds has proven beneficial in the prevention of short-interval deaths such as
10 heart failure, renal failure and brain hemorrhaging. However, the development of atherosclerosis or heart disease due to hypertension over a long period of time remains a problem. This implies that although high blood pressure is being reduced, the underlying cause of essential hypertension is not responding to this treatment.

Hypertension has been associated with elevated blood insulin levels, a condition
15 known as hyperinsulinemia. Insulin, a peptide hormone whose primary actions are to promote glucose utilization, protein synthesis and the formation and storage of neutral lipids, also acts to promote vascular cell growth and increase renal sodium retention, among other things. These latter functions can be accomplished without affecting glucose levels and are known causes of hypertension. Peripheral vasculature growth,
20 for example, can cause constriction of peripheral capillaries; while sodium retention increases blood volume. Thus, the lowering of insulin levels in hyperinsulinemics can prevent abnormal vascular growth and renal sodium retention caused by high insulin levels and thereby alleviate hypertension.

Cardiac hypertrophy is a significant risk factor in the development of sudden
25 death, myocardial infarction, and congestive heart failure. These cardiac events are due, at least in part, to increased susceptibility to myocardial injury after ischemia and reperfusion which can occur in out-patient as well as perioperative settings. There is an unmet medical need to prevent or minimize adverse myocardial perioperative outcomes, particularly perioperative myocardial infarction. Both non-cardiac and cardiac
30 surgery are associated with substantial risks for myocardial infarction or death. Some 7 million patients undergoing non-cardiac surgery are considered to be at risk, with incidences of perioperative death and serious cardiac complications as high as 20-25% in some series. In addition, of the 400,000 patients undergoing coronary by-pass

surgery annually, perioperative myocardial infarction is estimated to occur in 5% and death in 1-2%. There is currently no drug therapy in this area which reduces damage to cardiac tissue from perioperative myocardial ischemia or enhances cardiac resistance to ischemic episodes. Such a therapy is anticipated to be life-saving and
5 reduce hospitalizations, enhance quality of life and reduce overall health care costs of high risk patients.

Hepatic glucose production is an important target for NIDDM therapy. The liver is the major regulator of plasma glucose levels in the post absorptive (fasted) state, and the rate of hepatic glucose production in NIDDM patients is significantly
10 elevated relative to normal individuals. Likewise, in the postprandial (fed) state, where the liver has a proportionately smaller role in the total plasma glucose supply, hepatic glucose production is abnormally high in NIDDM patients.

Glycogenolysis is an important target for interruption of hepatic glucose production. The liver produces glucose by glycogenolysis (breakdown of the
15 glucose polymer glycogen) and gluconeogenesis (synthesis of glucose from 2- and 3-carbon precursors). Several lines of evidence indicate that glycogenolysis may make an important contribution to hepatic glucose output in NIDDM. First, in normal post absorptive man, up to 75% of hepatic glucose production is estimated to result from glycogenolysis. Second, patients having liver glycogen storage diseases,
20 including Hers' disease (glycogen phosphorylase deficiency), display episodic hypoglycemia. These observations suggest that glycogenolysis may be a significant process for hepatic glucose production.

Glycogenolysis is catalyzed in liver, muscle, and brain by tissue-specific isoforms of the enzyme glycogen phosphorylase. This enzyme cleaves the glycogen
25 macromolecule to release glucose-1-phosphate and a new shortened glycogen macromolecule. Two types of glycogen phosphorylase inhibitors have been reported to date: glucose and glucose analogs [Martin, J.L. et al. Biochemistry 1991, 30, 10101] and caffeine and other purine analogs [Kasvinsky, P.J. et al. J. Biol. Chem. 1978, 253, 3343-3351 and 9102-9106]. These compounds, and
30 glycogen phosphorylase inhibitors in general, have been postulated to be of potential use for the treatment of NIDDM by decreasing hepatic glucose production and lowering glycemia. [Blundell, T.B. et al. Diabetologia 1992, 35, Suppl. 2, 569-576 and Martin et al. Biochemistry 1991, 30, 10101].

-5-

The mechanism(s) responsible for the myocardial injury observed after ischemia and reperfusion is not fully understood. It has been reported (M. F. Allard, et al. Am. J. Physiol. 267, H66-H74, 1994) that "pre ischemic glycogen reduction...is associated with improved post ischemic left ventricular functional recovery in hypertrophied rat hearts".

Thus, although there are a variety of hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemia, hyperlipidemia, atherosclerosis and myocardial ischemia therapies there is a continuing need and a continuing search in this field of art for alternative therapies.

10


Summary of the Invention

This invention is directed to a glycogen phosphorylase inhibitor compound of Formula I useful for the treatment of diabetes, hyperglycemia, hypercholesterolemia, hyperinsulinemia, hypertension, hyperlipidemia, atherosclerosis and myocardial ischemia.

15

The compounds of this invention have the Formula I

20



Formula I

and the pharmaceutically acceptable salts and prodrugs thereof

25 wherein

the dotted line (—) is an optional bond;

A is $-C(H)=$, $-C((C_1-C_4)alkyl)=$, $-C(halo)=$ or $-N=$, when the dotted line (—) is a bond, or A is methylene or $-CH((C_1-C_4)alkyl)-$, when the dotted line (—) is not a bond;

30

R_1 , R_{10} or R_{11} are each independently H, halo, cyano, 4-, 6-, or 7-nitro, $(C_1-C_4)alkyl$, $(C_1-C_4)alkoxy$, fluoromethyl, difluoromethyl or trifluoromethyl;

R_2 is H;

R_3 is H or $(C_1-C_5)alkyl$;

-6-

R_4 is H, methyl, ethyl, n-propyl, hydroxy(C_1 - C_3)alkyl, (C_1 - C_3)alkoxy(C_1 - C_3)alkyl, phenyl(C_1 - C_4)alkyl, phenylhydroxy(C_1 - C_4)alkyl, (phenyl)((C_1 - C_4)-alkoxy)(C_1 - C_4)alkyl, thien-2- or -3-yl(C_1 - C_4)alkyl or fur-2- or -3-yl(C_1 - C_4)alkyl wherein said R_4 rings are mono-, di- or tri-substituted independently on carbon with H, halo, (C_1 - C_4)alkyl, (C_1 - C_4)alkoxy, trifluoromethyl, hydroxy, amino, cyano or 4,5-dihydro-1H-imidazol-2-yl; or

R_4 is pyrid-2-, -3- or -4-yl(C_1 - C_4)alkyl, thiazol-2-, -4- or -5-yl(C_1 - C_4)alkyl, imidazol-2-, -4- or -5-yl(C_1 - C_4)alkyl, pyrrol-2- or -3-yl(C_1 - C_4)alkyl, oxazol-2-, -4- or -5-yl(C_1 - C_4)alkyl, pyrazol-3-, -4- or -5-yl(C_1 - C_4)alkyl, isoxazol-3-, -4- or -5-yl(C_1 - C_4)alkyl, isothiazol-3-, -4- or -5-yl(C_1 - C_4)alkyl, pyridazin-3- or -4-yl(C_1 - C_4)alkyl, pyrimidin-2-, -4-, -5- or -6-yl(C_1 - C_4)alkyl, pyrazin-2- or -3-yl(C_1 - C_4)alkyl, 1,3,5-triazin-2-yl(C_1 - C_4)alkyl or indol-2-(C_1 - C_4)alkyl, wherein said preceding R_4 heterocycles are optionally mono- or di-substituted independently with halo, trifluoromethyl, (C_1 - C_4)alkyl, (C_1 - C_4)alkoxy, amino, hydroxy or cyano and said substituents are bonded to carbon; or

R_4 is R_{15} -carbonyloxymethyl, wherein said R_{15} is phenyl, thiazolyl, imidazolyl, 1H-indolyl, furyl, pyrrolyl, oxazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl or 1,3,5-triazinyl and wherein said preceding R_{15} rings are optionally mono- or di-substituted independently with halo, amino, hydroxy, (C_1 - C_4)alkyl, (C_1 - C_4)alkoxy or trifluoromethyl and said mono- or di-substituents are bonded to carbon;

R_5 is H, methyl, ethyl, n-propyl, hydroxymethyl or hydroxyethyl;

R_6 is carboxy, (C_1 - C_8)alkoxycarbonyl, benzyloxycarbonyl, $C(O)NR_8R_9$ or $C(O)R_{12}$ wherein

R_8 is H, (C_1 - C_8)alkyl, cyclo(C_3 - C_8)alkyl, cyclo(C_3 - C_8)alkyl(C_1 - C_5)alkyl, hydroxy or (C_1 - C_8)alkoxy; and

R_9 is H, cyclo(C_3 - C_8)alkyl, cyclo(C_3 - C_8)alkyl(C_1 - C_5)alkyl, cyclo(C_4 - C_7)alkenyl, cyclo(C_3 - C_7)alkyl(C_1 - C_5)alkoxy, cyclo(C_3 - C_7)alkyloxy, hydroxy, methylene-perfluorinated(C_1 - C_8)alkyl, phenyl, or a heterocycle wherein said heterocycle is pyridyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, pyridinyl, piperidinyl, morpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, thiochromanyl or

-7-

tetrahydrobenzothiazolyl wherein said heterocycle rings are carbon-nitrogen linked;

or

R_9 is (C_1-C_6) alkyl or (C_1-C_6) alkoxy wherein said (C_1-C_6) alkyl or (C_1-C_6) alkoxy is optionally monosubstituted with cyclo (C_4-C_7) alken-1-yl, phenyl, thienyl, pyridyl, 5 furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, piperidinyl, morpholinyl, thiomorpholinyl, 1-oxothiomorpholinyl, 1,1-dioxothiomorpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl or indolyl and wherein said (C_1-C_6) alkyl or (C_1-C_6) alkoxy are optionally additionally independently mono- or di- 10 substituted with halo, hydroxy, (C_1-C_5) alkoxy, amino, mono-N- or di-N,N- (C_1-C_5) alkylamino, cyano, carboxy, or (C_1-C_4) alkoxycarbonyl; and

wherein the R_9 rings are optionally mono- or di-substituted independently on carbon with halo, (C_1-C_4) alkyl, (C_1-C_4) alkoxy, hydroxy, hydroxy (C_1-C_4) alkyl, amino (C_1-C_4) alkyl, mono-N- or di-N,N- (C_1-C_4) alkylamino (C_1-C_4) alkyl, (C_1-C_4) alkoxy (C_1-C_4) alkyl, amino, mono-N- or di-N,N- (C_1-C_4) alkylamino, cyano, carboxy, 15 (C_1-C_5) alkoxycarbonyl, carbamoyl, formyl or trifluoromethyl and said R_9 rings may optionally be additionally mono- or di-substituted independently with (C_1-C_5) alkyl or halo;

with the proviso that no quaternized nitrogen on any R_9 heterocycle is 20 included;

R_{12} is morpholino, thiomorpholino, 1-oxothiomorpholino, 1,1-dioxothiomorpholino, thiazolidin-3-yl, 1-oxothiazolidin-3-yl, 1,1-dioxothiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, piperazin-4-yl, azetidin-1-yl, 1,2-oxazinan-2-yl, pyrazolidin-1-yl, isoxazolidin-2-yl, isothiazolidin-2-yl, 1,2-oxazetidin-2-yl, 25 oxazolidin-3-yl, 3,4-dihydroisoquinolin-2-yl, 1,3-dihydroisoindol-2-yl, 3,4-dihydro-2H-quinol-1-yl, 2,3-dihydro-benzo[1,4]oxazin-4-yl, 2,3-dihydro-benzo[1,4]thiazine-4-yl, 3,4-dihydro-2H-quinoxalin-1-yl, 3,4-dihydro-benzo[c][1,2]oxazin-1-yl, 1,4-dihydro-benzo[d][1,2]oxazin-3-yl, 3,4-dihydro-benzo[e][1,2]oxazin-2-yl, 3H-benzo[d]isoxazol-2-yl, 3H-benzo[c]isoxazol-1-yl or azepan-1-yl,

wherein said R_{12} rings are optionally mono-, di- or tri-substituted independently with halo, (C_1-C_5) alkyl, (C_1-C_5) alkoxy, hydroxy, amino, mono-N- or di-N,N- (C_1-C_5) alkylamino, formyl, carboxy, carbamoyl, mono-N- or di-N,N- (C_1-C_5) alkylcarbamoyl, (C_1-C_6) alkoxy (C_1-C_3) alkoxy, (C_1-C_5) alkoxycarbonyl, 30

benzyloxycarbonyl, (C₁-C₅)alkoxycarbonyl(C₁-C₅)alkyl, (C₁-C₄)alkoxycarbonylamino, carboxy(C₁-C₅)alkyl, carbamoyl(C₁-C₅)alkyl, mono-N- or di-N,N-(C₁-C₅)alkylcarbamoyl(C₁-C₅)alkyl, hydroxy(C₁-C₅)alkyl, (C₁-C₄)alkoxy(C₁-C₄)alkyl, amino(C₁-C₄)alkyl, mono-N- or di-N,N-(C₁-C₄)alkylamino(C₁-C₄)alkyl, oxo,

- 5 hydroxyimino or (C₁-C₆)alkoxyimino and wherein no more than two substituents are selected from oxo, hydroxyimino or (C₁-C₆)alkoxyimino and oxo, hydroxyimino or (C₁-C₆)alkoxyimino are on nonaromatic carbon; and

wherein said R₁₂ rings are optionally additionally mono- or di-substituted independently with (C₁-C₅)alkyl or halo;

- 10 with the proviso that when R₆ is (C₁-C₅)alkoxycarbonyl or benzyloxycarbonyl then R₁ is 5-halo, 5-(C₁-C₄)alkyl or 5-cyano and R₄ is (phenyl)(hydroxy)(C₁-C₄)alkyl, (phenyl)((C₁-C₄)alkoxy)(C₁-C₄)alkyl, hydroxymethyl or Ar(C₁-C₂)alkyl, wherein Ar is thien-2- or -3-yl, fur-2- or -3-yl or phenyl wherein said Ar is optionally mono- or di-substituted independently with halo; with the provisos that when R₄ is benzyl and R₅ is methyl, R₁₂ is not 4-hydroxy-piperidin-1-yl or when R₄ is benzyl and R₅ is methyl R₆ is not C(O)N(CH₃)₂;

with the proviso that when R₁ and R₁₀ and R₁₁ are H, R₄ is not imidazol-4-ylmethyl, 2-phenylethyl or 2-hydroxy-2-phenylethyl;

- with the proviso that when both R₈ and R₉ are n-pentyl, none of R₁ is 5-chloro, 5-bromo, 5-cyano, 5(C₁-C₅)alkyl, 5(C₁-C₅)alkoxy or trifluoromethyl;

with the proviso that when R₁₂ is 3,4-dihydroisoquinol-2-yl, said 3,4-dihydroisoquinol-2-yl is not substituted with carboxy((C₁-C₄)alkyl);

- with the proviso that when R₈ is H and R₉ is (C₁-C₆)alkyl, R₉ is not substituted with carboxy or (C₁-C₄)alkoxycarbonyl on the carbon which is attached to the nitrogen atom N of NHR₉; and

with the proviso that when R₈ is carboxy and R₁₁, R₁₀, R₁₁ and R₅ are all H, then R₄ is not benzyl, H, (phenyl)(hydroxy)methyl, methyl, ethyl or n-propyl.

A first group of preferred compounds of Formula I consists of those compounds wherein

- 30 R₁ is 5-H, 5-halo, 5-methyl, 5-cyano or 5-trifluoromethyl;
R₁₀ and R₁₁ are each independently H or halo;
A is -C(H)=;
R₂ and R₃ are H;

R_4 is H, methyl, phenyl(C_1 - C_2)alkyl, wherein said phenyl groups are mono- or di-substituted independently with H, halo, (C_1 - C_4)alkyl, (C_1 - C_4)alkoxy, trifluoromethyl, hydroxy, amino or cyano and wherein said R_4 groups are optionally additionally mono-substituted with halo; or

- 5 R_4 is thien-2- or -3-yl(C_1 - C_2)alkyl, pyrid-2-, -3- or -4-yl(C_1 - C_2)alkyl, thiazol-2-, -4- or -5-yl(C_1 - C_2)alkyl, imidazol-2-, -4- or -5-yl(C_1 - C_2)alkyl, fur-2- or -3-yl(C_1 - C_2)alkyl, pyrrol-2- or -3-yl(C_1 - C_2)alkyl, oxazol-2-, -4- or -5-yl(C_1 - C_2)alkyl, pyrazol-3-, -4- or -5-yl(C_1 - C_2)alkyl, isoxazol-3-, -4- or -5-yl(C_1 - C_2)alkyl, isothiazol-3-, -4- or -5-yl(C_1 - C_2)alkyl, pyridazin-3- or -4-yl(C_1 - C_2)alkyl, pyrimidin-2-, -4-, -5- or -6-yl(C_1 - C_2)alkyl, pyrazin-2-
10 or -3-yl(C_1 - C_2)alkyl or 1,3,5-triazin-2-yl(C_1 - C_2)alkyl wherein said preceding R_4 heterocycles are optionally mono- or di-substituted independently with halo, trifluoromethyl, (C_1 - C_4)alkyl, (C_1 - C_4)alkoxy, amino or hydroxy and said mono- or di-substituents are bonded to carbon;

R_5 is H; and

- 15 R_6 is $C(O)NR_8R_9$ or $C(O)R_{12}$.

Within the above first group of preferred compounds of Formula I is a first group of especially preferred compounds wherein

- R_4 is H, phenyl(C_1 - C_2)alkyl, thien-2- or -3-yl(C_1 - C_2)alkyl, fur-2- or -3-yl(C_1 - C_2)alkyl wherein said R_4 rings are mono- or di-substituted independently with H or
20 fluoro;

R_6 is $C(O)R_{12}$; and

- R_{12} is morpholino, thiomorpholino, 1-oxothiomorpholino, 1,1-dioxothiomorpholino, thiazolidin-3-yl, 1-oxothiazolidin-3-yl, 1,1-dioxothiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, piperazin-4-yl, azetidin-1-yl, 1,2-oxazin-
25 2-yl, isoxazolidin-2-yl, isothiazolidin-2-yl, 1,2-oxazetidin-2-yl, oxazolidin-3-yl, 1,3-dihydroisoindol-2-yl, or azepan-1-yl,

wherein said R_{12} rings are optionally mono- or di-substituted independently with halo, (C_1 - C_5)alkyl, (C_1 - C_5)alkoxy, hydroxy, amino, mono-N- or di-N,N-(C_1 - C_5)alkylamino, formyl, carboxy, carbamoyl, mono-N- or di-N,N-(C_1 -

- 30 C_5)alkylcarbamoyl, (C_1 - C_5)alkoxycarbonyl, hydroxy(C_1 - C_5)alkyl, amino(C_1 - C_4)alkyl, mono-N- or di-N,N-(C_1 - C_4)alkylamino(C_1 - C_4)alkyl, oxo, hydroxyimino or (C_1 - C_6)alkoxyimino with the proviso that only the R_{12} heterocycles thiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, piperazin-4-yl, azetidin-1-yl, 1,2-oxazin-

-10-

2-yl, isoxazolidin-2-yl, or oxazolidin-3-yl are optionally mono- or di-substituted with oxo, hydroxyimino, or (C₁-C₆)alkoxyimino; and

wherein said R₁₂ rings are optionally additionally mono- or di-substituted independently with (C₁-C₅)alkyl.

5 Within the above group of especially preferred compounds are the compounds

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxyimino-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [2-(cis-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

10 5-Chloro-1H-indole-2-carboxylic acid [2-((3S,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(cis-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

15 5-Chloro-1H-indole-2-carboxylic acid [2-(1,1-dioxo-thiazolidin-3-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid (2-oxo-2-thiazolidin-3-yl-ethyl)-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-(4-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide,

20 5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3RS)-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [2-oxo-2-((1RS)-oxo-1-thiazolidin-3-yl)-ethyl]-amide,

25 5-Chloro-1H-indole-2-carboxylic acid [(1S)-(2-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3S,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide,

30 5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxyimino-azetidin-1-yl)-2-oxo-ethyl]-amide or

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(4-hydroxyimino-piperidin-1-yl)-2-oxo-ethyl]-amide.

-11-

Within the above group of especially preferred compounds is a first group of particularly preferred compounds wherein

R_4 is H; and

R_{12} is thiazolidin-3-yl, 1-oxo-thiazolidin-3-yl, 1,1-dioxo-thiazolidin-3-yl or
5 oxazolidin-3-yl or said R_{12} substituents optionally mono- or di-substituted independently with carboxy, (C_1-C_5) alkoxycarbonyl, hydroxy (C_1-C_3) alkyl, amino (C_1-C_3) alkyl, mono-N- or di-N,N- (C_1-C_3) alkylamino (C_1-C_3) alkyl or

R_{12} is mono- or di-substituted pyrrolidin-1-yl wherein said substituents are independently carboxy, (C_1-C_5) alkoxycarbonyl, (C_1-C_5) alkoxy, hydroxy, hydroxy (C_1-C_3) alkyl, amino, amino (C_1-C_3) alkyl, mono-N- or di-N,N- (C_1-C_3) alkylamino (C_1-C_3) alkyl
10 or mono-N- or di-N,N- (C_1-C_4) alkylamino; and

the R_{12} rings are optionally additionally independently disubstituted with (C_1-C_5) alkyl.

Preferred compounds within the immediately preceding group of particularly
15 preferred compounds are compounds wherein

a. R_1 is 5-chloro;

R_{10} and R_{11} are H; and

R_{12} is cis-3,4-dihydroxy-pyrrolidin-1-yl;

b. R_1 is 5-chloro;

20 R_{10} and R_{11} are H; and

R_{12} is (3S,4S)-dihydroxy-pyrrolidin-1-yl;

c. R_1 is 5-chloro;

R_{10} and R_{11} are H; and

R_{12} is 1,1-dioxo-thiazolidin-3-yl;

25 d. R_1 is 5-chloro;

R_{10} and R_{11} are H; and

R_{12} is thiazolidin-3-yl; and

e. R_1 is 5-chloro;

R_{10} and R_{11} are H; and

30 R_{12} is 1-oxo-thiazolidin-3-yl.

Within the above group of especially preferred compounds is a second group of particularly preferred compounds wherein

-12-

R_4 is phenylmethyl, thien-2- or -3-ylmethyl wherein said R_4 rings are optionally mono- or di-substituted with fluoro; and

R_{12} is thiazolidin-3-yl, 1-oxo-thiazolidin-3-yl, 1,1-dioxo-thiazolidin-3-yl or oxazolidin-3-yl or said R_{12} substituents optionally mono- or di-substituted

5 independently with carboxy or (C_1-C_5) alkoxycarbonyl, hydroxy (C_1-C_3) alkyl, amino (C_1-C_3) alkyl or mono-N- or di-N,N- (C_1-C_3) alkylamino (C_1-C_3) alkyl

or R_{12} is mono- or di-substituted azetidin-1-yl or mono- or di-substituted pyrrolidin-1-yl or mono- or di-substituted piperidin-1-yl wherein said substituents are independently carboxy, (C_1-C_5) alkoxycarbonyl, hydroxy (C_1-C_3) alkyl, amino $(C_1-$

10 $C_3)$ alkyl, mono-N- or di-N,N- (C_1-C_3) alkylamino (C_1-C_3) alkyl, hydroxy, (C_1-C_5) alkoxy, amino, mono-N- or di-N,N- (C_1-C_5) alkylamino, oxo, hydroxyimino or (C_1-C_5) alkoxyimino; and

the R_{12} rings are optionally additionally mono- or di-substituted independently with (C_1-C_5) alkyl.

15 Preferred compounds within the immediately preceding group of particularly preferred compounds are compounds wherein

a. R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is 4-fluorobenzyl;

20 R_{12} is 4-hydroxypiperidin-1-yl; and
the stereochemistry of carbon (a) is (S);

b. R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

25 R_{12} is 3-hydroxypiperidin-1-yl; and
the stereochemistry of carbon (a) is (S);

c. R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

30 R_{12} is cis-3,4-dihydroxy-pyrrolidin-1-yl; and
the stereochemistry of carbon (a) is S;

d. R_1 is 5-chloro;

R_{10} and R_{11} are H; R_4 is benzyl;

-13-

- R_{12} is 3-hydroxyimino-pyrrolidin-1-yl; and
the stereochemistry of carbon (a) is (S);
- e. R_1 is 5-chloro;
 R_{10} and R_{11} are H;
5 R_4 is 2-fluorobenzyl;
 R_{12} is 4-hydroxypiperidin-1-yl; and
the stereochemistry of carbon (a) is (S);
- f. R_1 is 5-chloro;
 R_{10} and R_{11} are H;
10 R_4 is benzyl;
 R_{12} is (3S,4S)-dihydroxy-pyrrolidin-1-yl; and
the stereochemistry of carbon (a) is (S);
- g. R_1 is 5-chloro;
 R_{10} and R_{11} are H;
15 R_4 is benzyl;
 R_{12} is 3-hydroxy-azetidin-1-yl; and
the stereochemistry of carbon (a) is (S);
- h. R_1 is 5-chloro;
 R_{10} and R_{11} are H;
20 R_4 is benzyl;
 R_{12} is 3-hydroxyimino-azetidin-1-yl; and
the stereochemistry of carbon (a) is (S); and
- i. R_1 is 5-chloro;
 R_{10} and R_{11} are H;
25 R_4 is benzyl;
 R_{12} is 4-hydroxyimino-piperidin-1-yl; and
the stereochemistry of carbon (a) is (S).

A second group of especially preferred compounds within the first group of preferred compounds are the compounds wherein

- 30 R_4 is H, phenyl(C_1 - C_2)alkyl, thien-2- or -3-yl(C_1 - C_2)alkyl, fur-2- or -3-yl(C_1 - C_2)alkyl wherein said R_4 rings are mono- or di-substituted independently with H or fluoro;

R_6 is $C(O)NR_8R_9$; and

-14-

R_8 is H, (C_1-C_5) alkyl, hydroxy or (C_1-C_4) alkoxy; and

R_9 is H, cyclo (C_4-C_6) alkyl, cyclo (C_3-C_6) alkyl (C_1-C_5) alkyl, methylene-perfluorinated (C_1-C_3) alkyl, pyridyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, piperidinyl, benzothiazolyl or thiochromanyl; or

- 5 R_9 is (C_1-C_5) alkyl wherein said (C_1-C_5) alkyl is optionally substituted with cyclo (C_4-C_6) alkenyl, phenyl, thienyl, pyridyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, piperidinyl, morpholinyl, thiomorpholinyl, 1-oxothiomorpholinyl, or 1,1-dioxothiomorpholinyl and wherein said (C_1-C_5) alkyl or (C_1-C_4) alkoxy is optionally additionally independently mono- or di-substituted with halo, hydroxy, $(C_1-$
 10 $C_5)$ alkoxy, amino, mono-N- or di-N,N- (C_1-C_5) alkylamino, cyano, carboxy, or (C_1-C_4) alkoxycarbonyl; and

wherein the R_8 rings are optionally mono- or di-substituted independently on carbon with halo, (C_1-C_4) alkyl, (C_1-C_4) alkoxy, hydroxy, amino, mono-N- or di-N,N- (C_1-C_4) alkylamino, carbamoyl, (C_1-C_5) alkoxycarbonyl or carbamoyl.

- 15 Within the immediately preceding second group of especially preferred compounds are the compounds wherein

a. R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

20 R_8 is methyl; and

R_9 is 3-(dimethylamino)propyl;

b. the stereochemistry of carbon (a) is (S);

R_1 is 5-chloro;

R_{10} and R_{11} are H;

25 R_4 is benzyl;

R_8 is methyl; and

R_9 is 3-pyridyl;

c. the stereochemistry of carbon (a) is (S);

R_1 is 5-chloro;

30 R_{10} and R_{11} are H;

R_4 is benzyl;

R_8 is methyl; and

R_9 is 2-hydroxyethyl; and

-15-

d. the stereochemistry of carbon (a) is (S);

R_1 is 5-fluoro;

R_{10} and R_{11} are H;

R_4 is 4-fluorophenylmethyl;

5 R_8 is methyl; and

R_9 is 2-morpholinoethyl.

A third group of especially preferred compounds within the first group of preferred compounds are the compounds wherein

R_4 is H, phenyl(C_1 - C_2)alkyl, thien-2- or -3-yl(C_1 - C_2)alkyl, fur-2- or -3-yl(C_1 -
10 C_2)alkyl wherein said R_4 rings are mono- or di-substituted independently with H or fluoro;

R_6 is $C(O)NR_8R_9$; and

R_8 is H, (C_1 - C_5)alkyl, hydroxy or (C_1 - C_4)alkoxy; and

R_9 is (C_1 - C_4)alkoxy wherein said (C_1 - C_4)alkoxy is optionally substituted with
15 cyclo(C_4 - C_6)alkenyl, phenyl, thienyl, pyridyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, piperidinyl, morpholinyl, thiomorpholinyl, 1-oxothiomorpholinyl, or 1,1-dioxothiomorpholinyl and wherein said (C_1 - C_5)alkyl or (C_1 - C_4)alkoxy is optionally additionally independently mono- or di-substituted with halo, hydroxy, (C_1 - C_5)alkoxy, amino, mono-N- or di-N,N-(C_1 - C_5)alkylamino, cyano, carboxy, or (C_1 -
20 C_4)alkoxycarbonyl; and

wherein the R_9 rings are optionally mono- or di-substituted independently on carbon with halo, (C_1 - C_4)alkyl, (C_1 - C_4)alkoxy, hydroxy, amino, mono-N- or di-N,N-(C_1 - C_4)alkylamino, carbamoyl, (C_1 - C_5)alkoxycarbonyl or carbamoyl.

Within the immediately preceding third group of especially preferred
25 compounds are the compounds wherein

a. R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

R_8 is methyl; and

30 R_9 is 2-hydroxyethoxy;

b. the stereochemistry of carbon (a) is (S);

R_1 is 5-chloro;

R_{10} and R_{11} are H;

-16-

R_4 is 4-fluorophenylmethyl;

R_8 is methyl; and

R_9 is methoxy;

c. the stereochemistry of carbon (a) is (S);

5 R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

R_8 is methyl; and

R_9 is methoxy;

10 A second group of preferred compounds of Formula I are those compounds wherein

R_1 is 5-halo, 5-methyl, 5-cyano or trifluoromethyl;

R_{10} and R_{11} are each independently H or halo;

A is $-C(H)=$;

15 R_2 and R_3 are H;

R_4 is H, phenyl(C_1-C_2)alkyl, thien-2- or -3-yl(C_1-C_2)alkyl, fur-2- or -3-yl(C_1-C_2)alkyl wherein said rings are mono- or di-substituted independently with H or fluoro;

R_5 is H; and

20 R_8 is (C_1-C_5)alkoxycarbonyl.

A third group of preferred compounds of Formula I are those compounds wherein

R_1 is 5-halo, 5-methyl, 5-cyano or trifluoromethyl;

R_{10} and R_{11} are each independently H or halo;

25 A is $-C(H)=$;

R_2 and R_3 are H;

R_4 is H, methyl or phenyl(C_1-C_2)alkyl, wherein said phenyl groups are mono- or di-substituted independently with H, halo, (C_1-C_4)alkyl, (C_1-C_4)alkoxy, trifluoromethyl, hydroxy, amino or cyano and wherein said phenyl groups are

30 additionally mono- or di-substituted independently H or halo; or

R_4 is thien-2- or -3-yl(C_1-C_2)alkyl, pyrid-2-, -3- or -4-yl(C_1-C_2)alkyl, thiazol-2-, -4- or -5-yl(C_1-C_2)alkyl, imidazol-2-, -4- or -5-yl(C_1-C_2)alkyl, fur-2- or -3-yl(C_1-C_2)alkyl, pyrrol-2- or -3-yl(C_1-C_2)alkyl, oxazol-2-, -4- or -5-yl(C_1-C_2)alkyl, pyrazol-3-, -4- or -5-

-17-

yl(C₁-C₂)alkyl, isoxazol-3-, -4- or -5-yl(C₁-C₂)alkyl, isothiazol-3-, -4- or -5-yl(C₁-C₂)alkyl, pyridazin-3- or -4-yl(C₁-C₂)alkyl, pyrimidin-2-, -4-, -5- or -6-yl(C₁-C₂)alkyl, pyrazin-2- or -3-yl(C₁-C₂)alkyl or 1,3,5-triazin-2-yl(C₁-C₂)alkyl wherein said preceding R₄ heterocycles are optionally mono- or di-substituted independently with halo, trifluoromethyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, amino or hydroxy and said mono- or di-substituents are bonded to carbon;

R₅ is H; and

R₆ is carboxy.

Within the third group of preferred compounds is a first group of especially preferred compounds wherein


R₁₀ and R₁₁ are H; and

R₄ is H.

Particularly preferred within the immediately preceding especially preferred group is a compound wherein

R₁ is 5-chloro.

Another aspect of this invention is directed to intermediates useful for making some of the compounds of Formula I. The intermediates have the Formula QZ



Formula QZ

wherein

R₅ is H;

R₄ is H, phenylmethyl, thien-2- or -3-ylmethyl, fur-2- or -3-ylmethyl wherein said rings are optionally mono- or di-substituted with fluoro; and

R₁₂ is thiazolidin-3-yl, 1-oxothiazolidin-3-yl, 1,1-dioxothiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, azetidin-1-yl, 1,2-oxazinan-2-yl, isoxazolidin-2-yl, isothiazolidin-2-yl, 1,2-oxazetidin-2-yl or oxazolidin-3-yl,

wherein said R₁₂ rings are optionally mono- or di-substituted independently with halo, (C₁-C₅)alkyl, (C₁-C₅)alkoxy, hydroxy, amino, mono-N- or di-N,N-(C₁-C₅)alkylamino, formyl, carboxy, carbamoyl, mono-N- or di-N,N-(C₁-

-18-

C₅)alkylcarbamoyl, (C₁-C₅)alkoxycarbonyl, hydroxy(C₁-C₅)alkyl, amino(C₁-C₄)alkyl, mono-N- or di-N,N-(C₁-C₄)alkylamino(C₁-C₄)alkyl, oxo, hydroxyimino or (C₁-C₆)alkoxyimino with the proviso that only the R₁₂ heterocycles thiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, azetidin-1-yl, 1,2-oxazinan-2-yl, isoxazolidin-2-yl, or
5 oxazolidin-3-yl are optionally mono- or di-substituted independently with oxo, hydroxyimino, or (C₁-C₆)alkoxyimino; and

wherein said R₁₂ rings are optionally additionally mono- or di-substituted independently with (C₁-C₅)alkyl and

with the proviso that R₁₂ is not 2-carboxy-4-hydroxy- pyrrolidin-1-yl, 2-((C₁-
10 C₅)alkoxycarbonyl)-4-hydroxy-pyrrolidin-1-yl, 2-carboxy-piperidin-1-yl or 2-((C₁-C₅)alkoxycarbonyl)-piperidin-1-yl.

Particular compounds within the above group of intermediates are the compounds wherein

- a. R₄ is H; and
15 R₁₂ is thiazolidin-3-yl;
- b. R₄ is H; and
R₁₂ is 1,1-dioxo-thiazolidin-3-yl; and
- c. R₄ is H; and
R₁₂ is 1-oxo-thiazolidin-3-yl.

20 A first group of preferred compounds of Formula QZ are those compounds wherein

R₄ is phenylmethyl, said phenyl optionally mono- or di-substituted with fluoro; and

R₁₂ is 3-mono-substituted azetidin-1-yl, 3-mono- or 3,4-disubstituted
25 pyrrolidin-1-yl, 3-, 4-, or 5- mono- or di-substituted piperidin-1-yl, thiazolidin-3-yl, 1-oxo-thiazolidin-3-yl or 1,1-dioxothiazolidin-3-yl wherein said pyrrolidin-1-yl or piperidin-1-yl are mono- or di-substituted independently with hydroxy, oxo, hydroxyimino, amino, mono-N- or di-N,N-(C₁-C₄)alkylamino, (C₁-C₅)alkoxycarbonyl or carboxy

30 and said R₁₂ rings are optionally additionally mono- or di-substituted independently with (C₁-C₄)alkyl.

Particular compounds within the above immediately preceding group of preferred compounds are the compounds wherein

-19-

- a. R_4 is benzyl;
 $R_{1,2}$ is 3-hydroxypyrrolidin-3-yl; and
the stereochemistry of carbon (a) is (S);
- b. R_4 is benzyl;
5 $R_{1,2}$ is 3-hydroxyazetidin-1-yl; and
the stereochemistry of carbon (a) is (S);
- c. R_4 is benzyl;
 $R_{1,2}$ is 3,4-dihydroxypyrrolidin-1-yl; and
the stereochemistry of carbon (a) is (S);
- 10 d. R_4 is benzyl;
 $R_{1,2}$ is 4-hydroxypiperidin-1-yl; and
the stereochemistry of carbon (a) is (S);
- e. R_4 is 4-fluorophenylmethyl;
 $R_{1,2}$ is 4-hydroxypiperidin-1-yl; and
15 the stereochemistry of carbon (a) is (S); and
- f. R_4 is benzyl;
 $R_{1,2}$ is 4-hydroxyiminoazetidin-1-yl; and
the stereochemistry of carbon (a) is (S).

Yet another aspect of this invention is directed to a method for treating a
20 glycogen phosphorylase dependent disease or condition in a mammal by
administering to a mammal suffering from a glycogen phosphorylase dependent
disease or condition a glycogen phosphorylase dependent disease or condition
treating amount of a Formula I compound.

Yet another aspect of this invention is directed to a method for treating
25 hyperglycemia in a mammal by administering to a mammal suffering from
hyperglycemia a hyperglycemia treating amount of a Formula I compound.

Yet another aspect of this invention is directed to a method for treating
diabetes in a mammal by administering to a mammal suffering from diabetes a
diabetes treating amount of a Formula I compound. Included in the treatment of
30 diabetes is the prevention or attenuation of long term complications such as
neuropathy, nephropathy, retinopathy or cataracts.

Yet another aspect of this invention is directed to a method for treating
hypercholesterolemia in a mammal by administering to a mammal suffering from

-20-

hypercholesterolemia a hypercholesterolemia treating amount of a Formula I compound.

Yet another aspect of this invention is directed to a method for treating atherosclerosis in a mammal by administering to a mammal suffering from
5 atherosclerosis an atherosclerosis treating amount of a Formula I compound.

Yet another aspect of this invention is directed to a method for treating hyperinsulinemia in a mammal by administering to a mammal suffering from hyperinsulinemia a hyperinsulinemia treating amount of a Formula I compound.

Yet another aspect of this invention is directed to a method for treating
10 hypertension in a mammal by administering to a mammal suffering from hypertension a hypertension treating amount of a Formula I compound.

Yet another aspect of this invention is directed to a method for treating hyperlipidemia in a mammal by administering to a mammal suffering from hyperlipidemia a hyperlipidemia treating amount of a Formula I compound.

15 Yet another aspect of this invention is directed to a method for preventing a myocardial ischemic injury in a mammal by administering to a mammal at risk for perioperative myocardial ischemic injury a perioperative myocardial ischemic injury preventing amount of a Formula I compound.

Yet another aspect of this invention is directed to a method for preventing a
20 myocardial ischemic injury in a mammal by administering to a mammal at risk for perioperative myocardial ischemic injury a perioperative myocardial ischemic injury preventing amount of a glycogen phosphorylase inhibitor.

This invention is also directed to pharmaceutical compositions which comprise a therapeutically effective amount of a compound of Formula I and a
25 pharmaceutically acceptable carrier.

Preferred compositions include pharmaceutical compositions for the treatment of glycogen phosphorylase dependent diseases or conditions in mammals which comprise a glycogen phosphorylase dependent disease or condition treating amount of a compound of Formula I and a pharmaceutically acceptable carrier.

30 Another aspect of this invention is directed to pharmaceutical compositions for the treatment of diabetes which comprise a therapeutically effective amount of a glycogen phosphorylase inhibitor;

-21-

- one or more antidiabetic agents such as insulin and insulin analogs (e.g. LysPro insulin); GLP-1 (7-37) (insulinotropin) and GLP-1 (7-36)-NH₂; Sulfonylureas and Analogs: chlorpropamide, glibenclamide, tolbutamide, tolazamide, acetohexamide, glypizide®, glimepiride, repaglinide, meglitinide; Biguanides:
- 5 metformin, phenformin, buformin; α 2-Antagonists and Imidazolines: midaglizole, isaglidole, deriglidole, idazoxan, efaroan, fluparoxan; Other insulin secretagogues: linoglinide, A-4166; Glitazones: ciglitazone, pioglitazone, englitazone, troglitazone, darglitazone, BRL49653; Fatty Acid Oxidation Inhibitors: clomoxir, etomoxir; α -
- 10 Glucosidase inhibitors: acarbose, miglitol, emiglitate, voglibose, MDL-25,637, camiglibose, MDL-73,945; β -Agonists: BRL 35135, BRL 37344, Ro 16-8714, ICI D7114, CL 316,243; Phosphodiesterase Inhibitors: L-386,398; Lipid-lowering Agents: benfluorex; Antiobesity Agents: fenfluramine; Vanadate and vanadium complexes (e.g. naglivan®) and peroxovanadium complexes; Amylin Antagonists; Glucagon Antagonists; Gluconeogenesis Inhibitors; Somatostatin Analogs; Antilipolytic Agents:
- 15 nicotinic acid, acipimox, WAG 994; and

optionally a pharmaceutically acceptable carrier.

Preferred pharmaceutical compositions within the immediately preceding group are those compositions wherein the glycogen phosphorylase inhibitor is a compound of Formula I.

- 20 Another aspect of this invention is a method of treating diabetes in a mammal with the above described combination compositions.

Glycogen phosphorylase dependent diseases or conditions refers to disorders which are mediated, initiated or maintained, in whole or in part, by the cleavage of the glycogen macromolecule by glycogen phosphorylase enzymes to

25 release glucose-1-phosphate and a new shortened glycogen molecule. These disorders are ameliorated by reduction of or characterized by an elevation of glycogen phosphorylase activity. Examples include diabetes, hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemia, hyperlipidemia, atherosclerosis and myocardial ischemia.

- 30 The term glycogen phosphorylase inhibitor refers to any substance or agent or any combination of substances and/or agents which reduces, retards, or eliminates the enzymatic action of glycogen phosphorylase. The currently known enzymatic action of glycogen phosphorylase is the degradation of glycogen by

-22-

catalysis of the reversible reaction of a glycogen macromolecule and inorganic phosphate to glucose-1-phosphate and a glycogen macromolecule which is one glucosyl residue shorter than the original glycogen macromolecule (forward direction of glycogenolysis).

- 5 The term "treating" as used herein includes preventative (e.g., prophylactic) and palliative treatment.

By halo is meant chloro, bromo, iodo, or fluoro.

By alkyl is meant straight chain or branched saturated hydrocarbon.

- Exemplary of such alkyl groups (assuming the designated length encompasses the particular example) are methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tertiary butyl,
10 pentyl, isopentyl, hexyl and isohexyl.

- By alkoxy is meant straight chain or branched saturated alkyl bonded through an oxy. Exemplary of such alkoxy groups (assuming the designated length encompasses the particular example) are methoxy, ethoxy, propoxy, isopropoxy,
15 butoxy, isobutoxy, tertiary butoxy, pentoxy, isopentoxy, hexoxy and isohexoxy.

The expression "pharmaceutically-acceptable anionic salt" refers to nontoxic anionic salts containing anions such as (but not limited to) chloride, bromide, iodide, sulfate, bisulfate, phosphate, acetate, maleate, fumarate, oxalate, lactate, tartrate, citrate, gluconate, methanesulfonate and 4-toluene-sulfonate.

- 20 The expression "pharmaceutically-acceptable cationic salt" refers to nontoxic cationic salts such as (but not limited to) sodium, potassium, calcium, magnesium, ammonium or protonated benzathine (N,N'-dibenzylethylenediamine), choline, ethanolamine, diethanolamine, ethylenediamine, meglamine (N-methyl-glucamine), benethamine (N-benzylphenethylamine), piperazine or tromethamine (2-amino-2-
25 hydroxymethyl-1,3-propanediol).

- The expression "prodrug" refers to compounds that are drug precursors, which following administration, release the drug in vivo via some chemical or physiological process (e.g., a prodrug on being brought to the physiological pH is converted to the desired drug form). Certain exemplary prodrugs upon cleavage
30 release the corresponding free acid, and such hydrolyzable ester-forming residues of the compounds of this invention include but are not limited to carboxylic acid substituents (e.g., R₆ is carboxy, or R₈, R₉ or R₁₂ contains carboxy) wherein the free hydrogen is replaced by (C₁-C₄)alkyl, (C₂-C₁₂)alkanoyloxymethyl, 1-(alkanoyloxy)ethyl

-23-

having from 4 to 9 carbon atoms, 1-methyl-1-(alkanoyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxycarbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxycarbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxycarbonyloxy)ethyl having from 5 to 8 carbon atoms, N-

- 5 (alkoxycarbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxycarbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C₁-C₂)alkylamino(C₂-C₃)alkyl (such as β -dimethylaminoethyl), carbamoyl-(C₁-C₂)alkyl, N,N-di(C₁-C₂)alkylcarbamoyl-(C₁-C₂)alkyl and piperidino-, pyrrolidino- or morpholino(C₂-C₃)alkyl.
- 10 Other exemplary prodrugs release an alcohol of Formula I wherein the free hydrogen of the hydroxy substituent (e.g., R₈, R₉ or R₁₂ contains hydroxy) is replaced by (C₁-C₆)alkanoyloxymethyl, 1-((C₁-C₆)alkanoyloxy)ethyl, 1-methyl-1-((C₁-C₆)alkanoyloxy)ethyl, (C₁-C₆)alkoxycarbonyloxymethyl, N-(C₁-C₆)alkoxycarbonylaminomethyl, succinoyl, (C₁-C₆)alkanoyl, α -amino(C₁-C₄)alkanoyl, arylacyl and α -aminoacyl, or α -aminoacyl- α -aminoacyl wherein said α -aminoacyl moieties are independently any of the naturally occurring L-amino acids found in proteins, P(O)(OH)₂, -P(O)(O(C₁-C₆)alkyl)₂ or glycosyl (the radical resulting from detachment of the hydroxyl of the hemiacetal of a carbohydrate).

- Other exemplary prodrugs include but are not limited to derivatives of
- 20 Formula I wherein R₂ is a free hydrogen which is replaced by R-carbonyl, RO-carbonyl, NRR'-carbonyl where R and R' are each independently (C₁-C₁₀)alkyl, (C₃-C₇)cycloalkyl, benzyl, or R-carbonyl is a natural α -aminoacyl or natural α -aminoacyl-natural α -aminoacyl, -C(OH)C(O)OY wherein (Y is H, (C₁-C₆)alkyl or benzyl), -C(OY₀)Y₁ wherein Y₀ is (C₁-C₄) alkyl and Y₁ is ((C₁-C₆)alkyl, carboxy(C₁-C₆)alkyl, amino(C₁-C₄)alkyl or mono-N- or di-N,N-(C₁-C₆)alkylaminoalkyl, -C(Y₂)Y₃ wherein Y₂ is H or methyl and Y₃ is mono-N- or di-N,N-(C₁-C₆)alkylamino, morpholino, piperidin-1-yl or pyrrolidin-1-yl.

- Other exemplary prodrugs include but are not limited to derivatives of
- Formula I bearing a hydrolyzable moiety at R₃, which release a compound of
- 30 formula I wherein R₃ is a free hydrogen on hydrolysis. Such hydrolyzable moieties at R₃ are/include 1-hydroxy(C₁-C₆)alkyl or 1-hydroxy-1-phenylmethyl.

Other exemplary prodrugs include cyclic structures such as compounds of Formula I wherein R₂ and R₃ are a common carbon, thus forming a five-membered

ring. The linking carbon may be mono- or di-substituted independently with H, (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl or phenyl.

As used herein, the expressions "reaction-inert solvent" and "inert solvent" refers to a solvent which does not interact with starting materials, reagents, intermediates or products in a manner which adversely affects the yield of the desired product.

The chemist of ordinary skill will recognize that certain compounds of this invention will contain one or more atoms which may be in a particular stereochemical or geometric configuration, giving rise to stereoisomers and configurational isomers. All such isomers and mixtures thereof are included in this invention. Hydrates of the compounds of this invention are also included as an aspect of this invention.

The chemist of ordinary skill will recognize that certain combinations of heteroatom-containing substituents listed in this invention define compounds which will be less stable under physiological conditions (e.g. those containing acetal or aминаl linkages). Accordingly, such compounds are less preferred.

The term "R_x ring" wherein x is an integer, for example, "R₉ ring", "R₁₂ ring" or "R₄ ring" as used herein in reference to substitution on the ring refers to moieties wherein the ring is R_x and also wherein the ring is contained within R_x.

As used herein the term mono-N- or di-N,N-(C₁-C_x)alkyl... refers to the (C₁-C_x)alkyl moiety taken independently when it is di-N,N-(C₁-C_x)alkyl....(x refers to integers).


Other features and advantages will be apparent from the specification and claims which describe the invention.

Detailed Description of the Invention


In general the compounds of Formula I can be made by processes which include processes known in the chemical arts, particularly in light of the description contained herein. Certain processes for the manufacture of Formula I compounds are provided as further features of the invention and are illustrated by the following reaction schemes.

-25-


REACTION SCHEME I




REACTION SCHEME II




REACTION SCHEME III



REACTION SCHEME IV




REACTION SCHEME V




-30-


REACTION SCHEME VI



10



20



According to Reaction Scheme I the Formula I compounds, wherein R_1 , R_{10} , R_{11} , A, R_2 , R_3 , R_4 , R_5 and R_6 are as defined above may be prepared by either of two general processes. In the first process the desired Formula I compound may be prepared by coupling the appropriate Formula II indole-2-carboxylic acid, indoline-2-
5 carboxylic acid or benzimidazole-2-carboxylic acid with the appropriate Formula III amine (i.e., acylating the amine). In the second process the desired Formula I compound may be prepared by coupling the appropriate Formula IV compound (i.e., a Formula I compound wherein R_6 is carboxy) with the appropriate alcohol or formula R_8R_9NH or $R_{12}H$ amine, wherein R_8 , R_9 and R_{12} are as defined above (i.e.,
10 acylating the amine or alcohol). The first process (coupling Formula II compounds with Formula III compounds) is typically preferred when R_4 is not H and R_5 is H.

Typically, the Formula II compound is combined with the Formula III compound (or Formula IV compound is combined with the appropriate amine (e.g., $R_{12}H$ or R_8R_9NH)) or alcohol in the presence of a suitable coupling agent. A suitable
15 coupling agent is one which transforms a carboxylic acid into a reactive species which forms an amide or ester linkage on reaction with an amine or alcohol, respectively.

The coupling agent may be a reagent which effects this condensation in a one pot process when mixed together with the carboxylic acid and amine or alcohol.
20 If the acid is to be condensed with an alcohol it is preferable to employ a large excess of the alcohol as the reaction solvent, with or without 1.0 to 1.5 equivalent added dimethylaminopyridine. Exemplary coupling reagents are 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride-hydroxybenzotriazole (DEC/HBT), carbonyldiimidazole, dicyclohexylcarbodiimide/hydroxybenzotriazole
25 (HBT), 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), carbonyldiimidazole/HBT, propanephosphonic anhydride (propanephosphonic acid anhydride, PPA) and diethylphosphorylcyanide. The coupling is performed in an inert solvent, preferably an aprotic solvent at a temperature of about -20°C to about 50°C for about 1 to about 48 hours, in the optional presence of a tertiary amine
30 base such as triethylamine. Exemplary solvents include acetonitrile, dichloromethane, ethyl acetate, dimethylformamide and chloroform or mixtures thereof. An example of a suitable coupling procedure is Procedure A, contained herein (just prior to the EXAMPLES).

-32-

The coupling agent may also be that agent which converts the carboxylic acid to an activated intermediate which is isolated and/or formed in a first step and allowed to react with the amine or alcohol in a second step. Examples of such coupling agents and activated intermediates are thionyl chloride or oxalyl chloride to form the acid chloride, cyanuric fluoride to form an acid fluoride or an alkyl chloroformate such as isobutyl or isopropenyl chloroformate (with a tertiary amine base) to form a mixed anhydride of the carboxylic acid. If the coupling agent is oxalyl chloride it is advantageous to employ a small amount of dimethylformamide as cosolvent with another solvent (such as dichloromethane) to catalyze the formation of the acid chloride. This acid chloride may be coupled by mixing with the Formula III intermediate in an appropriate solvent together with an appropriate base. Appropriate solvent/base combinations are for example, dichloromethane, dimethylformamide or acetonitrile or mixtures thereof in the presence of a tertiary amine base e.g., triethylamine. Other appropriate solvent/base combinations include water or a (C₁-C₆) alcohol or a mixture thereof together with a cosolvent such as dichloromethane, tetrahydrofuran or dioxane and a base such as sodium or potassium carbonate, sodium potassium or lithium hydroxide or sodium bicarbonate in sufficient quantity to consume the acid liberated in the reaction. Use of a phase transfer catalyst (typically 1 to 10 mole %) such as a quaternary ammonium halide (e.g. tetrabutylammonium bromide or methyl trioctylammonium chloride) is advantageous when a mixture of only partially miscible cosolvents is employed (e.g. dichloromethane-water or dichloromethane-methanol). Use of these coupling agents and appropriate selection of solvents and temperatures are known to those skilled in the art or can be readily determined from the literature. These and other exemplary conditions useful for coupling carboxylic acids are described in Houben-Weyl, Vol XV, part II, E. Wunsch, Ed., G. Theime Verlag, 1974, Stuttgart, and M. Bodansky, Principles of Peptide Synthesis, Springer-Verlag Berlin 1984, and The Peptides. Analysis, Synthesis and Biology (ed. E. Gross and J. Meienhofer), vols 1-5 (Academic Press NY 1979-1983).

The Formula IV compounds wherein R₁, R₁₀, R₁₁, A, R₂, R₃, R₄ and R₅ are as defined above may be prepared from the corresponding Formula V ester (i.e., Formula I compounds wherein R₆ is (C₁-C₅)alkoxycarbonyl or benzyloxycarbonyl) by hydrolysis with aqueous alkali at a temperature of about

-33-

-20°C to about 100°C, typically at about 20°C, for about 30 minutes to about 24 hours.

Alternatively, Formula IV compounds are prepared by activation of a Formula II indole carboxylic acid with a coupling agent (as described above) which gives an
5 activated intermediate (such as an acid chloride, acid fluoride, or mixed anhydride) which is then allowed to react with a compound of Formula III wherein R_3 , R_4 and R_5 are as described above and R_6 is carboxy, in a suitable solvent in the presence of a suitable base. Suitable solvents include water, or methanol or a mixture thereof, together with a cosolvent such as dichloromethane, tetrahydrofuran, or
10 dioxane. Suitable bases include sodium, potassium or lithium hydroxides, sodium or potassium bicarbonate, sodium or potassium carbonate, or potassium carbonate together with tetrabutyl ammonium bromide (1 equivalent) in sufficient quantity to consume the acid liberated in the reaction (generally that quantity sufficient to maintain the pH of the reaction at greater than 8). The base may be added
15 incrementally together with the activated intermediate to effect proper pH control of the reaction. The reaction is conducted generally between -20°C and 50°C. Isolation procedures are tailored by one skilled in the art to remove impurities, but typically consist of removal of water-miscible cosolvents by evaporation, extraction of impurities at high pH with an organic solvent, acidification to low pH (1-2) and
20 filtration, or extraction of the desired product with a suitable solvent such as ethyl acetate or dichloromethane.

The Formula V compound may be prepared by coupling the appropriate Formula III compound wherein R_6 is alkoxycarbonyl and the appropriate Formula II compound in an analogous procedure to that described above (e.g., Procedure A).

25 Alternatively, Formula I compounds which contain sulfur atoms in the sulfoxide or sulfone oxidation state may be prepared from the corresponding Formula I compounds having the sulfur atom in the unoxidized form, by treatment with a suitable oxidizing agent, such as m-chloroperoxybenzoic acid in dichloromethane at a temperature of about 0°C to about 25°C for about 1 to about
30 48 hours using about 1 to about 1.3 equivalent for conversion to the sulfoxide oxidation state and greater than about 2 equivalents for conversion to the sulfone oxidation state.

Some of the preparation methods described herein may require protection of remote functionality (i.e., primary amine, secondary amine, carboxyl in Formula I precursors). The need for such protection will vary depending on the nature of the remote functionality and the conditions of the preparation methods. The need for
5 such protection is readily determined by one skilled in the art. The use of such protection/deprotection methods is also within the skill in the art. For a general description of protecting groups and their use, see T.W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, New York, 1991.

For example, in Reaction Scheme I certain Formula I compounds contain
10 primary amine, secondary amine or carboxylic acid functionality in the part of the molecule defined by R_6 which may interfere with the intended coupling reaction of Reaction Scheme I, if the Formula III intermediate or $R_{1,2}H$ or R_8R_9NH amine is left unprotected. Accordingly, the primary amine, secondary amine or carboxylic acid functionality may be protected, where it is present in the R_6 moieties of the Formula
15 III intermediate R_8R_9NH or $R_{1,2}H$ amine by an appropriate protecting group during the coupling reaction of Reaction Scheme I. The product of such coupling reaction in such a case is a Formula I compound containing the protecting group. This protecting group is removed in a subsequent step to provide the Formula I compound. Suitable protecting groups for amine and carboxylic acid protection
20 include those protecting groups commonly used in peptide synthesis (such as N-t-butoxycarbonyl, N-carbobenzyloxy, and 9-fluorenylmethylenoxycarbonyl for amines and lower alkyl or benzyl esters for carboxylic acids) which are not chemically reactive under the coupling conditions described above (and immediately preceding the Examples herein as Procedure A) and can be removed without chemically
25 altering other functionality in the Formula I compound.

The starting indole-2-carboxylic acids and indoline-2-carboxylic acids used in Reaction Scheme I, when not commercially available or known in the prior art (such art is extensively published), are available by conventional synthetic methods. For example, according to Reaction Scheme II the Formula VII indole ester (wherein A is
30 not nitrogen) may be prepared from the Formula VI compound (wherein Q is selected to achieve the desired A as defined above, except for N) via a Fischer Indole synthesis (see The Fischer Indole Synthesis Robinson, B. (Wiley, New York, 1982)) followed by saponification of the resulting Formula VII indole ester to yield the

corresponding Formula VIII acid. The starting aryl hydrazone may be prepared by condensation of a readily available hydrazine with the appropriate carbonyl derivative or via the Japp-Klingeman reaction (see Organic Reactions, Phillips, R. R., 1959, 10, 143).

- 5 Alternatively, the Formula VIIIa indole 2-carboxylic acid may be prepared by condensation of a Formula IX ortho methyl nitro compound with an oxalate ester to yield the Formula X indole ester followed by reduction of the nitro group and subsequent hydrolysis.

 This three step process is known as the Reissert indole synthesis (Reissert, 10 *Chemische Berichte* 1897, 30, 1030). Conditions for accomplishing this sequence, and references thereto, are described in the literature (Kermack, et al., *J. Chem. Soc.* 1921, 119, 1602; Cannon et al., *J. Med. Chem.* 1981, 24, 238; Julian, et al in *Heterocyclic Compounds*, vol 3 (Wiley, New York, NY, 1962, R.C. Elderfield, ed.) p 18). An example of the specific implementation of this sequence is Examples 10A- 15 10C herein.

 3-Halo-5-chloro-1H-indole-2-carboxylic acids may also be prepared by halogenation of 5-chloro-1H-indole-2-carboxylic acids.

 According to Reaction Scheme III the Formula XI benzimidazole-2-carboxylic acid intermediates may be prepared by condensation of a Formula XIII ortho- 20 diamino compound with glycolic acid, followed by oxidation of the resulting Formula XII benzimidazole-2-methanol (Bistrzycki, A. and Przeworski, G. *Ber.* 1912, 45, 3483).

 Alternatively, (to Reaction Scheme II) the Formula XIV substituted indolines may be prepared by reduction of the corresponding Formula XV indoles with a reducing agent such as magnesium in methanol at a temperature of about 25°C to 25 about 65°C for about 1 to about 48 hours (Reaction Scheme III).

 Formula XVI indoline carboxylic acids are prepared by saponification of the corresponding Formula XVII ester (Reaction Scheme III). The Formula XVII ester is prepared by reduction of the corresponding Formula VII indole ester with a reducing agent such as magnesium in methanol as described for the conversion of the 30 Formula XV compound to the Formula XIV compound above.

 The following paragraphs describe ways to prepare the various amines which are used in the above Reaction Schemes.

According to Reaction Scheme IV a Formula XXIII alpha-amino acid may be protected on nitrogen with an appropriate protecting group (P_1) (e.g., t-Boc) to form a Formula XXIV compound. One skilled in the art can readily select an appropriate protecting group and a method for its introduction. For example, two common

5 protecting groups are t-Boc (introduced by treating the amino acid with di-t-butylidicarbonate in a preferably protic suitable solvent or solvent mixture at high pH) and CBZ (introduced by treating the amino acid with benzylchloroformate in a suitable, preferably protic solvent or solvent mixture and base). The Formula XXIV compound is coupled (in an analogous procedure to the coupling process

10 described in Reaction Scheme I) with an appropriate R_8R_9NH or HR_{12} amine to form a Formula XXV compound, which is then deprotected resulting in the Formula IIIb compound (i.e., Formula III compound wherein R_6 is $C(O)R_{12}$ or $C(O)NR_8R_9$). If the protecting group is t-Boc by treatment of the Formula XXV compound with an acid in a suitable, preferably aprotic, solvent. Acids for this deprotection include HCl,

15 $MeSO_3H$ or trifluoroacetic acid.

According to Reaction Scheme V a Formula XXXI compound (N-protected Formula III amine where R_6 is (C_1-C_8) alkoxycarbonyl or benzyloxycarbonyl) may be prepared from the corresponding Formula XXX unprotected amino acid via N-protection (yielding a Formula XXXIII protected amino acid) followed by

20 esterification. For example, the Formula XXXIII compound may be esterified with the appropriate alcohol and an acid catalyst such as hydrogen chloride or thionyl chloride, or in the case of tert-butanol by treatment of the amino acid with isobutylene and an acid catalyst such as concentrated sulfuric acid or by treatment with an alkyl halide (e.g., methyl iodide) and base (e.g., potassium carbonate).

25 Alternatively, the esterification may precede the protection step.

According to Reaction Scheme VI the Formula XXX compounds wherein R_3 is not H utilized in Reaction Scheme V may be prepared as follows. The Formula XLI amino acids may be prepared by N-alkylation of the Formula XL protected (P_1) amino acids by treatment with an appropriate base and alkylating agent. Specific

30 procedures for this alkylation are described by Benoiton, Can. J. Chem. 1977, 55, 906-910, and Hansen, J. Org. Chem. 1985, 50 945-950. For example, when R_3 is methyl, and P_1 is Boc, sodium hydride and methyl iodide in tetrahydrofuran are

-37-

utilized. Deprotection of the Formula XLI compound yields the desired Formula XXX compound.

Alternatively, a Formula XLII amino acid may be N-alkylated by a three-step sequence involving reductive benzylation (such as with benzaldehyde, Pd/C-
5 catalyzed hydrogenation) to give the mono-N-benzyl derivative and reductive amination with the appropriate carbonyl compound (for example with formaldehyde and sodium cyanoborohydride to introduce R_3 as methyl) to give the N-Benzyl, N- R_3 -substituted amino acid. The N-benzyl protecting group is conveniently removed (for example by hydrogenation with an appropriate catalyst) to yield the Formula XXX
10 compound. Specific conditions for this three step alkylation procedure are described by Reinhold et al., J. Med. Chem., 1968, 11, 258-260.

The immediately preceding preparation may also be used to introduce an R_3 moiety into a Formula IIIa intermediate (which is a Formula III intermediate wherein R_3 is H).

15 The amino acids used in the schemes herein (e.g., XL, XLII), if not commercially available, or reported in the literature, may be prepared by a variety of methods known to those skilled in the art. For example, the Strecker synthesis or variations thereof may be used. Accordingly, an aldehyde (R_4 CHO), sodium or potassium cyanide and ammonium chloride react to form the corresponding
20 aminonitrile. The aminonitrile is hydrolyzed with mineral acid to form the desired Formula XLII $R_4C(NH_2)COOH$ amino acid. Alternatively, the Bucherer-Berg method may be used wherein a hydantoin is formed by heating an aldehyde (R_4 CHO) with ammonium carbonate and potassium cyanide followed by hydrolysis (for example, with barium hydroxide in refluxing dioxane) with acid or base to form the desired
25 Formula XLII $R_4C(NH_2)COOH$ amino acid.

Other methods for synthesis of α -amino acids are also reported in the literature which would permit one skilled in the art to prepare the desired Formula XLII $R_4C(NH_2)COOH$ intermediate necessary for the synthesis of Formula I compounds.

30 Suitable methods for the synthesis and/or resolution of Formula XLII compounds are found in reviews by Duthaler (Tetrahedron 1994, 50, 1539-1650), or by Williams (R. M. Williams, Synthesis of optically active amino acids. Pergamon: Oxford, U.K., 1989).

A specific method for the synthesis of a Formula XLII intermediate in either enantiomeric form from the corresponding R_4X ($X = Cl, Br, \text{ or } I$) intermediate is the procedure of Pirrung and Krishnamurthy (J. Org. Chem. 1993, 58, 957-958), or by the procedure of O'Donnell, et al. (J. Am. Chem. Soc. 1989, 111, 2353-2355). The
5 required R_4X intermediates are readily prepared by many methods familiar to the chemist skilled in the art. For example, those compounds when R_4X is $ArCH_2X$ may be prepared by radical halogenation of the compound $ArCH_3$ or by formylation of the arene $Ar-H$ and conversion of the alcohol to the bromide.

Another specific method for the synthesis of Formula XLII intermediates in
10 either enantiomeric form is that of Corey and Link (J. Am. Chem. Soc. 1992, 114, 1906-1908). Thus, an intermediate of formula R_4COCCl_3 is reduced enantiospecifically to intermediate $R_4CH(OH)CCl_3$, which is converted on treatment with azide and base to an intermediate $R_4CH(N_3)COOH$, which is reduced by catalytic hydrogenation to the desired Formula XLII compound. The requisite
15 trichloromethyl ketone R_4COCCl_3 is obtained by reaction of the aldehyde R_4CHO with trichloromethide anion followed by oxidation (Gallina and Giordano, Synthesis 1989, 466-468).

A compound of the formula R_8NH_2 or R_9NH_2 is monoalkylated with a carbonyl compound corresponding to R_8 or R_9 , respectively, under appropriate
20 reductive amination conditions, to give a formula R_8R_9NH amine. To avoid dialkylation, it may be preferable to protect the amines (R_8NH_2 or R_9NH_2) with a suitable protecting group P_T to give $R_8(P_T)NH$ or $R_9(P_T)NH$, for example by reaction with benzaldehyde and a reducing agent. The protected amines are monoalkylated with a carbonyl compound corresponding to R_9 or R_8 respectively, under suitable
25 reductive amination conditions, to give $R_8R_9N(P_T)$. The protecting group (P_T) is removed (e.g. by exhaustive catalytic hydrogenation when P_T is benzyl) to give a compound of formula R_8R_9NH . Appropriate reductive amination conditions are available from the literature to one skilled in the art. These conditions include those reported by Borch et al. (J. Am. Chem. Soc. 1971, 2897-2904) and those reviewed
30 by Emerson (Organic Reactions, Wiley: New York, 1948 (14), 174), Hutchins et al. (Org. Prep. Proced. Int 1979 (11), 20, and Lane et al. (Synthesis 1975, 135). Reductive amination conditions favoring N-monoalkylation include those reported by Morales, et al. (Synthetic Communications 1984, 1213-1220) and Verardo et al.

(Synthesis 1992 121-125). The R_8NH_2 or R_9NH_2 amines may also be monoalkylated with R_8X or R_9X , respectively, where X is chloride, bromide, tosylate or mesylate.

Alternatively, an intermediate of formula $R_8(P_7)NH$ or $R_9(P_7)NH$ may be alkylated with R_8X or R_9X , and the protecting group removed to give a compound of formula

5 R_8R_9NH .

Additional methods may be used to prepare formula R_8R_9NH amines wherein R_8-NH or R_9-NH are oxygen-nitrogen linked. Thus a readily available compound of formula (C_1-C_4) alkoxycarbonyl-NHOH or $NH_2CONHOH$ is dialkylated on nitrogen and oxygen by treatment with base and excess suitable alkylating agent ($R-X$) to give the
10 corresponding (C_1-C_4) alkoxycarbonyl-N(R)OR which is then hydrolyzed to give a compound of formula R_8R_9NH (wherein $R_8=R_9=R$). Suitable conditions, base, and alkylating agent include those described by Goel and Krolls (Org. Prep. Proced. Int. 1987, 19, 75-78) and Major and Fleck (J. Am. Chem. Soc. 1928, 50, 1479).

Alternatively, N-hydroxyurea ($NH_2CONH(OH)$) may be sequentially alkylated, first on
15 oxygen to give $NH_2CONH(OR')$, then on nitrogen to give $NH_2CON(R'')(OR')$, by successive treatment with the alkylating agents $R'X$ and $R''X$, respectively, in the presence of a suitable base. Suitable base and alkylating agents include those described by Kreutzkamp and Messinger (Chem. Ber. 100, 3463-3465 (1967) and Danen et al (J. Am. Chem. Soc. 1973, 95, 5716-5724). Hydrolysis of these alkylated
20 hydroxyurea derivatives yields the amines $R'ONH_2$ and $R'ONHR''$, which correspond to certain formula R_8R_9NH amines. The chemist skilled in the art can adapt the procedures described in this paragraph to other alkylating agents R , R' and $R''-X$ to prepare other amines of formula R_8R_9NH wherein R_8-N or R_9-N are oxygen-nitrogen linked. Uno et al (SynLett 1991, 559-560) describe the BF_3 -catalyzed addition of an
25 organometallic reagent $R-Li$ to an O-alkyl oxime of formula $R'CH=N-OR''$, to give compounds of formula $R'RCH-NH(OR'')$. This route may also be used to give compounds of formula R_8R_9NH wherein one of R_8-NH or R_9-NH are oxygen-nitrogen linked.

Prodrugs of this invention where a carboxyl group in a carboxylic acid of
30 Formula I is replaced by an ester may be prepared by combining the carboxylic acid with the appropriate alkyl halide in the presence of a base such as potassium carbonate in an inert solvent such as dimethylformamide at a temperature of about 0 to 100°C for about 1 to about 24 hours. Alternatively the acid is combined with

-40-

appropriate alcohol as solvent in the presence of a catalytic amount of acid such as concentrated sulfuric acid at a temperature of about 20 to 120°C, preferably at reflux, for about 1 hour to about 24 hours. Another method is the reaction of the acid with a stoichiometric amount of the alcohol in the presence of a catalytic
5 amount of acid in an inert solvent such as tetrahydrofuran, with concomitant removal of the water being produced by physical (e.g. Dean-Stark trap) or chemical (e.g. molecular sieves) means.

Prodrugs of this invention where an alcohol function has been derivatized as an ether may be prepared by combining the alcohol with the appropriate alkyl
10 bromide or iodide in the presence of a base such as potassium carbonate in an inert solvent such as dimethylformamide at a temperature of about 0 to 100°C for about 1 to about 24 hours. Alkanoylaminomethyl ethers may be obtained by reaction of the alcohol with a bis-(alkanoylamino)methane in the presence of a catalytic amount of acid in an inert solvent such as tetrahydrofuran, according to a
15 method described in US 4,997, 984. Alternatively, these compounds may be prepared by the methods described by Hoffman et al. in J. Org. Chem. 1994, 59, 3530.

The dialkylphosphate esters may be prepared by reaction of the alcohol with a dialkyl chlorophosphate in the presence of a base in an inert solvent such as
20 tetrahydrofuran. The dihydrogen phosphates may be prepared by reaction of the alcohol with a diaryl or dibenzyl chlorophosphate as described above, followed by hydrolysis or hydrogenation in the presence of a noble metal catalyst, respectively.

Glycosides are prepared by reaction of the alcohol and a carbohydrate in an inert solvent such as toluene in the presence of acid. Typically the water formed in
25 the reaction is removed as it is being formed as described above. An alternate procedure is the reaction of the alcohol with a suitably protected glycosyl halide in the presence of base followed by deprotection.

N-(1-hydroxyalkyl) amides, N-(1-hydroxy-1-(alkoxycarbonyl)methyl) amides or compounds where R₂ has been replaced by C(OH)C(O)OY may be prepared by the
30 reaction of the parent amide or indole with the appropriate aldehyde under neutral or basic conditions (e.g. sodium ethoxide in ethanol) at temperatures between 25 and 70°C. N-alkoxymethyl indoles or N-1-(alkoxy)alkyl indoles can be obtained by reaction of the N-unsubstituted indole with the necessary alkyl halide in the

presence of a base in an inert solvent. 1-(N,N-dialkylaminomethyl) indole, 1-(1-(N,N-dialkylamino)ethyl) indole and N,N-dialkylaminomethyl amides (e.g. $R_3 = CH_2N(CH_3)_2$) may be prepared by the reaction of the parent N-H compound with the appropriate aldehyde and amine in an alcoholic solvent at 25 to 70°C.

- 5 The prodrugs of this invention where R_2 and R_3 are a common carbon may be prepared by reaction of the parent compound (drug) with benzaldehyde or a ketone or its dimethyl acetal in an inert solvent in the presence of a catalytic amount of acid with concomitant water or methanol removal.

- The starting materials and reagents for the above described reaction
10 schemes (e.g., amines, substituted indole carboxylic acids, substituted indoline carboxylic acids, amino acids), although the preparation of most of which are described above, are also readily available or can be easily synthesized by those skilled in the art using conventional methods of organic synthesis. For example, many of the intermediates used herein to prepare compounds of Formula I are, are
15 related to, or are derived from amino acids found in nature, in which there is a large scientific interest and commercial need, and accordingly many such intermediates are commercially available or are reported in the literature or are easily prepared from other commonly available substances by methods which are reported in the literature. Such intermediates include, for example, Formula XXX, Formula XLII,
20 Formula XXXII and Formula XXXIII compounds.

- Some compounds of Formula I have asymmetric carbon atoms and therefore are enantiomers or diastereomers. Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods known per se, for example, by chromatography and/or fractional
25 crystallization. Enantiomers (e.g., of Formula III, VIII or IX) can be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., alcohol), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. All such isomers, including diastereomers, enantiomers and
30 mixtures thereof are considered as part of this invention.

 Although many compounds of this invention are not ionizable at physiological conditions, some of the compounds of this invention are ionizable at physiological conditions. Thus, for example some of the compounds of this

invention are acidic and they form a salt with a pharmaceutically acceptable cation. All such salts are within the scope of this invention and they can be prepared by conventional methods. For example, they can be prepared simply by contacting the acidic and basic entities, usually in a stoichiometric ratio, in either an aqueous, non-
5 aqueous or partially aqueous medium, as appropriate. The salts are recovered either by filtration, by precipitation with a non-solvent followed by filtration, by evaporation of the solvent, or, in the case of aqueous solutions, by lyophilization, as appropriate.

In addition, some of the compounds of this invention are basic, and they
10 form a salt with a pharmaceutically acceptable anion. All such salts are within the scope of this invention and they can be prepared by conventional methods. For example, they can be prepared simply by contacting the acidic and basic entities, usually in a stoichiometric ratio, in either an aqueous, non-aqueous or partially aqueous medium, as appropriate. The salts are recovered either by filtration, by
15 precipitation with a non-solvent followed by filtration, by evaporation of the solvent, or, in the case of aqueous solutions, by lyophilization, as appropriate. In addition, when the compounds of this invention form hydrates or solvates they are also within the scope of the invention.

The utility of the compounds of the present invention as medical agents in
20 the treatment of metabolic diseases (such as are detailed herein) in mammals (e.g., humans) is demonstrated by the activity of the compounds of this invention in conventional assays and the in vitro and in vivo assays described below. Such assays also provide a means whereby the activities of the compounds of this invention can be compared with the activities of other known compounds. The
25 results of these comparisons are useful for determining dosage levels in mammals, including humans, for the treatment of such diseases.

The purified human liver glycogen phosphorylase a (HLGPa) is obtained by the following procedure.

Expression and fermentation:

30 The HLGP cDNA is expressed from plasmid pKK233-2 (Pharmacia Biotech. Inc., Piscataway, New Jersey) in E. coli strain XL-1 Blue (Stratagene Cloning Systems, LaJolla, CA). The strain is inoculated into LB medium (consisting of 10 g tryptone, 5 g yeast extract, 5 g NaCl, and 1 ml 1N NaOH per liter) plus 100 mg/L

-43-

ampicillin, 100 mg/L pyridoxine and 600 mg/L MnCl_2 and grown at 37°C to a cell density of $\text{OD}_{550} = 1.0$. At this point, the cells are induced with 1 mM isopropyl-1-thio- β -D-galactoside (IPTG). Three hours after induction the cells are harvested by centrifugation and cell pellets are frozen at -70°C until needed for purification.

5 Purification of Glycogen Phosphorylase:

The cells in pellets described above are resuspended in 25 mM β -glycerophosphate (pH 7.0) with 0.2 mM DTT, 1 mM MgCl_2 , plus the following protease inhibitors:

	0.7 $\mu\text{g/mL}$	Pepstatin A
10	0.5 $\mu\text{g/mL}$	Leupeptin
	0.2 mM	Phenylmethylsulfonyl fluoride (PMSF), and
	0.5 mM	EDTA,

lysed by pretreatment with 200 $\mu\text{g/mL}$ lysozyme and 3 $\mu\text{g/mL}$ DNAase followed by sonication in 250 mL batches for 5 x 1.5 minutes on ice using a Branson Model 450
 15 ultrasonic cell disrupter (Branson Sonic Power Co., Danbury CT). The lysates are cleared by centrifugation at 35,000 X g for one hour followed by filtration through 0.45 micron filters. HLGP in the soluble fraction of the lysates (estimated to be less than 1% of the total protein) is purified by monitoring the enzyme activity (as described in HLGPa Activity Assay section, below) from a series of chromatographic
 20 steps detailed below.

Immobilized Metal Affinity Chromatography (IMAC):

This step is based on the method of Luong et al (Luong et al. Journal of Chromatography (1992) 584, 77-84.). 500 mL of the filtered soluble fraction of cell lysates (prepared from approximately 160 g of original cell pellet) are loaded onto a
 25 130 mL column of IMAC Chelating-Sepharose (Pharmacia LKB Biotechnology, Piscataway, New Jersey) which has been charged with 50 mM CuCl_2 and 25 mM β -glycerophosphate, 250 mM NaCl and 1 mM imidazole at pH 7 equilibration buffer. The column is washed with equilibration buffer until the A_{280} returns to baseline. The sample is then eluted from the column with the same buffer containing 100 mM
 30 imidazole to remove the bound HLGP and other bound proteins. Fractions containing the HLGP activity are pooled (approximately 600 mL), and ethylenediaminetetraacetic acid (EDTA), DL-dithiothreitol (DTT), phenylmethylsulfonyl fluoride (PMSF), leupeptin and pepstatin A are added to obtain 0.3 mM, 0.2 mM, 0.2

-44-

mM, 0.5 μ g/mL and 0.7 μ g/mL concentrations respectively. The pooled HLGP is desalted over a Sephadex G-25 column (Sigma Chemical Co., St. Louis, Missouri) equilibrated with 25 mM Tris-HCl (pH 7.3), 3 mM DTT buffer (Buffer A) to remove imidazole and is stored on ice until the second chromatographic step.

5 5' - AMP-Sepharose Chromatography:

The desalted pooled HLGP sample (approximately 600mL) is next mixed with 70 mL of 5'-AMP Sepharose (Pharmacia LKB Biotechnology, Piscataway, New Jersey) which has been equilibrated with Buffer A (see above). The mixture is gently agitated for one hour at 22°C then packed into a column and washed with Buffer A
10 until the A280 returns to baseline. HLGP and other proteins are eluted from the column with 25 mM Tris-HCl, 0.2 mM DTT and 10 mM adenosine 5'-monophosphate (AMP) at pH 7.3 (Buffer B). HLGP-containing fractions are pooled following identification by determining enzyme (described below) activity and visualizing the M_r approximately 97 kdal HLGP protein band by sodium dodecyl sulfate polyacrylamide
15 gel electrophoresis (SDS-PAGE) followed by silver staining (2D-silver Stain II "Daiichi Kit", Daiichi Pure Chemicals Co., LTD., Tokyo, Japan) and then pooled. The pooled HLGP is dialyzed into 25 mM β -glycerophosphate, 0.2 mM DTT, 0.3 mM EDTA, 200mM NaCl, pH 7.0 buffer (Buffer C) and stored on ice until use.

Determination of HLGP Enzyme Activity:

20 A) Activation of HLGP: Conversion of HLGPb to HLGPa

Prior to the determination of HLGP enzyme activity, the enzyme is converted from the inactive form as expressed in *E. coli* strain XL-1 Blue (designated HLGPb) (Stragene Cloning Systems, La Jolla, California) to the active form (designated HLGPa) by phosphorylation of HLGP using phosphorylase kinase as follows:

25 HLGPb reaction with Immobilized Phosphorylase Kinase

Phosphorylase kinase (Sigma Chemical Company, St. Louis, MO) is immobilized on Affi-Gel 10 (BioRad Corp., Melville, NY) as per the manufacturer's instructions. In brief, the phosphorylase kinase enzyme (10 mg) is incubated with washed Affi-Gel beads (1 mL) in 2.5 mL of 100 mM HEPES and 80 mM CaCl_2 at pH
30 7.4 for 4 hours at 4°C. The Affi-Gel beads are then washed once with the same buffer prior to blocking with 50 mM HEPES and 1 M glycine methyl ester at pH 8.0 for one hour at room temperature. Blocking buffer is removed and replaced with 50 mM HEPES (pH 7.4), 1 mM β -mercaptoethanol and 0.2% NaN_3 for storage. Prior to

use to convert HLGPb to HLGP_a, the Affi-Gel immobilized phosphorylase kinase beads are equilibrated by washing in the buffer used to perform the kinase reaction, consisting of 25mM β-glycerophosphate, 0.3 mM DTT, and 0.3mM EDTA at pH 7.8 (kinase assay buffer).

- 5 The partially purified, inactive HLGPb obtained from 5'-AMP-Sepharose chromatography above is diluted 1:10 with the kinase assay buffer then mixed with the aforementioned phosphorylase kinase enzyme immobilized on the Affi-Gel beads. NaATP is added to 5 mM and MgCl₂ to 6 mM. The resulting mixture is mixed gently at 25°C for 30 to 60 minutes. The sample is removed from the beads
- 10 and the percent activation of HLGPb by conversion to HLGP_a is estimated by determining HLGP enzyme activity in the presence and absence of 3.3 mM AMP. The percentage of total HLGP enzyme activity due to HLGP_a enzyme activity (AMP-independent) is then calculated as follows:

$$15 \quad \% \text{ of total HLGP as HLGP}_a = \frac{\text{HLGP activity -AMP}}{\text{HLGP activity +AMP}}$$

B) HLGP_a Activity Assay:

- The hypoglycemic activity (also the other disease/condition treating/preventing activities described herein) of the compounds of this invention
- 20 can be indirectly determined by assessing the effect of the compounds of this invention on the activity of the activated form of glycogen phosphorylase (GP_a) by one of two methods; glycogen phosphorylase a activity is measured in the forward direction by monitoring the production of glucose-1-phosphate from glycogen or by following the reverse reaction, measuring glycogen synthesis from glucose-1-
- 25 phosphate by the release of inorganic phosphate. All reactions are run in triplicate in 96-well microtiter plates and the change in absorbance due to formation of the reaction product is measured at the wavelength specified below in a MCC/340 MKII Elisa Reader (Lab Systems, Finland), connected to a Titertech Microplate Stacker (ICN Biomedical Co, Huntsville, Alabama).
- 30 To measure the HLGP_a enzyme activity in the forward direction, the production of glucose-1-phosphate from glycogen is monitored by the multienzyme coupled general method of Pesce et al. [Pesce, M.A., Bodourian, S.H., Harris, R.C. and Nicholson, J.F. (1977) Clinical Chemistry 23, 1711-1717] modified as follows: 1

-46-

- to 100 μ g phosphorylase a, 10 units phosphoglucomutase and 15 units glucose-6-phosphate dehydrogenase (Boehringer Mannheim Biochemicals, Indianapolis, IN) are diluted to 1 mL in Buffer A (described hereinafter). Buffer A is at pH 7.2 and contains 50 mM HEPES, 100 mM KCl, 2.5 mM ethyleneglycoltetraacetic acid (EGTA), 2.5 mM $MgCl_2$, 3.5 mM KH_2PO_4 and 0.5 mM dithiothreitol. 20 μ l of this stock is added to 80 μ l of Buffer A containing 0.47 mg/mL glycogen, 9.4 mM glucose, 0.63 mM of the oxidized form of nicotinamide adenine dinucleotide phosphate ($NADP^+$). The compounds to be tested are added as 5 μ L of solution in 14% dimethylsulfoxide (DMSO) prior to the addition of the enzymes. The basal rate of HLGPa enzyme activity in the absence of inhibitors is determined by adding 5 μ L of 14% DMSO and a fully-inhibited rate of HLGPa enzyme activity is obtained by adding 20 μ L of 50 mM of the positive control test substance, caffeine. The reaction is followed at room temperature by measuring the conversion of oxidized $NADP^+$ to reduced NADPH at 340 nm.
- 15 To measure HLGPa enzyme activity in the reverse direction, the conversion of glucose-1-phosphate into glycogen plus inorganic phosphate is measured by the general method described by Engers et al. [Engers, H.D., Shechosky, S. and Madsen, N.B. (1970) Can. J. Biochem. 48, 746-754] modified as follows: 1 to 100 μ g HLGPa is diluted to 1 mL in Buffer B (described hereinafter). Buffer B is at pH 7.2 and contains 50 mM HEPES, 100 mM KCl, 2.5 mM EGTA, 2.5 mM $MgCl_2$, and 0.5 mM dithiothreitol. 20 μ L of this stock is added to 80 μ L of Buffer B with 1.25 mg/mL glycogen, 9.4 mM glucose, and 0.63 mM glucose-1-phosphate. The compounds to be tested are added as 5 μ L of solution in 14% DMSO prior to the addition of the enzyme. The basal rate of HLGPa enzyme activity in the absence of added inhibitors is determined by adding 5 μ L of 14% DMSO and a fully-inhibited rate of HLGPa enzyme activity is obtained by adding 20 μ L of 50 mM caffeine. This mixture is incubated at room temperature for 1 hour and the inorganic phosphate released from the glucose-1-phosphate is measured by the general method of Lanzetta et al. [Lanzetta, P.A., Alvarez, L.J., Reinach, P.S. and Candia, O.A. (1979) Anal. Biochem. 100, 95-97] modified as follows: 150 μ L of 10 mg/mL ammonium molybdate, 0.38 mg/mL malachite green in 1 N HCl is added to 100 μ L of the enzyme mix. After a 20 minute incubation at room temperature, the absorbance is measured at 620 nm.

-47-

The compounds of this invention are readily adapted to clinical use as hypoglycemic agents. The hypoglycemic activity of the compounds of this invention can be determined by the amount of test compound that reduces glucose levels relative to a vehicle without test compound in male ob/ob mice. The test also allows
5 the determination of an approximate minimal effective dose (MED) value for the in vivo reduction of plasma glucose concentration in such mice for such test compounds.

Since the concentration of glucose in blood is closely related to the development of diabetic disorders, these compounds by virtue of their hypoglycemic
10 action, prevent, arrest and/or regress diabetic disorders.

Five to eight week old male C57BL/6J-ob/ob mice (obtained from Jackson Laboratory, Bar Harbor, ME) are housed five per cage under standard animal care practices. After a one week acclimation period, the animals are weighed and 25 microliters of blood are collected from the retro-orbital sinus prior to any treatment.
15 The blood sample is immediately diluted 1:5 with saline containing 0.025% sodium heparin, and held on ice for metabolite analysis. Animals are assigned to treatment groups so that each group has a similar mean for plasma glucose concentration. After group assignment, animals are dosed orally each day for four days with the vehicle consisting of either: 1) 0.25% w/v methyl cellulose in water without pH
20 adjustment; or 2) 0.1% Pluronic® P105 Block Copolymer Surfactant (BASF Corporation, Parsippany, NJ) in 0.1% saline without pH adjustment. On day 5, the animals are weighed again and then dosed orally with the test compound or the vehicle alone. All drugs are administered in vehicle consisting of either: 1) 0.25% w/v methyl cellulose in water without pH adjustment; or 2) 10% DMSO/0.1% Pluronic®
25 P105 (BASF Corporation, Parsippany, NJ) in 0.1% saline without pH adjustment. The animals are then bled from the retro-orbital sinus three hours later for determination of blood metabolite levels. The freshly collected samples are centrifuged for two minutes at 10,000 x g at room temperature. The supernatant is analyzed for glucose, for example, by the Abbott VP™ (Abbott Laboratories, Diagnostics Division, Irving, TX) and VP Super System® Autoanalyzer (Abbott
30 Laboratories, Irving, TX), using the A-Gent™ Glucose-UV Test reagent system (Abbott Laboratories, Irving, TX) (a modification of the method of Richterich and Dauwalder,

Schweizerische Medizinische Wochenschrift, 101, 860 (1971)) (hexokinase method) using a 100 mg/dL standard. Plasma glucose is then calculated by the equation:

Plasma glucose (mg/dL) = Sample value x 5 x 1.784 = 8.92 x Sample value

where 5 is the dilution factor and 1.784 is the plasma hematocrit adjustment

5 (assuming the hematocrit is 44%).

The animals dosed with vehicle maintain substantially unchanged hyperglycemic glucose levels (e.g., greater than or equal to 250 mg/dL), animals treated with test compounds at suitable doses have significantly depressed glucose levels. Hypoglycemic activity of the test compounds is determined by statistical
10 analysis (unpaired t-test) of the mean plasma glucose concentration between the test compound group and vehicle-treated group on day 5. The above assay carried out with a range of doses of test compounds allows the determination of an approximate minimal effective dose (MED) value for the in vivo reduction of plasma glucose concentration.

15 The compounds of this invention are readily adapted to clinical use as hyperinsulinemia reversing agents, triglyceride lowering agents and hypocholesterolemic agents. Such activity can be determined by the amount of test compound that reduces insulin, triglycerides or cholesterol levels relative to a control vehicle without test compound in male ob/ob mice.

20 Since the concentration of cholesterol in blood is closely related to the development of cardiovascular, cerebral vascular or peripheral vascular disorders, the compounds of this invention by virtue of their hypocholesterolemic action, prevent, arrest and/or regress atherosclerosis.

Since the concentration of insulin in blood is related to the promotion of
25 vascular cell growth and increased renal sodium retention, (in addition to the other actions e.g., promotion of glucose utilization) and these functions are known causes of hypertension, the compounds of this invention by virtue of their hypoinsulinemic action, prevent, arrest and/or regress hypertension.

Since the concentration of triglycerides in blood contributes to the overall
30 levels of blood lipids, the compounds of this invention by virtue of their triglyceride lowering activity prevent, arrest and/or regress hyperlipidemia.

Five to eight week old male C57BL/6J-ob/ob mice (obtained from Jackson Laboratory, Bar Harbor, ME) are housed five per cage under standard animal care

practices and fed standard rodent diet ad libitum. After a one week acclimation period, the animals are weighed and 25 microliters of blood are collected from the retro-orbital sinus prior to any treatment. The blood sample is immediately diluted 1:5 with saline containing 0.025% sodium heparin, and held on ice for plasma glucose analysis. Animals are assigned to treatment groups so that each group has a similar mean for plasma glucose concentration. The compound to be tested is administered by oral gavage as an about 0.02% to 2.0% solution (weight/volume (w/v)) in either 1) 10% DMSO/0.1% Pluronic® P105 Block Copolymer Surfactant (BASF Corporation, Parsippany, NJ) in 0.1% saline without pH adjustment or 2) 0.25% w/v methylcellulose in water without pH adjustment. Single daily dosing (s.i.d.) or twice daily dosing (b.i.d.) is maintained for 1 to 15 days. Control mice receive the 10% DMSO/0.1% Pluronic® P105 in 0.1% saline without pH adjustment or the 0.25% w/v methylcellulose in water without pH adjustment only.

Three hours after the last dose is administered, the animals are sacrificed by decapitation and trunk blood is collected into 0.5 mL serum separator tubes containing 3.6 mg of a 1:1 weight/weight sodium fluoride: potassium oxalate mixture. The freshly collected samples are centrifuged for two minutes at 10,000 x g at room temperature, and the serum supernatant is transferred and diluted 1:1 volume/volume with a 1TIU/mL aprotinin solution in 0.1% saline without pH adjustment.

The diluted serum samples are then stored at -80°C until analysis. The thawed, diluted serum samples are analyzed for insulin, triglycerides, and cholesterol levels. Serum insulin concentration is determined using Equate® RIA INSULIN kits (double antibody method; as specified by the manufacturer) purchased from Binax, South Portland, ME. The inter assay coefficient of variation is $\leq 10\%$. Serum triglycerides are determined using the Abbott VP™ and VP Super System® Autoanalyzer (Abbott Laboratories, Irving, TX), using the A-Gent™ Triglycerides Test reagent system (Abbott Laboratories, Diagnostics Division, Irving, TX) (lipase-coupled enzyme method; a modification of the method of Sampson, et al., Clinical Chemistry 21, 1983 (1975)). Serum total cholesterol levels are determined using the Abbott VP™ and VP Super System® Autoanalyzer (Abbott Laboratories, Irving, TX), and A-Gent™ Cholesterol Test reagent system (cholesterol esterase-coupled enzyme method; a modification of the method of Allain, et al. Clinical Chemistry 20, 470

(1974)) using a 100 and 300 mg/dL standards. Serum insulin, triglycerides, and total cholesterol levels are then calculated by the equations,

$$\text{Serum insulin } (\mu\text{U/mL}) = \text{Sample value} \times 2$$

$$\text{Serum triglycerides (mg/dL)} = \text{Sample value} \times 2$$

5
$$\text{Serum total cholesterol (mg/dL)} = \text{Sample value} \times 2$$

where 2 is the dilution factor.

The animals dosed with vehicle maintain substantially unchanged, elevated serum insulin (e.g. 225 $\mu\text{U/mL}$), serum triglycerides (e.g. 225 mg/dl), and serum total cholesterol (e.g. 160 mg/dL) levels, while animals treated with test compounds of this
10 invention generally display reduced serum insulin, triglycerides, and total cholesterol levels. The serum insulin, triglycerides, and total cholesterol lowering activity of the test compounds are determined by statistical analysis (unpaired t-test) of the mean serum insulin, triglycerides, or total cholesterol concentration between the test compound group and the vehicle-treated control group.

15 Activity in providing protection from damage to heart tissue for the compounds of this invention can be demonstrated in vitro along the lines presented in Butwell et al., Am. J. Physiol., 264, H1884-H1889, 1993 and Allard et al., Am. J. Physio., 1994, 267, H66-H74. Experiments are performed using an isovolumic isolated rat heart preparation, essentially as described in the above-referenced article. Normal male
20 Sprague-Dawley rats, male Sprague-Dawley rats treated to possess cardiac hypertrophy by an aortic banding operation, acutely diabetic male BB/W rats, or non-diabetic BB/W age matched control rats are pretreated with heparin (1000 u, i.p.), followed by pentobarbital (65 mg/kg, i.p.). After deep anesthesia is achieved as determined by the absence of a foot reflex, the heart is rapidly excised and placed into iced saline. The
25 heart is retrogradely perfused through the aorta within 2 minutes. Heart rate and ventricular pressure are determined using a latex balloon in the left ventricle with high pressure tubing connected to a pressure transducer. The heart is perfused with a perfusate solution consisting of (mM) NaCl 118, KCl 4.7, CaCl_2 1.2, MgCl_2 1.2, NaHCO_3 25, glucose 11. The perfusion apparatus is tightly temperature-controlled with heated
30 baths used for the perfusate and for the water jacketing around the perfusion tubing to maintain heart temperature at 37°C. Oxygenation of the perfusate is provided by a pediatric hollow fiber oxygenator (Capiax, Terumo Corp., Tokyo, Japan) immediately proximal to the heart. Hearts are exposed to perfusion solution \pm test compound for

about 10 minutes or more, followed by 20 minutes of global ischemia and 60 minutes of reperfusion in the absence of the test compound. The heart beats of the control and test compound treated hearts are compared in the period following ischemia. The left ventricular pressure of the control and test compound treated hearts are compared in the period following ischemia. At the end of the experiment, hearts are also perfused and stained to determine the ratio of infarct area relative to the area at risk (%IA/AAR) as described below.

The therapeutic effects of the compounds of this invention in preventing heart tissue damage otherwise resulting from an ischemic insult can also be demonstrated in vivo along lines presented in Liu et al., *Circulation*, Vol. 84, No. 1, (July 1991), as described specifically herein. The in vivo assay tests the cardioprotection of the test compound relative to the control group which receives saline vehicle. As background information, it is noted that brief periods of myocardial ischemia followed by coronary artery reperfusion protects the heart from subsequent severe myocardial ischemia (Murry et al., *Circulation* 74:1124-1136, 1986). Cardioprotection, as indicated by a reduction in infarcted myocardium, can be induced pharmacologically using intravenously administered adenosine receptor agonists in intact, anesthetized rabbits studied as an in situ model of myocardial ischemic preconditioning (Liu et al., *Circulation* 84:350-356, 1991). The in vivo assay tests whether compounds can pharmacologically induce cardioprotection, i.e., reduced myocardial infarct size, when parenterally administered to intact, anesthetized rabbits. The effects of the compounds of this invention can be compared to ischemic preconditioning using the A1 adenosine agonist, N⁶-1-(phenyl-2R-isopropyl) adenosine (PIA) that has been shown to pharmacologically induce cardioprotection in intact anesthetized rabbits studied in situ (Liu et al., *Circulation* 84:350-356, 1991). The exact methodology is described below.

Surgery: New Zealand White male rabbits (3-4 kg) are anesthetized with sodium pentobarbital (30 mg/kg, i.v.). A tracheotomy is performed via a ventral midline cervical incision and the rabbits are ventilated with 100% oxygen using a positive pressure ventilator. Catheters are placed in the left jugular vein for drug administration and in the left carotid artery for blood pressure measurements. The hearts are then exposed through a left thoracotomy and a snare (00 silk) placed around a prominent branch of the left coronary artery. Ischemia is induced by pulling the snare tight and clamping it in place. Releasing the snare allowed the affected area to reperfuse. Myocardial

-52-

ischemia is evidenced by regional cyanosis; reperfusion was evidenced by reactive hyperemia.

Protocol: Once arterial pressure and heart rate has been stable for at least 30 minutes the experiment is started. Ischemic preconditioning is induced by twice occluding the coronary artery for 5 min followed by a 10 min reperfusion. Pharmacological preconditioning is induced by twice infusing test compound over, for example 5 minutes and allowing 10 minutes before further intervention or by infusing the adenosine agonist, PIA (0.25 mg/kg). Following ischemic preconditioning, pharmacological preconditioning or no conditioning (unconditioned, vehicle control) the artery is occluded for 30 minutes and then reperfused for two hours to induce myocardial infarction. The test compound and PIA are dissolved in saline or other suitable vehicle and delivered at 1 to 5 ml/kg, respectively.

Staining (Liu et al., Circulation 84:350-356, 1991): At the end of the 2 hour reperfusion period, the hearts are quickly removed, hung on a Langendorff apparatus, and flushed for 1 minute with normal saline heated to body temperature (38°C). The silk suture used as the snare is then tied tightly to reocclude the artery and a 0.5% suspension of fluorescent particles (1-10 μ m) is infused with the perfusate to stain all of the myocardium except the area at risk (nonfluorescent ventricle). The hearts are then quickly frozen and stored overnight at -20°C. On the following day, the hearts are cut into 2 mm slices and stained with 1% triphenyl tetrazolium chloride (TTC). Since TTC reacts with living tissue, this stain differentiates between living (red stained) tissue, and dead tissue (unstained infarcted tissue). The infarcted area (no stain) and the area at risk (no fluorescent particles) are calculated for each slice of left ventricle using a pre-calibrated image analyzer. To normalize the ischemic injury for differences in the area at risk between hearts, the data is expressed as the ratio of infarct area vs. area at risk (%IA/AAR). All data is expressed as Mean \pm SEM and compared statistically using single factor ANOVA or unpaired t-test. Significance is considered as $p < 0.05$.

Administration of the compounds of this invention can be via any method which delivers a compound of this invention preferentially to the liver and/or cardiac tissues. These methods include oral routes, parenteral, intraduodenal routes, etc. Generally, the compounds of the present invention are administered in single (e.g., once daily) or multiple doses.

-53-

However, the amount and timing of compound(s) administered will, of course, be dependent on the particular disease/condition being treated, the subject being treated, on the severity of the affliction, on the manner of administration and on the judgment of the prescribing physician. Thus, because of patient to patient variability, the dosages given below are a guideline and the physician may titrate doses of the drug to achieve the activity (e.g., glucose lowering activity) that the physician considers appropriate for the patient. In considering the degree of activity desired, the physician must balance a variety of factors such as starting level, other risk (cardiovascular) factors, presence of preexisting disease, and age of the patient and the patient's motivation.

In general an effective dosage for the activities of this invention, for example, the blood glucose, triglycerides, and cholesterol lowering activities and hyperinsulinemia reversing activities of the compounds of this invention is in the range of 0.005 to 50 mg/kg/day, preferably 0.01 to 25 mg/kg/day and most preferably 0.1 to 15 mg/kg/day.

Generally, the compounds of this invention are administered orally, but parenteral administration (e.g., intravenous, intramuscular, subcutaneous or intramedullary) may be utilized, for example, where oral administration is inappropriate for the instant target or where the patient is unable to ingest the drug. Topical administration may also be indicated, for example, where the patient is suffering from gastrointestinal disorders or whenever the medication is best applied to the surface of a tissue or organ as determined by the attending physician.

The compounds of the present invention are generally administered in the form of a pharmaceutical composition comprising at least one of the compounds of this invention together with a pharmaceutically acceptable vehicle or diluent. Thus, the compounds of this invention can be administered individually or together in any conventional oral, parenteral or transdermal dosage form.

For oral administration a pharmaceutical composition can take the form of solutions, suspensions, tablets, pills, capsules, powders, and the like. Tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate are employed along with various disintegrants such as starch and preferably potato or tapioca starch and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia.

Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type are also employed as fillers in soft and hard-filled gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the compounds of this invention can be combined with various sweetening agents, flavoring agents, coloring agents, emulsifying agents and/or suspending agents, as well as such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

10 For purposes of parenteral administration, solutions in sesame or peanut oil or in aqueous propylene glycol can be employed, as well as sterile aqueous solutions of the corresponding water-soluble salts. Such aqueous solutions may be suitably buffered, if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose. These aqueous solutions are especially suitable for

15 intravenous, intramuscular, subcutaneous and intraperitoneal injection purposes. In this connection, the sterile aqueous media employed are all readily obtainable by standard techniques well-known to those skilled in the art.

For purposes of transdermal (e.g., topical) administration, dilute sterile, aqueous or partially aqueous solutions (usually in about 0.1% to 5% concentration),

20 otherwise similar to the above parenteral solutions, are prepared.

Methods of preparing various pharmaceutical compositions with a certain amount of active ingredient are known, or will be apparent in light of this disclosure, to those skilled in this art. For examples, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easter, Pa., 15th Edition (1975).

25 Pharmaceutical compositions according to the invention may contain 0.1%-95% of the compound(s) of this invention, preferably 1%-70%. In any event, the composition or formulation to be administered will contain a quantity of a compound(s) according to the invention in an amount effective to treat the disease/condition of the subject being treated, i.e., a glycogen phosphorylase

30 dependent disease/condition.

General Experimental Procedures for Examples 1-99 and 166-172

NMR spectra were recorded on a Varian XL-300 (Varian Co., Palo Alto, California) or Bruker AM-300 spectrometer (Bruker Co., Billerica, Massachusetts) at

about 23°C at 300 MHz for proton and 75.4 MHz for carbon nuclei. Chemical shifts are expressed in parts per million downfield from trimethylsilane. Resonances designated as exchangeable did not appear in a separate NMR experiment where the sample was shaken with several drops of D₂O in the same solvent. FAB-MS

- 5 spectra were obtained on a VG70-2505 spectrometer (V4 analytical LTD., Wythanshaw, Manchester, U.K.) using a liquid matrix consisting of 3:1 dithiothreitol/dithioerythritol. Thermospray MS (TSPMS) were obtained on a Fisons Trio-1000 spectrometer (Fisons Co., Valencia, California) using ammonia ionization. Chemical ionization mass spectra were obtained on a Hewlett-Packard 5989
- 10 instrument (Hewlett-Packard Co., Palo Alto, California) (ammonia ionization, PBMS). Where the intensity of chlorine or bromine-containing ions are described the expected intensity ratio was observed (approximately 3:1 for ³⁵Cl/³⁷Cl-containing ions) and 1:1 for ⁷⁹Br/⁸¹Br-containing ions) and the intensity of only the lower mass ion is given.

- 15 HPLC was performed with 214 nM detection on a 250 x 4.6 mm Rainin Microsorb C-18 column (Rainin Co., Woburn, Massachusetts) eluted isocratically by a two-pump/mixer system supplying the indicated mixture of acetonitrile and aqueous pH 2.1 (with H₃PO₄) 0.1M KH₂PO₄, respectively, at 1.5 mL/min. Samples were injected in a 1:1 mixture of acetonitrile and pH 7.0 phosphate buffer (0.025M
- 20 in each Na₂HPO₄ and KH₂PO₄). Percent purities refer to percent of total integrated area usually over a 10 to 15 minute run. Melting points are uncorrected and were determined on a Buchi 510 melting point apparatus (Buchi Laboratorums-Technik Ag., Flawil, Switzerland) where melting points of 120.5-122°C for benzoic acid and 237.5-240.5°C for p-chlorobenzoic acid (Aldrich 99+% grades) were obtained.

- 25 Column chromatography was performed with Amicon silica gel (30 µm, 60A pore size) (Amicon D Vision, W.R. Grace & Co., Beverly, Mass.) in glass columns under low nitrogen pressure. Unless otherwise specified, reagents were used as obtained from commercial sources. Dimethylformamide, 2-propanol, tetrahydrofuran, and dichloromethane used as reaction solvents were the anhydrous grade supplied by
- 30 Aldrich Chemical Company (Milwaukee, Wisconsin). Microanalyses were performed by Schwarzkopf Microanalytical Laboratory, Woodside, NY. The terms "concentrated" and coevaporated refer to removal of solvent at water aspirator pressure on a rotary evaporator with a bath temperature of less than 45°C.

-56-

Reactions conducted at "0-20°C" or "0-25°C" were conducted with initial cooling of the vessel in an insulated ice bath which was allowed to warm to room temperature over several hours. The abbreviation "min" and "h" stand for "minutes" and "hours" respectively.

5

Procedure A (Peptide Coupling Using DEC)

An 0.1-0.7 M solution of the primary amine (1.0 equiv, or a primary amine hydrochloride and 1.0 to 1.3 equivalents of triethylamine per equiv HCl) in dichloromethane (unless other solvent specified), is treated sequentially at 25°C with 0.95 to 1.2 equivalent of the specified carboxylic acid, 1.2 to 1.8 equivalent
10 hydroxybenzotriazole hydrate (usually 1.5 equivalent relative to the carboxylic acid), and 0.95-1.2 equivalent (corresponding in mole ratio to the carboxylic acid) 1-(3-dimethylaminopropyl)3-ethylcarbodiimide hydrochloride (DEC) and the mixture is stirred for 14 to 20 hours. (See Note 1 below). The mixture is diluted with ethyl acetate, washed 2 to 3 times with 1 or 2N NaOH, 2 to 3 times with 1 or 2N HCl (Note 2), the
15 organic layer dried over MgSO₄, and concentrated giving crude product which is purified by chromatography on silica gel, trituration, or recrystallization, as specified using the specified solvents. Purified products were analyzed by RP-HPLC and found to be of greater than 95% purity unless otherwise noted. Reactions conducted at 0 to 25°C were conducted with initial cooling of the vessel in an insulated ice bath which
20 was allowed to warm to room temperature over several hours.

Note 1: On larger scale couplings (>50 mL solvent) the mixture was concentrated at this point and the residue dissolved in ethyl acetate.

Note 2: If the product contained ionizable amine functionality the acid wash was omitted. Exceptions in the use of Procedure A are noted individually (where
25 appropriate below) usually in parentheses, immediately following mention of Procedure A.

Example 1

(2S)-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid methyl ester

L-Phenylalanine methyl ester hydrochloride (77.0 mmol) and 5-chloro-1H-indole-
30 2-carboxylic acid (77 mmol) were coupled according to procedure A (0 - 25 °C) and the product purified by chromatography on silica gel in 10 and 20 % ethyl acetate-hexanes giving the title substance as an off-white solid (22.12 g, 81 %): mp 156 - 157 °C; HPLC (60/40) 9.5 minutes (98 %); PBMS 357/359 (MH⁺, 100 %).

-57-

^1H NMR (CDCl_3) δ 9.40 (br, 1H), 7.60 (d, 1H, $J = \text{ca } 1 \text{ Hz}$), 7.35 (d, 1H, $J = 8.9 \text{ Hz}$), 7.3 - 7.2 (m, 4H), 7.13 (m, 2H), 6.74 (d, 1H, $J = 1.7 \text{ Hz}$), 6.62 (d, 1H, $J = 7.5 \text{ Hz}$), 5.11 (m, 1H), 3.77 (s, 3H), 3.26 (m, 2H);

Anal. Calcd for $\text{C}_{19}\text{H}_{17}\text{ClN}_2\text{O}_3$: C, 63.96; H, 4.80; N, 7.85.

5 Found: C, 64.24; H, 4.84; N, 8.02.

Example 2

2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid

Aqueous 2M LiOH (33.10 ml) was added to a solution of (2S)-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid methyl ester (21.47 g, 60 mmol) in THF (140 ml) at 0 - 5 °C. After 0.5 hour, the mixture was partially concentrated, acidified to pH 1 - 2 with 6N HCl, concentrated to dryness, and the solids washed with water and then ether to yield a colorless solid (18.78 g, 91%): mp 248 - 255°C; HPLC (60/40) 5.21 minutes (98%); TSPMS 343/345 (MH^+ , 100%).

^1H NMR ($\text{DMSO}-d_6$) δ 12.85 (br, 1H), 11.75 (d, 1H, $J = < 1 \text{ Hz}$), 8.84 (d, 1H, $J = 8.4 \text{ Hz}$), 7.35 - 7.14 (m, 7H), 4.65 (m, 1H), 3.20 (A of AB, 1H, $J = 4.5, 13.9 \text{ Hz}$), 3.07 (B of AB, 1H, $J = 10.8, 13.8 \text{ Hz}$);

Anal. Calcd for $\text{C}_{18}\text{H}_{15}\text{ClN}_2\text{O}_3$: C, 63.07; H, 4.41; N, 8.17.

Found: C, 62.90; H, 4.60; N, 8.04.

Example 3

[(5-Chloro-1H-indole-2-carbonyl)-amino]-acetic acid methyl ester

Glycine methyl ester hydrochloride (50 mmol) and 5-chloro-1H-indole-2-carboxylic acid (50 mmol) were coupled according to procedure A, substituting the following workup: the reaction mixture was stirred in ethyl acetate (250 mL), hexanes (50 mL) and 1N NaOH (50 mL) and the suspension was filtered. The solid was washed with 1N NaOH, 1N HCl, with water, ethyl acetate, and dried: Yield 11.5 g, 86%; mp 252-254°C with decomposition;

^1H NMR ($\text{DMSO}-d_6$) δ 11.87 (br, 1H), 9.05 (t, 1H, $J = 6.0 \text{ Hz}$), 7.72 (d, 1H, $J = 2.0 \text{ Hz}$), 7.45 (d, 1H, $J = 8.7 \text{ Hz}$), 7.19 (dd, 1H, $J = 2.0, 8.7 \text{ Hz}$), 4.05 (d, 2H, $J = 6.0 \text{ Hz}$), 3.91 (s, 3H);

30 Anal. Calcd for $\text{C}_{12}\text{H}_{11}\text{ClN}_2\text{O}_3$: C, 54.05; H, 4.16; N, 10.50.

Found: C, 54.11; H, 4.23; N, 10.56.

-58-

Example 4

[(5-Chloro-1H-indole-2-carbonyl)-amino]-acetic acid

1N NaOH (35 ml) was added to a suspension of [(5-chloro-1H-indole-2-carbonyl)-amino]-acetic acid methyl ester (8.0 g, 30 mmol) in THF (100 ml) and the
5 resulting mixture stirred for 18 hours at 25 °C. The solution was acidified with 6N HCl (7 mL), the mixture concentrated, the solids suspended in water, filtered, and washed with water (7.42 g, 98 %): HPLC (60/40) 2.89 minutes (100 %;)

¹H NMR (300 MHz, DMSO-d₆) δ 12.68 (br, 1H), 11.85 (br, 1H), 8.95 (t, 1H, J = 5.9 Hz), 7.72 (d, 1H, J = 2.0 Hz), 7.44 (d, 1H, J = 8.7 Hz), 7.19 (dd, 1H, J = 2.0, 8.7 Hz), 7.14
10 (d, 1H, J = <2 Hz), 3.96 (d, 2H, J = 5.9 Hz)

Anal. Calcd for C₁₁H₉N₂O₃Cl: C, 52.29; H, 3.59; N, 11.09.

Found: C, 52.26; H, 3.73; N, 11.20.

Example 5

5-Chloro-1H-indole-2-carboxylic acid [2-((3R)-hydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide

3-Pyrrolidinol (1.25 mmol) and [(5-chloro-1H-indole-2-carbonyl)-amino]-acetic acid (1.19 mmol) were coupled according to Procedure A with the following workup: the reaction mixture was diluted with ethyl acetate and 2 N HCl, stirred 1 hour, the mixture filtered, and the resulting solids washed successively with 2 N HCl, 2 N NaOH,
20 2 N HCl, dried, triturated with 1:1 ether / hexanes and dried, giving an off-white solid: Yield 280 mg, 73 %; HPLC (60/40) 4.66 minutes (96 %); PBMS 322/324 (MH⁺, 100 %).

¹H NMR (DMSO-d₆) δ 11.87 (br, 1H), 8.71 (q, 1H), 7.71 (d, 1H, J = 2.1 Hz), 7.45 (d, 1H, J = 8.8 Hz), 7.19 (dd, 1H, J = 3.1, 8.8 Hz), 7.16 (s, 1H), 5.07 (d, 0.5H, J = 3.6 Hz),
25 4.97 (d, 0.5H, J = 3.1 Hz), 4.35 (m, 1H), 4.27 (m, 1H), 4.10 (t, 1H), 4.03 (d, 1H), 3.59 (m, 1H), 3.49 - 3.27 (m, 2H), 2.04 - 1.79 (m, 2H).

Example 6

5-Chloro-1H-indole-2-carboxylic acid [2-(Cis-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide

30 (3R,4S)-3,4-Dihydroxypyrrolidine hydrochloride (the cis, or meso isomer, 1.79 mmol) and [(5-chloro-1H-indole-2-carbonyl)-amino]-acetic acid (0.85 mmol) were coupled according to procedure A (1:1 CH₂Cl₂ / DMF reaction solvent) with the following workup: the reaction mixture was concentrated, the residue suspended in 10

-59-

ml EtOAc and 10 ml 2 N NaOH, the solids filtered and washed successively with aqueous 1 N NaOH, EtOAc, aqueous 1 N HCl, H₂O, and ether. This washing sequence was repeated and the resulting solids were suspended in EtOAc, stirred for 1 hour, filtered and dried: Yield 252 mg, 88 %; HPLC (60/40) 2.33 minutes (93 %);

5 TSPMS 338/340 (MH⁺, 100 %);

¹H NMR (DMSO-d₆) δ 11.82 (s, 1H), 8.72 (t, 1H), 7.73 (d, 1H), 7.45 (d, 1H), 7.20 (dd, 1H), 7.15 (s, 1H), 5.05 (d, 1H), 4.98 (d, 1H), 4.10 (m, 1H), 4.03 (m, 3H), 3.68 (dd, 1H), 3.42 (dd, 1H), 3.33 (dd, 1H), 3.23 (dd, 1H).

Example 7

10 5-Chloro-1H-indole-2-carboxylic acid [2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide
4-Hydroxypiperidine (0.83 mmol) and [(5-chloro-1H-indole-2-carbonyl)-amino]-acetic acid (0.8 mmol) were coupled according to procedure A (dimethylformamide-dichloromethane reaction solvent) with the following workup: the reaction mixture was stirred with ethyl acetate and aqueous 2 N HCl, the resulting suspension filtered and
15 the collected solid washed successively with aqueous 2 N HCl, aqueous 2 N NaOH, ether and dried : Yield 180 mg, 68 %; TSPMS 336/338 (MH⁺, 100 %);
¹H NMR (DMSO-d₆) δ 11.84 (br, 1H), 8.68 (br, 1H), 7.71 (d, 1H), 7.43 (d, 1H), 7.17 (dd, 1H), 7.14 (s, 1H), 4.80 (br, 1H), 4.15 (m, 2H), 3.91 (m, 1H), 3.72 (m, 2H), 3.20 (m, 1H), 3.05 (m, 1H), 1.75 (m, 2H), 1.48 (m, 1H), 1.38 (m, 1H).

20

Example 8

5-Chloro-1H-indole-2-carboxylic acid [1-benzyl-2-(3-hydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide

Racemic 3-pyrrolidinol (2.0 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (1 mmol) were coupled according to procedure A (0 - 25 °C
25 reaction temperature, washing first with acid then base), and the product purified by column chromatography on silica gel eluted with 0.5 - 16% ethanol in dichloromethane to give a colorless foam: Yield 260 mg, 63 %; HPLC (60/40) 100 %, 3.86 minutes; PBMS 412/414 (MH⁺, 100 %);
Anal. Calcd for C₂₂H₂₂ClN₃O₃ + 0.2 H₂O: C, 63.60; H, 5.43; N, 10.11.
30 Found: C, 63.90; H, 5.93; N, 10.11.

Example 9

5-Chloro-1H-indole-2-carboxylic acid (1-diethylcarbamoyl-2-phenyl-ethyl)-amide

Diethylamine (1.2 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (0.6 mmol) were coupled according to procedure A (0 - 25 °C for 5 days) substituting the following workup: the crude product was suspended in 1:1 chloroform / dichloromethane, sonicated, filtered to remove solids, concentrated, and the residue purified by column chromatography on silica gel eluted with 10, 20 and 30 % ethyl acetate in hexanes: Yield 14 mg, 6 %; HPLC (60/40) 8.88 minutes (98 %); PBMS 398/400 (MH+, 100 %);

10 ¹H NMR (CDCl₃) δ 9.31 (br, 1H), 7.61 (d, 1H), 7.32 (d, 1H, J = 8.7 Hz), 7.28-7.18 (m, 7H), 6.87 (d, 1H, J = 1.4 Hz), 5.26 (m, 1H), 3.6 (m, 1-1.5H), 3.2-2.9 (m, 4.5-5H), 1.07 (t, 3H, J = 7.2 Hz), 1.02 (t, 3H, J = 7.2 Hz).

Anal. Calcd for C₂₂H₂₄ClN₃O₂ + 0.25 H₂O: C, 65.66; H, 6.14; N, 10.44.
Found: C, 65.61; H, 6.20; N, 10.11.

15

Example 10

4-{2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl}-piperazine-1-carboxylic acid tert-butyl ester

1-Piperazinecarboxylic acid t-butyl ester (1.2 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (0.6 mmol) were coupled according to

20 procedure A (0 - 25 °C reaction temperature, reaction time 4 days, extraction with acid first, then base), and the crude product purified by column chromatography on silica gel eluted with 30 % ethyl acetate in hexanes to give a colorless foam: Yield 290 mg, 95 %; HPLC (70/30) 6.23 min (99 %); PBMS 512/514 (MH+, 100 %);

¹H NMR (CDCl₃) δ 9.32 (br, 1H), 7.60 (d, 1H, J = 1.9 Hz), 7.32 (d, 1H, J = 8.7 Hz),

25 7.3-7.15 (m, ca. 7H), 6.87 (d, 1H, J = 1.5 Hz), 5.33 (m, 1H), 3.65 - 2.9 (overlapping m, 9H), 2.70 (m, 1H), 1.43 (s, 9H).

Anal. Calcd for C₂₇H₃₁ClN₄O₄: C, 63.46; H, 6.11; N, 10.96.
Found: C, 63.33; H, 5.97; N, 10.97.

Example 11

30

5-Chloro-1H-indole-2-carboxylic acid [1-benzyl-2-(4-methylamino-piperidin-1-yl)-2-oxo-ethyl]-amide

Dimethylamine hydrochloride (1.1 mmol), sodium acetate (2.1 mmol), activated 3Å molecular sieves, and sodium cyanoborohydride (0.25 mmol) were added in this

-61-

order to 5-chloro-1H-indole-2-carboxylic acid [1-benzyl-2-oxo-2-(4-oxo-piperidin-1-yl)-ethyl]-amide (0.21 mmol) in methanol (2 mL) at 0 °C. After 18 hours, the mixture was concentrated, the residue taken up in ethyl acetate, the resulting solution washed with 2N NaOH and brine, dried with Na₂SO₄ and concentrated. The product was purified by chromatography on silica gel eluted with 1-8% ethanol in dichloromethane containing 0.5 % NH₄OH followed by trituration with ether: Yield 82 %; HPLC (60/40) 2.79 minutes (98 %); PBMS 439/441 (MH⁺, 100 %);

¹H NMR (DMSO-d₆) δ 11.75 (br, 1H), 8.94 (d, 0.5H, J = 8.8 Hz), 8.90 (d, 0.5H), 7.71 (d, 1H, J = 1.8 Hz), 7.40 (d, 1H, J = 8.7 Hz), 7.31 - 7.20 (m, 6-7H), 7.17 (dd, 1H, J = 2.1, 8.7 Hz), 5.15 (m, 1H), 4.22 (m, 0.5H), 4.08 (m, 0.5H), 3.96 (m, 0.5H), 3.85 (m, 0.5H), 3.2 - 2.9 (m, 4H), 2.78 (m, 0.5H), 2.72 (m, 0.5H), 2.25 (s, 1.5H), 2.24 (s, 1.5H), 1.75 (m, 2H), 1.3 - 0.8 (m, 2H).

Anal. Calcd for C₂₄H₂₇ClN₄O₂ + 1.0 H₂O: C, 63.08; H, 6.40; N, 12.26.

Found: C, 63.18; H, 6.16; N, 12.46.

15

Example 12

5-Chloro-1H-indole-2-carboxylic acid (1-benzyl-2-morpholin-4-yl-2-oxo-ethyl)-amide

Morpholine (0.33mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (0.30 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 48 hour reaction time). The crude product was chromatographed on silica gel eluted with 1:1 ethyl acetate / hexanes, the desired fractions concentrated, the residue dissolved in chloroform and methanol and the resulting solution stirred 18 hours with approx. 128 mg dimethylaminopyridine-polystyrene resin (Fluka Chemical Co.). The solution was filtered, concentrated and the residue trituated with ether: Yield, 51 %; HPLC (60/40) 5.92 min (98 %);

25 PBMS 412/414 (MH⁺, 100%);

¹H NMR (DMSO-d₆) δ 11.75 (br, 1H), 8.95 (d, 1H), 7.72 (d, 1H), 7.39 (d, 1H, J = 8.7 Hz), 7.35 - 7.15 (m, 7H), 5.13 (m, 1H), 3.65-3.10 (m, 8H), 3.05 (m, 2H).

Anal. Calcd for C₂₂H₂₂ClN₃O₃ + 0.33H₂O: C, 63.23; H, 5.47; N, 10.06.

Found: C, 63.28; H, 5.32; N, 10.10.

30

Example 13

5-Chloro-1H-indole-2-carboxylic acid (1-butylcarbamoyl-2-phenyl-ethyl)-amide

n-Butyl amine (0.66 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (0.60 mmol) were coupled according to procedure A (0 - 25 °C

-62-

- reaction temperature). The crude product was dissolved in chloroform and methanol and the resulting solution stirred 18 hours with 50 mg dimethylaminopyridine-poly-styrene resin (Fluka Chemical Co.), the solution filtered, concentrated and the solids triturated with ether: Yield 83 %; HPLC (60/40) 8.88 minutes (92 %); mp 192 - 193 °C; TSPMS 398/400 (MH+, 100 %);
- ¹H NMR (DMSO-*d*₆) δ 11.71 (br, 1H), 8.70 (d, 1H, J = 8.3 Hz), 8.10 (t, 1H), 7.72 (d, 1H, J = 2.0 Hz), 7.39 (d, 1H, J = 8.7 Hz), 7.35 - 7.15 (m, 7H), 4.70 (m, 1H), 3.13-2.93 (m, 4H), 1.38 (m, 2H), 1.25 (m, 2H), 0.86 (t, 3H, J = 7.2 Hz);
- Anal. Calcd for C₂₂H₂₄ClN₃O₂: C, 66.41; H, 6.08; N, 10.56.
- 10 Found: C, 66.15; H, 6.04; N, 10.52.

Example 14

5-Chloro-1H-indole-2-carboxylic acid [1-benzyl-2-oxo-2-(4-oxo-piperidin-1-yl)-ethyl]-amide

- 4-Piperidone monohydrate (2.0 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (1.0 mmol) were coupled according to procedure A (0 - 25 °C reaction temperature) substituting the following workup: the reaction mixture was diluted with ethyl acetate, the resulting solution washed with 2 N NaOH and 2 N HCl, the suspension filtered and the solids dried: Yield 111 mg, 26 %; HPLC (60/40) 8.88 minutes (92 %); PBMS 424/426 (MH+, 100 %); mp 258 - 261 °C; PBMS 424/426
- 20 (MH+, 100 %);
- ¹H NMR (DMSO-*d*₆) δ 11.75 (br, 1H), 9.03 (d, 1H, J = 8.1 Hz), 7.72 (d, 1H, J = 1.9 Hz), 7.39 (d, 1H, J = 8.7 Hz), 7.4 - 7.15 (m, 7H), 5.20 (m, 1H, J = 8.2 Hz), 3.88 (m, 1H), 3.73 (m, 3H), 3.1 (m, 3H), 2.5 - 2.22 (m, 3H), 2.05 (m, 1H).
- Anal. Calcd for C₂₃H₂₂ClN₃O₃ + 0.75 H₂O: C, 63.16; H, 5.42; N, 9.61.
- 25 Found: C, 63.11; H, 5.15; N, 9.53.

Example 15

5-Chloro-1H-indole-2-carboxylic acid (1-benzyl-2-oxo-2-pyrrolidin-1-yl-ethyl)-amide

- Pyrrolidine (0.35 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (0.31 mmol) were coupled according to procedure A (0 - 25 °C reaction
- 30 temperature, 140 hour reaction time) and the crude product triturated with ether: Yield 89 mg, 71 %; HPLC (70/30) 7.57 minutes (98 %); PBMS 396/398 (MH+, 100 / 80 %);
- Anal. Calcd for C₂₂H₂₂ClN₃O₂ + 0.33 H₂O: C, 65.75; H, 5.68; N, 10.48.
- Found: C, 65.56; H, 5.81; N, 10.44.

-63-

Example 16

5-Chloro-1H-indole-2-carboxylic acid [1-[(3-dimethylamino-propyl)-methyl-carbamoyl]-2-phenyl-ethyl]-amide

N,N,N'-Trimethyl-1,3-diaminopropane (0.31 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (0.28 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 120 hour reaction time) and the product purified by chromatography on silica gel eluted with 1 - 8 % ethanol in dichloromethane containing 0.5 % ammonium hydroxide: Yield 86 mg, 69 %; HPLC (40/60) 7.57 minutes (>99 %); mp 187 - 190.5 °C; TSPMS 441/443 (MH+, 100 %);

Anal. Calcd for $C_{24}H_{29}ClN_4O_2 + 0.25 H_2O$: C, 64.71; H, 6.68; N, 12.58.

Found: C, 64.73; H, 6.94; N, 12.86.

Example 17

5-Chloro-1H-indole-2-carboxylic acid [1-(3-morpholin-4-yl-propylcarbamoyl)-2-phenyl-ethyl]-amide

4-(3-Aminopropyl)morpholine (0.34 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (0.30 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) substituting the following workup: the reaction was diluted with ethyl acetate, the resulting solution washed three times with 2 N NaOH and once with brine, dried over Na_2SO_4 , and concentrated. The residue was stirred under ether for 1 hour, the solid filtered and dried: Yield 125 mg, 87 %; HPLC (60/40) 2.85 minutes (98 %); PBMS 469/471 (MH+, 100 / 90 %);

Anal. Calcd for $C_{25}H_{29}ClN_4O_3 + 0.25 H_2O$: C, 63.42; H, 6.28; N, 11.83.

Found: C, 63.31; H, 6.57; N, 12.04.

Example 18

5-Chloro-1H-indole-2-carboxylic acid (1-dimethylcarbamoyl-2-phenyl-ethyl)-amide

Dimethylamine hydrochloride (0.96 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (0.90 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 60 hour reaction time, washed first with acid, then base), and the resulting solid triturated with ether: Yield 320 mg, 99 %; HPLC (60/40) 5.87 minutes (100 %);

mp 224 - 225 °C; PBMS 370/372 (MH+, 100 %).

A sample was recrystallized from hot ethyl acetate for analysis (mp 224 - 225 °C).

Anal. Calcd for $C_{20}H_{20}ClN_3O_2 + 0.5 C_4H_8O_2$: C, 63.80; H, 5.84; N, 10.15.

Found: C, 63.81; H, 5.80; N, 10.21.

Example 19

5-Chloro-1H-indole-2-carboxylic acid [2-((3R,4R)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide

5 (3R,4R)-3,4-dihydroxypyrrolidine (from unnatural tartaric acid by the procedure described in US Patent No. 4634775, 1.0 mmol) and [(5-chloro-1H-indole-2-carbonyl)-amino]-acetic acid (1.1 mmol) were coupled according to Procedure A (dimethylformamide reaction solvent) substituting the following workup: the reaction mixture was concentrated, diluted with 20 mL ethyl acetate and 20 mL 2 N NaOH, the suspension
10 stirred 0.5 hours, filtered, and the resulting solids washed successively with 2N NaOH, water, 1N HCl and ethyl acetate: Yield 77 %; HPLC (40/60) 10.7 minutes (99 %); TSPMS 338/340 (MH+, 100 %);
¹H NMR (DMSO-d₆) δ 11.84 (br, 1H, exchanges), 8.72 (t, 1H, exchanges), 7.72 (d, 1H, J = 1.9 Hz), 7.44 (d, 1H, J = 8.7 Hz), 7.19 (dd, 1H, J = 2.1, 8.8 Hz), 7.16 (s, 1H), 5.26
15 (d, 1H, J = 4.4 Hz, exchanges), 5.17 (d, 1H, J = 3.2 Hz, exchanges), 4.04 (m, 3H), 3.92 (m, 1H), 3.66 (dd, 1H, J = 4.0, 10.8 Hz), 3.42 - 3.28 (m, 3H).;
Anal. Calcd for C₁₅H₁₆ClN₃O₄ + 0.25 H₂O: C, 52.64; H, 4.86; N, 12.28.
Found: C, 52.61; H, 4.85; N, 12.23.

Example 20

20 5-Chloro-1H-indole-2-carboxylic acid [2-((3S,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide

(3S,4S)-3,4-Dihydroxypyrrolidine (from naturally occurring tartaric acid by the procedure described in US Patent No. 4634775, 1.0 mmol) and [(5-chloro-1H-indole-2-carbonyl)-amino]-acetic acid (1.0 mmol) were coupled according to Procedure A
25 (dimethylformamide reaction solvent) substituting the following workup: the reaction was diluted with ethyl acetate and 2N NaOH, the resulting suspension filtered, the solids washed with ethyl acetate, water and dried: Yield 135 mg, 40 %; HPLC (40/60) 7.29 minutes (98 %); TSPMS 338/340 (MH+, 100 %);
¹H NMR (DMSO-d₆) δ 12.1 (br, 1H), 8.86 (br, 1H), 7.71 (d, 1H, J = 2 Hz), 7.43 (d, 1H, J = 8.8 Hz), 7.17 (dd, 1H, J = 2, 8.8 Hz), 7.13 (s, 1H), 5.35 (br, 1H, exchanges with
30 D₂O), 5.28 (br, 1H, exchanges with D₂O), 4.03 (m, 3H), 3.92 (s, 1H), 3.66 (dd, 1H, J = 4, 11 Hz), 3.4 - 3.2 (m, 3H).
Anal. Calcd for C₁₅H₁₆ClN₃O₄ + 1.5 H₂O: C, 49.39; H, 5.25; N, 11.52.

-65-

Found: C, 49.50; H, 5.04; N, 11.27.

Example 21

5-Chloro-1H-indole-2-carboxylic acid [1-benzyl-2-(4-methoxymethoxy-piperidin-1-yl)-2-oxo-ethyl]-amide

- 5 4-Methoxymethoxy-piperidine (1.0 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (1.0 mmol) were coupled according to Procedure A and the product purified by chromatography on silica gel eluted with 1:1 ethyl acetate-hexanes: Yield 241 mg, 50 %; HPLC (60/40) 7.67 minutes (94 %); PBMS 470/472 (MH⁺, 100 %);
- 10 Anal. Calcd for C₂₅H₂₈ClN₃O₄: C, 63.89; H, 6.01; N, 8.94.
Found: C, 63.91; H, 6.00; N, 8.95.

Example 22

5-Chloro-1H-indole-2-carboxylic acid [2-phenyl-1-(2,2,6,6-tetramethyl-piperidin-4-ylcarbamoyl)-ethyl]-amide

- 15 2,2,6,6-Tetramethyl-piperidine (1.0 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (1.0 mmol) were coupled according to Procedure A. The resulting yellow foam was triturated with ether, 1:5 dichloromethane-ether. The resulting solid was dissolved in dichloromethane and the resulting solution treated with 0.20 mL 4 N HCl in dioxane. A precipitate formed which was filtered, washed with
- 20 dichloromethane and dried: Yield 220 mg, 42 %; HPLC (60/40) 3.19 minutes (96 %); PBMS 481/483 (MH⁺, 100 %);
- Anal. Calcd for C₂₇H₃₃ClN₄O₂ + HCl + 1.5 H₂O: C, 59.56; H, 6.85; N, 10.29.
Found: C, 59.30; H, 6.90; N, 10.22.

Example 23

(1-{2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl}-pyrrolidin-(3RS)-yl)-carbamic acid tert-butyl ester

- 25 Racemic Pyrrolidine-3-carbamic acid tert-butyl ester (1.0 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (1.0 mmol) were coupled according to Procedure A and the product purified by chromatography on silica gel eluted with 1:1 ethyl acetate-hexanes: Yield 302 mg, 59 %; PBMS 511/513 (MH⁺, 100 %);
- 30 Anal. Calcd for C₂₇H₃₁ClN₄O₄: C, 63.46; H, 6.11; N, 10.96.
Found: C, 63.32; H, 6.26; N, 10.89.

Example 24

5-Chloro-1H-indole-2-carboxylic acid (2-morpholin-4-yl-2-oxo-ethyl)-amide

Morpholine (1.0 mmol) and [(5-chloro-1H-indole-2-carbonyl)-amino]-acetic acid (1.0 mmol) were coupled according to Procedure A. The resulting solid was
5 suspended in ether, filtered and dried to give a beige solid: Yield, 264 mg, 71 %; HPLC (60/40) 3.28 minutes (100 %); TSPMS 322/324 (MH+, 100 %);
¹H NMR (DMSO-d₆) δ 11.85 (s, 1H), 8.68 (t, 1H), 7.72 (d, 1H, J = 2.0 Hz), 7.43 (d, 1H, J = 8.8 Hz), 7.19 (dd, 1H, J = 2.1, 8.8 Hz), 7.16 (s, 1H), 4.17 (d, 2H, J = 5.7 Hz), 3.65-3.45 (m, 8H).
10 Anal. Calcd for C₁₅H₁₆ClN₃O₃ + 0.25 H₂O: C, 55.22; H, 5.10; N, 12.88.
Found: C, 55.22; H, 5.08; N, 12.82.

Example 25

5-Chloro-1H-indole-2-carboxylic acid [(methoxy-methyl-carbamoyl)-methyl]-amide

Methoxymethylamine hydrochloride (1.0 mmol) and [(5-chloro-1H-indole-2-
15 carbonyl)-amino]-acetic acid (1.0 mmol) were coupled according to Procedure A. The resulting solid was suspended in ether, filtered and dried: Yield 158 mg, 53 %; PBMS 296/298 (MH+, 100 %);
¹H NMR (DMSO-d₆) δ 11.82 (s, 1H), 8.77 (t, 1H, J = 6 Hz), 7.73 (d, 1H, J = 2.0 Hz), 7.43 (d, 1H, J = 8.7 Hz), 7.19 (dd, 1H, J = 2.0, 8.7 Hz), 7.16 (s, 1H), 4.22 (d, 2H, J =
20 5.7 Hz), 3.76 (s, 3H), 3.14 (s, 3H).
Anal. Calcd for C₁₃H₁₄ClN₃O₃: C, 52.80; H, 4.77; N, 14.21.
Found: C, 52.51; H, 4.82; N, 14.01.

Example 26

5-Chloro-1H-indole-2-carboxylic acid [1-benzyl-2-25 (4-dimethylamino-piperidin-1-yl)-2-oxo-ethyl]-amide

4-Dimethylaminopiperidine (1.0 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-
amino]-3-phenyl-propionic acid (1.0 mmol) were coupled according to Procedure A. The residue was purified by chromatography on silica gel eluted with 5-30% ethanol in dichloromethane containing 0.5 % ammonium hydroxide, followed by trituration with
30 ether: Yield 21 mg, 5 %; PBMS 453/455 (MH+, 100 %);
¹H NMR (DMSO-d₆, partial) δ 11.75 (br, 1H), 8.94 (m, 1H), 7.72 (d, 1H, J = 2 Hz), 7.45-7.10 (m, 8H), 5.17 (m, 1H), 4.63 (m), 4.38 (m), 4.03 (m), 3.50 (m), 3.15 - 2.8 (m), 2.51 (s, 3H), 2.50 (s, 3H).

-67-

Example 27

5-Chloro-1H-indole-2-carboxylic acid (1-benzyl-2-oxo-2-piperazin-1-yl-ethyl)-amide

- Trifluoroacetic acid (4 ml) was added to 4-{2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl}-piperazine-1-carboxylic acid tert-butyl ester (0.6 mmol) at 0 °C and the resulting solution was stirred for 0.3 hours and concentrated. The residue was partitioned between ethyl acetate and 2 N NaOH, the organic layer separated and washed with brine, dried over Na₂SO₄, concentrated and the resulting solid triturated with ether: Yield 189 mg, 77 %; HPLC (60/40) 2.63 minutes (99 %); mp 166.5 - 168 °C; TSPMS 411/413 (MH⁺, 100 %);
- Anal. Calcd for C₂₂H₂₃ClN₄O₂ + 0.5 H₂O: C, 62.93; H, 5.76; N, 13.34.
Found: C, 62.64; H, 5.52; N, 13.34.

Example 28

5-Chloro-1H-indole-2-carboxylic acid [2-((3RS)-amino-pyrrolidin-1-yl)-1-benzyl-2-oxo-ethyl]-amide

- 4N HCl in 1,4-dioxane (5 ml) was added to (1-{2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl}-pyrrolidin-(3RS-yl)-carbamic acid tert-butyl ester (0.5 mmol). The resulting solution was stirred at 25 °C for 0.5 hours, concentrated and the residue triturated with ether: Yield 190 mg, 85 %; HPLC (60/40) 2.62 minutes (98 %); PBMS 411/413 (MH⁺, 100 %);
- Anal. Calcd for C₂₂H₂₃ClN₄O₂ + HCl + 1.7 H₂O: C, 55.28; H, 5.78; N, 11.72.
Found: C, 55.14; H, 5.86; N, 11.45.

Example 29

1-{(2RS)-[5-Chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl}-pyrrolidine-(2S)-carboxylic acid

- Trifluoroacetic acid was added to 1-{2(RS)-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl}-pyrrolidine-(2S)-carboxylic acid tert-butyl ester (1.0 mmol) at 25 °C. After 1.5 hours, the reaction was concentrated and the residue triturated first with ether then with a mixture of ether and hexanes. Yield 360 mg, 82 %; HPLC (60/40) 4.84 minutes (99 %); PBMS 440/442 (MH⁺, 40 %), 396/398 (MH⁺, 100 %);
- Anal. Calcd for C₂₃H₂₂ClN₃O₄ + 0.8 H₂O: C, 60.81; H, 5.24; N, 9.25.
Found: C, 60.74; H, 5.42; N, 8.96.

-68-

Example 29a

1-{2(R,S)-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl}-
pyrrolidine-(2S)-carboxylic acid tert-butyl ester

L-Proline-t-butyl ester (2.0 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-
5 3-phenyl-propionic acid (2.0 mmol) were coupled according to Procedure A and the
crude product purified by chromatography on silica gel eluting with 1:2 ethyl acetate:
hexanes: Yield 611 mg, 62 %; HPLC (60/40) 13.45 minutes (57 %) and 14.46 minutes
(41 %).

Example 30

10 5-Chloro-1H-indole-2-carboxylic acid ((1S)-methylcarbamoyl-2-thiazol-
4-yl-ethyl)-amide

(S)-2-Amino-N-methyl-3-thiazol-4-yl-propionamide hydrochloride (0.6 mmol) and
5-chloro-1H-indole-2-carboxylic acid (0.51 mmol) were coupled according to Procedure
A (0 - 25 °C reaction temperature, reaction solvent dimethyl-formamide). The crude
15 product was stirred in ether for 0.5 hours then filtered giving a beige solid: 182 mg, 98
%; HPLC (60/40) 3.41 minutes (98 %); mp >260 °C (dec); TSPMS 363/365 (MH+,
100 %);

¹H NMR (DMSO-d₆) δ 11.82 (br, 1H), 9.0 (d, 1H), 8.82 (br, 1H), 8.10 (br, 1H), 7.70 (m,
1H), 7.44 - 7.38 (m, 2H), 7.21 - 7.15 (m, 2H), 4.80 (m, 1H), 3.24 (m, 1H), 3.05 (m, 1H),
20 2.60 (d, 3H).

Example 30a

(S)-2-Amino-N-methyl-3-thiazol-4-yl-propionamide hydrochloride

(S)-(1-Methylcarbamoyl-2-thiazol-4-yl-ethyl)-carbamic acid tert-butyl ester (248
mg, 0.87 mmol) was dissolved in 4 M HCl-dioxane at 0°C. The resulting mixture was
25 stirred for 1 hour at 25 °C, concentrated and the residue triturated with ether. Yield,
202 g, 102 %; HPLC (70/30) 2.41 minutes (96 %);

Example 30b

(S)-2-(N-t-Butoxycarbonylamino)-N-methyl-3-thiazol-4-yl-propionamide

Methylamine hydrochloride (1.2 mmol) and Boc-L-3-(4-thiazolyl)alanine (1.0
30 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, acid
wash omitted) and the product used without further purification. Yield, 250 mg, 88 %.

Example 31**(±)-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-hydroxy-propionic acid methyl ester**

D,L-Serine methyl ester hydrochloride (2.1 mmol) and 5-chloro-1H-indole-2-carboxylic acid (2.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, washed with acid first then with saturated NaHCO₃) and the product purified by chromatography on silica gel eluted with 10, 20, 40 and 60% ethyl acetate in hexanes: Yield 565 mg, 95 %; HPLC (60/40) 3.46 minutes (98 %); mp 153 - 155 °C; TSPMS 297/299 (MH⁺, 100 / 40 %);
Anal. Calcd for C₁₃H₁₃ClN₂O₄: C, 52.62; H, 4.42; N, 9.44.
10 Found: C, 52.62; H, 4.54; N, 9.53.

Example 32**5-Chloro-1H-indole-2-carboxylic acid ((1S)-dimethylcarbamoyl-2-thiazol-4-yl-ethyl)-amide**

(S)-2-Amino-N,N-dimethyl-3-thiazol-4-yl-propionamidehydrochloride(0.43mmol)
15 and 5-chloro-1H-indole-2-carboxylic acid (0.40 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) and the crude product triturated first with 1:1 ether-hexanes, then with hexanes. Yield 115 mg, 75 %; HPLC (60/40) 3.72 minutes (99 %); mp 198 - 202 °C (shrinks on insertion at 192 °C); PBMS 377/379 (MH⁺, 100 %);
20 ¹H NMR (DMSO-d₆) δ 11.75 (s, 1H), 9.02 (d, 1H, J = 2 Hz), 8.9 (d, 1H, J = 8.2 Hz), 7.7 (d, 1H, J = 1.8 Hz), 7.41 (d, 1H, J = 6.7 Hz), 7.39 (s, 1H), 7.22 (s, 1H), 7.17 (dd, 1H, J = 2.2, 8.7 Hz), 5.30 (m, 1H), 3.24 (dd, A of AB, 1H, J = 7, 13 Hz), 3.16 (dd, B of AB, 1H, J = 8.5, 16 Hz), 3.07 (s, 3H), 2.84 (s, 3H).
Anal. Calcd for C₁₇H₁₇ClN₄O₂S + 0.125 H₂O: C, 53.86; H, 4.59; N, 14.78.
25 Found: C, 53.92; H, 4.47; N, 14.42.

Example 32a**(S)-2-Amino-N,N-dimethyl-3-thiazol-4-yl-propionamide hydrochloride**

(S)-(1-Dimethylcarbamoyl-2-thiazol-4-yl-ethyl)-carbamic acid tert-butyl ester was dissolved in 4M HCl-dioxanes at 0 °C and stirred at 25 °C for 2 hours. The mixture
30 was concentrated and the residue triturated with ether. Yield, 3.06 g, 105 %; HPLC (70/30) 2.12 minutes (97 %); PBMS 200 (MH⁺, 100 %).

-70-

Example 32b

(S)-(1-Dimethylcarbamoyl-2-thiazol-4-yl-ethyl)-carbamic acid tert-butyl ester

Dimethylamine hydrochloride (1.2 mmol) and Boc-L-3-(4-thiazolyl)alanine (1.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) and
5 the product purified by chromatography on silica gel eluted with 1 - 16 % ethanol in dichloromethane containing 0.5 % ammonium hydroxide. Yield 124 mg, 41 %.

Example 33

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3R,4S)-
dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide

10 (3R,4S)-Dihydroxypyrrolidine hydrochloride (0.5 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.5 mmol) were coupled according to Procedure A and the product purified by chromatography on silica gel eluted with 2 - 10 % ethanol in dichloromethane. Yield 180 mg, 86 %; HPLC (60/40) 3.14 minutes (98 %); TSPMS 428/430 (MH⁺, 100 %);

15 ¹H NMR (DMSO-d₆) δ 11.75 (br, 1H), 8.94 (d, 1H, J = 8 Hz), 7.72 (s, 1H), 7.4-7.1 (m, 8H), 5.03 (d, 0.5H, J = 5 Hz), 4.95 (d, 0.5H, J = 5 Hz), 4.90 (d, 1H, J = 5 Hz), 4.87 (m, 1H), 4.08 (m, 0.5H), 4.00 (m, 0.5H), 3.88 (m, 1.5H), 3.5 - 3.3 (m, 2.5H), 3.2 (m, 0.5H), 3.0 (m, 2H).

Anal. Calcd for C₂₂H₂₂ClN₃O₄ + 0.25 H₂O: C, 61.11; H, 5.25; N, 9.72.

20 Found: C, 60.91; H, 5.46; N, 9.43.

Example 33a

(Cis-3,4-)-Dihydroxypyrrolidine hydrochloride

Cis-3,4-Dihydroxy-pyrrolidine-1-carboxylic acid tert-butyl ester (1.99g, 9.8 mmol) was dissolved in 4M HCl-dioxane at 5 °C and the resulting suspension stirred at 25 °C
25 for 1 hour. The mixture was concentrated and the residue triturated with ether giving a light purple powder (1.30 g, 95 %).

Example 33b

Cis-3,4-Dihydroxy-pyrrolidine-1-carboxylic acid tert-butyl ester

A solution of crude 2,5-dihydro-pyrrole-1-carboxylic acid tert-butyl ester (10.5g,
30 62.1 mmol) in tetrahydro-furan (300 ml) was treated sequentially with osmium tetroxide (2.5 % in t-butanol, 6 mL) and N-methylmorpholine-N-oxide at 25 °C. After 48 hours, aqueous 10 % sodium thiosulfate solution was added and the mixture was stirred 30 minutes, partially concentrated to remove tetrahydro-furan, and the resulting aqueous

-71-

mixture extracted twice with ether. The ether extracts were washed with 10 % sodium thiosulfate, 0.1 N HCl, dried and concentrated giving a dark orange oil which was chromatographed on silica eluted with 1 %, 2 %, 4 %, 8 % and 10 % ethanol - dichloromethane giving an amber syrup (4.09 g).

5

Example 33c**2,5-Dihydro-pyrrole-1-carboxylic acid tert-butyl ester**

Di-t-butylidicarbonate (83 g, 380 mmol) was added to a solution of 3-pyrroline (containing 35 % pyrrolidine, 25 g, 362 mmol) in tetrahydrofuran (500 mL) at 0 °C. The mixture was stirred at 25 °C for 1 hour and concentrated giving 76.2 g of a yellow
10 oil which was used without purification.

Example 34**(3S)-[(5-Chloro-1H-indole-2-carbonyl)-amino]-4-****(4-hydroxy-piperidin-1-yl)-4-oxo-butyric acid tert-butyl ester**

(S)-3-Amino-4-(4-hydroxy-piperidin-1-yl)-4-oxo-butyric acid tert-butyl ester (0.8
15 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.8 mmol) were coupled according to Procedure A and the product purified by chromatography on silica gel eluted with 25, 40, 50, 75 and 100 % ethyl acetate in hexanes. Yield 330 mg, 94 %; HPLC (60/40) 4.18 minutes (97 %); TSPMS 450/452 (MH+, 100 %).

Example 34a

20

(S)-3-Amino-4-(4-hydroxy-piperidin-1-yl)-4-oxo-butyric acid tert-butyl ester

Diethylamine (1.0 mmol) was added to (S)-3-(9H-fluoren-9-ylmethoxycarbonyl-amino)-4-(4-hydroxy-piperidin-1-yl)-4-oxo-butyric acid tert-butyl ester in dimethyl-formamide (5 ml) at 25 °C. After 1 hour, the reaction mixture was concentrated, the residue suspended in 1:1 ether/dichloromethane, filtered and concentrated. The
25 residue was purified by chromatography on silica gel eluted with 1-50% ethanol in dichloromethane containing 0.5 % ammonium hydroxide. Yield 217 mg, 80 %.

Example 34b**(S)-3-(9H-Fluoren-9-ylmethoxycarbonylamino)-4-(4-hydroxy-****piperidin-1-yl)-4-oxo-butyric acid**

30

4-Hydroxypiperidine (2.1 mmol) and N-FMOC-L-aspartic acid- β -t-butyl ester (2.0 mmol) were coupled according to Procedure A (96 hour reaction time, washed with acid only) and the product purified by chromatography on silica gel eluted with 1 - 4

-72-

% ethanol in dichloromethane. Yield 516 mg, 52 %; HPLC (60/40) 5.33 minutes (93 %).

Example 35

5-Chloro-1H-indole-2-carboxylic acid [(1R)-benzyl-2-

(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

(R)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one hydrochloride (3.1 mmol) and 5-chloro-1H-indole-2-carboxylic acid (3.4 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 60 hour reaction time) and the product purified by chromatography on silica gel eluted with 50, 75 and 100 % ethyl acetate in hexanes followed by trituration with 1:1 ether-hexanes. Yield 1.1 g, 84 %; HPLC (60/40) 4.06 minutes (99 %); PBMS 426/428 (MH⁺, 100 %);
Anal. Calcd for C₂₃H₂₄ClN₃O₃ + 0.25 H₂O: C, 64.18; H, 5.74; N, 9.76.
Found: C, 64.28; H, 5.94; N, 9.41.

Example 35a

(R)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one hydrochloride

(R)-2-(N-t-butoxycarbonylamino)-1-(4-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one (12.5 mmol) was dissolved in 4M HCl-dioxane at 0 °C and the resulting suspension stirred at 25 °C for 1 hour. The mixture was concentrated and the residue triturated with ether. Yield, 3.44 g, 97 %.

Example 35b

(R)-2-(N-t-butoxycarbonylamino)-1-(4-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one

(R)-2-(N-t-butoxycarbonylamino)-3-phenyl-propan-1-one (14 mmol) and 4-hydroxypiperidine (21.5 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, washed with acid first then base) and the product used without further purification. Yield 4.7 g, 94 %; HPLC (60/40) 3.52 minutes (98 %).

Example 36

1H-Indole-2-carboxylic acid [2-(1,1-dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-amide

2-Amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-ethanone hydrochloride (1.0 mmol) and 1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 60 hour reaction time) with the following workup: the reaction mixture was diluted with ethyl acetate and 2 N NaOH, the resulting precipitate was collected and washed with 2N NaOH, 1N HCl and water. Yield 135 mg, 42 %; HPLC (60/40) 2.97 minutes (97 %); PBMS 322 (MH⁺, 100 %);

-73-

Anal. Calcd for $C_{14}H_{15}N_3O_4S + 0.25 H_2O$: C, 51.60; H, 4.79; N, 12.90.

Found: C, 51.31; H, 4.66; N, 12.88.

Example 36a

2-Amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-ethanone hydrochloride

- 5 [2-(1,1-Dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (11 mmol) was dissolved in 4M HCl-dioxane at 0 °C and the resulting suspension stirred at 25 °C for 1 hour. The mixture was concentrated and the residue triturated with ether. Yield, 2.3 g, 100 %.

Example 36b

- 10 [2-(1,1-Dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester

m-Chloroperoxybenzoic acid (35 mmol) was added slowly to (2-oxo-2-thiazolidin-3-yl-ethyl)-carbamic acid tert-butyl ester (14 mmol) in dichloromethane (35 ml) at 0°C. After foaming subsided, the mixture was stirred an additional 2.5 hours at 25 °C. The mixture was diluted with ethyl acetate, the resulting solution washed three times with
15 a 1:1 mixture of saturated aqueous $NaHCO_3$ and 10 % aqueous NaS_2O_3 solution, once with saturated $NaHCO_3$, dried, concentrated and the residue triturated with 1:1 ether/hexanes. Yield, 3.6 g, 92 %.

Example 36c

(2-Oxo-2-thiazolidin-3-yl-ethyl)-carbamic acid tert-butyl ester

- 20 Thiazolidine (85 mmol) and Boc-glycine (57 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 60 hour reaction time) and the product used without purification. Yield 12.7g, 90 %.

Example 37

5-Fluoro-1H-indole-2-carboxylic acid [(1S)-benzyl-

- 25 2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

(S)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one hydrochloride (0.65 mmol) and 5-fluoro-1H-indole-2-carboxylic acid (0.73 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) and the product purified by chromatography on silica gel eluted with 20, 30, 40, 50, 75 and 100 % ethyl acetate in
30 hexanes. Yield 228 mg, 84 %; HPLC (60/40) 3.57 minutes (98 %); PBMS 410 (MH+, 100 %);

Anal. Calcd for $C_{23}H_{24}FN_3O_3 + 0.25 H_2O$: C, 66.73; H, 5.97; N, 10.15.

Found: C, 66.68; H, 6.19; N, 9.94.

-74-

Example 38

1H-Indole-2-carboxylic acid [(1S)-benzyl-2-(4-hydroxy-piperidin-1-yl)-
2-oxo-ethyl]-amide

(S)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one hydrochloride (3.4 mmol) and 1H-indole-2-carboxylic acid (3.7 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 48 hour reaction time). The product was purified by chromatography on silica gel eluted with 50, 75 and 100 % ethyl acetate in hexanes, followed by trituration with 1:1 ether - hexanes. Yield 1.14 g, 86 %; HPLC (60/40) 3.52 minutes (98 %); PBMS 392 (MH+, 100 %);

10 Anal. Calcd for $C_{23}H_{25}N_3O_3 + 0.25 H_2O$: C, 69.77; H, 6.49; N, 10.61.
Found: C, 69.99; H, 6.72; N, 10.47.

Example 39

5-Fluoro-1H-indole-2-carboxylic acid [(1S)-(4-fluoro-benzyl)-2-morpholin-
4-yl-2-oxo-ethyl]-amide

15 (S)-2-Amino-3-(4-fluoro-phenyl)-1-morpholin-4-yl-propan-1-one hydrochloride (0.48 mmol) and 5-fluoro-1H-indole-2-carboxylic acid (0.48 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 48 hour reaction time, washed with acid first then base) and the product purified by chromatography on silica gel eluted with 20, 30, 40, 50 and 75 % ethyl acetate in hexanes. Yield 189 mg, 95%;

20 HPLC (60/40) 4.76 minutes (97 %); PBMS 414 (MH+, 100 %);
 1H NMR ($CDCl_3$) δ 9.23 (br, 1H), 7.4 - 7.1 (m, 5H), 7.1 - 6.94 (m, 3H), 6.9 (d, 1H, J = 2 Hz), 5.30 (m, 1H), 3.72 - 3.48 (m, 5H), 3.42 (m, 1H), 3.03 (m, 4H).
Anal. Calcd for $C_{22}H_{21}F_2N_3O_3$: C, 63.92; H, 5.12; N, 10.16.
Found: C, 64.30; H, 5.34; N, 9.82.

25

Example 39a

(S)-2-Amino-3-(4-fluoro-phenyl)-1-morpholin-4-yl-propan-1-one hydrochloride

(S)-2-(N-t-Butoxycarbonylamino)-3-(4-fluoro-phenyl)-1-morpholin-4-yl-propan-1-one (3.1 mmol) was dissolved in 4M HCl-dioxane at 0 °C and the resulting suspension stirred at 25 °C for 1 hour. The mixture was concentrated and the residue triturated

30 with ether. Yield, 776 mg, 88 %; HPLC (60/40) 2.31 minutes (99 %).

Example 39b

(S)-2-(N-t-Butoxycarbonylamino)-3-(4-fluoro-phenyl)-1-morpholin-4-yl-propan-1-one

Morpholine (3.7 mmol) and (S)-Boc-4-fluoro-phenyl-alanine (3.5 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 60 hour reaction
5 time, washed first with acid, then base) and the product purified by chromatography on silica gel eluted with 20, 30 and 40 % ethyl acetate in hexanes. Yield 1.08 g oil, 87 %.

Example 40

5-Fluoro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(1,1-dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-amide

10

(S)-2-Amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-3-phenyl-propan-1-one hydrochloride (1.0 mmol) and 5-fluoro-1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, reaction time 48
15 hour, washed with acid first, then base) and the product purified by chromatography on silica gel eluted with 20, 30, 40 and 50 % ethyl acetate in hexanes. Yield 404 mg, 94 %; HPLC (60/40) 4.74 min (98 %); PBMS 430 (MH⁺, 100 %);
¹H NMR (CDCl₃) δ 9.53 (br, 0.5H), 9.44 (br, 0.5H), 7.44 (d, 0.5H, J = 9 Hz), 7.4-7.1 (m, 7H), 7.02 (M, 1H), 6.84 (s, 0.5H), 6.81 (s, 0.5H), 5.20 (m, 0.5H), 4.96 (m, 0.5H), 4.68 (d, 0.5H, J = 11 Hz), 4.52 (d, A of AB, 0.5H, J = 11.5 Hz), 4.37 (d, B of AB, 0.5H, J = 11.5
20 Hz), 4.20 (m, 0.5H), 4.03 (m, 0.5H), 3.80 (m, 0.5H), 3.50 (d, 0.5H, J = 11 Hz), 3.3-3.0 (m, 4H), 2.69 (m, 0.5H).

Example 40a

(S)-2-Amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-3-phenyl-propan-1-one hydrochloride

(S)-2-(N-t-Butoxycarbonylamino)-1-(1,1-dioxo-1-thiazolidin-3-yl)-3-phenyl-propan-
25 1-one was dissolved in 4M HCl-dioxanes at 0 °C. The solution was stirred at 25 °C for 1 hour, concentrated and the residue triturated with ether. Yield, 866 mg, 84 %.

Example 40b

(S)-2-(N-t-Butoxycarbonylamino)-1-(1,1-dioxo-1-thiazolidin-3-yl)-3-phenyl-propan-1-one

30 A solution of m-chloroperoxybenzoic acid (9 mmol) and (S)-(1-benzyl-2-oxo-2-thiazolidin-3-yl-ethyl)-carbamic acid-tert-butyl ester (3 mmol) in dichloromethane (9 ml) were heated at reflux for 6 hours. The mixture was diluted with ethyl acetate, the resulting solution washed three times with a 1:1 mixture of 10 % aqueous Na₂S₂O₃ and

-76-

saturated aqueous NaHCO_3 , dried and concentrated. The resulting foam was purified by chromatography on silica gel eluted with 20, 30 and 40 % ethyl acetate in hexanes giving a colorless foam (979 mg, 89 % yield).

Example 40c

5 (S)-(1-Benzyl-2-oxo-2-thiazolidin-3-yl-ethyl)-carbamic acid tert-butyl ester

Thiazolidine (38 mmol) and Boc-L-phenylalanine (19 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, washed with acid then base) and the product used without further purification. Yield 5.5 g, 86 %.

Example 41

10 5-Fluoro-1H-indole-2-carboxylic acid [2-(1,1-dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-amide

2-Amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-ethanone hydrochloride (1.0 mmol) and 5-fluoro-1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 48 hour reaction time) with the following workup.

15 The reaction mixture was diluted with ethyl acetate and 1N HCl, the resulting suspension was filtered and the collected solid washed with 2 N HCl, 2 N NaOH and water. The filtered solid was boiled in acetone, filtered and dried. Yield 134 mg, 40 %; HPLC (60/40) 3.06 minutes (97 %); mp 239 - 241 °C (with discoloration); PBMS 340 (MH⁺, 70 %), 357 (100 %)

20 ¹H NMR (DMSO-d₆) δ 11.74 (s, 1H), 8.82 (m, 1H), 7.43 (m, 2H), 7.17 (s, 1H), 7.05 (dt, 1H, J = 3, 9 Hz), 4.86 (s, 1.2H), 4.52 (s, 0.8H), 4.27 (d, 0.8H, J = 5.5 Hz), 4.13 (d, 1.2H, J = 6 Hz, superimposed on m, 1.2H), 3.86 (t, 1.2H, J = 7.4 Hz), 3.58 (t, 0.8H, J = 7 Hz), 3.46 (t, 1.2H, J = 7.2 Hz).

Anal. Calcd for C₁₄H₁₄FN₃O₄S + 0.6 H₂O: C, 48.02; H, 4.38; N, 12.00.

25 Found: C, 47.99; H, 4.04; N, 12.00.

Example 42

5-Cyano-1H-indole-2-carboxylic acid ((1S)-benzyl-2-oxo-2-thiazolidin-3-yl-ethyl)-amide

(S)-2-Amino-3-phenyl-1-thiazolidin-3-yl-propan-1-one hydrochloride (4.0 mmol) and 5-cyano-1H-indole-2-carboxylic acid (4.0 mmol) were coupled according to

30 Procedure A (0 - 25 °C reaction temperature, 48 hours reaction time) with the following workup: the reaction mixture was diluted with ethyl acetate and 2 N HCl, the resulting precipitate collected by filtration, washed with 2 N HCl and 2 N NaOH. The crude product was purified by chromatography on silica gel eluted with 30, 40 and 50 % ethyl

-77-

acetate in hexanes. Yield 1.22 g, 75 %; HPLC (60/40) 4.74 minutes (97 %); PBMS 405 (MH⁺, 100 %);

Anal. Calcd for C₂₂H₂₀N₄O₂S + 0.5 H₂O: C, 63.90; H, 5.12; N, 13.55.

Found: C, 64.18; H, 5.04; N, 13.47.

5

Example 42a

(S)-2-Amino-3-phenyl-1-thiazolidin-3-yl-propan-1-one hydrochloride

(S)-(1-Benzyl-2-oxo-2-thiazolidin-3-yl-ethyl)-carbamic acid tert-butyl ester (16 mmol) was dissolved in 4 M HCl-dioxanes at 0 °C, the solution stirred at 25 °C for 1 hour, the reaction concentrated and the residue triturated with ether. Yield, 4.2 g, 95 %.

10

Example 42b

5-Cyano-1H-indole-2-carboxylic acid

5-Cyano-1H-indole-2-carboxylic acid ethyl ester (1.71 g, 8.0 mmol) was added to a solution of ethanol (10 mL) and potassium hydroxide (2 g) and the resulting mixture heated at reflux for 1 hour. Water was added to dissolve the precipitate, and 6 N HCl was added to bring the pH to 1. A precipitate formed. The mixture was cooled in an ice bath, filtered, and the resulting colorless solid washed with cold water and dried (1.51 g). A portion (1.4 g) was suspended in hot acetic acid (40 mL) and cooled giving a solid which was filtered, washed with cold ethyl acetate and dried: Yield 980 mg (70 %); HPLC (60/40) 3.09 minutes (97 %).

20

Example 43

1H-Indole-2-carboxylic acid [(1S)-benzyl-2-(1,1-dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-amide

(S)-2-Amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-3-phenyl-propan-1-one hydrochloride (0.56 mmol) and 1H-indole-2-carboxylic acid (0.56 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) and the product triturated with 1:1 ether-hexanes. Yield 213 mg, 92 %; HPLC (60/40) 4.15 minutes (99 %); PBMS 412 (MH⁺, 100 %);

25

Anal. Calcd for C₂₁H₂₁N₃O₄S + 0.5 H₂O: C, 59.99; H, 5.27; N, 9.99.

30 Found: C, 60.25; H, 5.27; N, 9.98.

-78-

Example 44

5-Chloro-1H-indole-2-carboxylic acid [2-(1,1-dioxo-1-thiazolidin-3-yl)-
2-oxo-ethyl]-amide

- 2-Amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-ethanone hydrochloride (0.6 mmol) and
5 5-chloro-1H-indole-2-carboxylic acid (0.6 mmol) were coupled according to Procedure
A (0 - 25 °C reaction temperature, 120 hour reaction time) with the following workup:
the reaction mixture was diluted with ethyl acetate and 2N HCl, the resulting precipitate
was collected by filtration followed by washing with 2 N HCl, 2 N NaOH, water and
ether. Yield 110 mg, 52 %; HPLC (60/40) 3.37 minutes (99 %); mp 236 - 239 °C
10 (dec); PBMS 356/358 (MH+, 100 %);
¹H NMR (acetone-d₆) δ 11.0 (br, 1H), 8.0 (br, 1H), 7.66 (d, 1H, J = 2 Hz), 7.55 (d, 1H,
J = 8.7 Hz), 7.21 (dd, 1H, J = 2.0, 8.7 Hz), 7.15 (d, 1H, J = 2 Hz), 4.77 (s, 1.1H), 4.49
(s, 0.9H), 4.37 (d, 0.9H, J = 5.3 Hz), 4.27 (d, ca. 1H, J = 5.3 Hz, superimposed on m,
ca. 1H), 4.04 (t, 1.1H, J = 7 Hz), 3.54 (t, 0.9H, J = 7 Hz), 3.40 (t, 1.1H, J = 7 Hz).
15 Anal. Calcd for C₁₄H₁₄ClN₃O₄S + 1.6 H₂O: C, 43.72; H, 4.51; N, 10.93.
Found: C, 44.05; H, 3.88; N, 10.99:

Example 45

5-Chloro-1H-indole-2-carboxylic acid (2-oxo-2-thiazolidin-3-yl-ethyl)-amide

- 2-Amino-1-thiazolidin-3-yl-ethanone hydrochloride (3.1 mmol) and 5-chloro-1H-
20 indole-2-carboxylic acid (3.4 mmol) were coupled according to Procedure A (0-25°C,
120 hour reaction time) substituting the following workup: the reaction mixture was
stirred with ethyl acetate and 2N HCl, filtered, and the filtered solid washed with 2N
HCl, 2N NaOH and ether. Yield 988 mg, 98 %; HPLC (70/30) 3.25 minutes (99 %);
mp 253 - 255 °C (dec, darkening at 243 °C); PBMS 324/326 (MH+, 100 %);
25 ¹H NMR (acetone-d₆) δ 11.03 (br, 1H), 7.88 (br, 1H), 7.66 (d, 1H, J = 2 Hz), 7.54 (d,
1H, J = 8.3 Hz), 7.21 (dd, 1H, J = 2, 8.3 Hz), 4.67 (s, 0.8H), 4.53 (s, 1.2 H), 4.24 (m,
2H), 3.87 (t, 1.2H, J = 7 Hz), 3.78 (t, 0.8H, J = 7 Hz), 3.18 (t, 1.2H, J = 7 Hz), 3.05 (t,
0.8H, J = 7 Hz).
A sample was recrystallized from acetic acid for analysis (mp 262 - 264 °C):
30 Anal. Calcd for C₁₄H₁₄ClN₃O₂S: C, 51.93; H, 4.36; N, 12.98.
Found: C, 51.78; H, 4.38; N, 12.95.

-79-

Example 45a**2-Amino-1-thiazolidin-3-yl-ethanone hydrochloride**

(2-Oxo-2-thiazolidin-3-yl-ethyl)-carbamic acid tert-butyl ester (5.41 g, 22 mmol) was dissolved in 4M HCl-dioxane (80 mL) at 0 °C. The resulting solution was stirred at 25 °C for 2 hours, concentrated and the residue triturated with ether. Yield, 3.9 g, 97 %.

Example 46**5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-****(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide**

(S)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one hydrochloride (0.8 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.9 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 48 hour reaction time) and the product purified by chromatography on silica gel eluted with 50, 75 and 100 % ethyl acetate in hexanes followed by trituration from 1:1 ether-hexanes. Yield 266 mg, 76 %; HPLC (60/40) 4.09 minutes (99 %); PBMS 426/428 (MH⁺, 100 %);
Anal. Calcd for C₂₃H₂₄ClN₃O₃ + 0.33 H₂O: C, 63.96; H, 5.76; N, 9.73.
Found: C, 63.90; H, 5.74; N, 9.58.

Example 46a**(S)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one hydrochloride**

(S)-[1-Benzyl-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (3.66 g, 10.5 mmol) was dissolved in 4M HCl-dioxane (39 mL) at 0 °C. The mixture was stirred at 25 °C for 1 hour, concentrated and the residue triturated with ether. Yield 3.06 g, 102 %.

Example 46b**(S)-[1-Benzyl-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester**

4-Hydroxypiperidine (75 mmol) and Boc-L-phenylalanine (38 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 144 hour reaction time) and the product used without further purification. Yield 12.2 g, 96 %; HPLC (60/40) 3.45 minutes (97 %).

-80-

Example 47

5-Bromo-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(1,1-dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-amide

(S)-2-Amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-3-phenyl-propan-1-one hydrochloride (0.3 mmol) and 5-bromo-1H-indole-2-carboxylic acid (0.3 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, washed with acid first, then base) and the product purified by chromatography on silica gel eluted with 30, 40 and 50 % ethyl acetate in hexanes. The product was collected as an off-white foam and triturated with 1:1 ether - hexanes to give 107 mg, 73 %; HPLC (60/40) 6.21 minutes (99 %); PBMS 490/492 (MH⁺, 100 %);

¹H NMR (CDCl₃) δ 9.53 (br, 0.5H), 9.44 (br, 0.5H), 7.78 (d, 0.5H, J = 2 Hz), 7.76 (d, 0.5H, J = 2 Hz), 7.4 - 7.2 (m, 7H), 7.10 (d, 0.5H, J = 9 Hz), 7.02 (d, 0.5H, J = 9 Hz), 6.86 (s, 0.5H), 6.81 (s, 0.5H), 5.21 (m, 0.5H), 4.95 (m, 0.5H), 4.62 (d, 0.5H, J = 11 Hz), 4.47 (d, A of AB, 0.5H, J = 13 Hz), 4.38 (d, B of AB, 0.5H, J = 13 Hz), 4.20 (m, 0.5H), 4.03 (m, 0.5H), 3.82 (m, 0.5H), 3.44 (d, 0.5H, J = 11 Hz), 3.33 - 3.0 (m, 4H), 2.70 (m, 0.5H).

Anal. Calcd for C₂₁H₂₀BrN₃O₄S + 0.2 H₂O: C, 51.06; H, 4.16; N, 8.51.

Found: C, 51.44; H, 4.36; N, 7.93.

Example 48

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-oxo-2-(3-oxo-pyrrolidin-1-yl)-ethyl]-amide

(S)-1-(2-Amino-3-phenyl-propionyl)-pyrrolidin-3-one hydrochloride (0.6 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.6 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, washed with acid first, then base) and the product purified by chromatography on silica gel eluted with 40 and 50 % ethyl acetate in hexanes, followed by trituration of the resulting foam with ether. Yield 112 mg, 45%; HPLC (60/40) 5.13 minutes (>99 %); PBMS 410/412 (MH⁺, 100 %);

¹H NMR (CDCl₃) δ 9.19 (m, 1H), 7.60 (m, 1H), 7.3-7.15 (m, 8H), 6.86 (m, 1H), 4.23 (m, 0.5H), 4.95 (m, 0.5H), 4.0 - 3.7 (m, 3H), 3.27 (m, 1H), 3.15 (m, 1H), 3.05 (m, 0.5H), 2.85 (d, 0.5H, J = 28 Hz), 2.45 (m, 1.5H), 2.15 (m, 0.5H).

Anal. Calcd for C₂₂H₂₀ClN₃O₃ + 0.55 H₂O: C, 62.95; H, 5.07; N, 10.01.

Found: C, 63.31; H, 5.09; N, 9.61.

SUBSTITUTE SHEET (RULE 26)

Example 48a**(S)-1-(2-Amino-3-phenyl-propionyl)-pyrrolidin-3-one hydrochloride**

(S)-[1-Benzyl-2-oxo-2-(3-oxo-pyrrolidin-1-yl)-ethyl]-carbamic acid tert-butyl ester (552 mg, 1.7 mmol) was dissolved in 4M HCl-dioxane (6.2 mL) at 0°C. The mixture
5 was stirred at 25 °C for 1 hour, concentrated and the residue triturated with ether to give a light brown solid. Yield, 482 mg, 108 %.

Example 48b**(S)-[1-Benzyl-2-oxo-2-(3-oxo-pyrrolidin-1-yl)-ethyl]-carbamic acid tert-butyl ester**

A solution of dimethyl sulfoxide (4.07 g, 52 mmol) and oxalyl chloride (3.61 g,
10 28 mmol) were added slowly in that order to dichloromethane (50 ml) at -78 °C. (S)-[1-Benzyl-2-(3-hydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (24 mmol) in dichloromethane (30 ml) was added to the above solution via canula, the reaction temperature brought to -30 °C for 0.5 hours, then lowered to -78 °C followed by the addition of triethylamine (118 mmol). The reaction was then warmed to 25 °C,
15 diluted with ethyl acetate, washed three times with 1:1 saturated NaHCO₃ / brine, the organics dried over MgSO₄ and concentrated. The resulting foam was purified by chromatography on silica gel eluted with 30, 40 and 50 % ethyl acetate in hexanes to give a light yellow foam (7.5 g, 95 % yield).

Example 48c

20 **(S)-[1-Benzyl-2-(3RS)-hydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester**

(±)-3-Pyrrolidinol (75 mmol) and Boc-L-phenylalanine (38 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, washed with acid first, then base) and the product used without further purification. Yield 12.2 g, 96 %; HPLC
25 (60/40) 3.45 minutes (96 %).

Example 49**5-Chloro-1H-indole-2-carboxylic acid ((1S)-benzyl-2-oxo-2-thiazolidin-3-yl-ethyl)-amide**

(S)-2-Amino-3-phenyl-1-thiazolidin-3-yl-propan-1-one hydrochloride (2.6 mmol) and 5-chloro-1H-indole-2-carboxylic acid (2.6 mmol) were coupled according to
30 Procedure A (0 - 25 °C reaction temperature, 96 hour reaction time, washed with acid first then base). The crude product was triturated with 1:1 ether-hexanes and dried. Yield 966 mg, 91 %; HPLC (60/40) 7.99 minutes (97 %); PBMS 414/416 (MH⁺, 100 %);

-82-

¹H NMR (CDCl₃) δ 9.26 (br, 1H), 7.59 (m, 1H), 7.35 - 7.20 (m, 8H), 6.84 (m, 1H), 5.14 (m, 1H), 4.61 (d, A of AB, 0.6H, J = 10.3 Hz), 4.52 (d, 0.4H, J = 11.6 Hz), 4.42 (d, B of AB, 0.6H, J = 10.3 Hz), 3.88 (m, 0.4H), 3.80 - 3.65 (m, ca 1.5H), 3.2 (m, ca. 2.5H), 3.04 (m, 0.4H), 2.95-2.8 (m, 1.2H), 2.63 (m, 0.6H).

- 5 Anal. Calcd for C₂₁H₂₀ClN₃O₂S + 0.6 H₂O: C, 59.39; H, 5.03; N, 9.89.
Found: C, 59.39; H, 4.96; N, 9.52.

Example 50

5-Chloro-1H-indole-2-carboxylic acid ((1S)-benzyl-2-oxo-2-thiomorpholin-4-yl-ethyl)-amide

- 10 (S)-2-Amino-3-phenyl-1-thiomorpholin-4-yl-propan-1-one hydrochloride (2.6 mmol) and 5-chloro-1H-indole-2-carboxylic acid (2.6 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, washed with acid first, then base). The crude product was then triturated with 1:1 ether - hexanes and dried. Yield 1.03 g, 94 %; HPLC (60/40) 8.74 minutes (99 %); PBMS 428/430 (MH+, 100 %);
- 15 Anal. Calcd for C₂₂H₂₂ClN₃O₂S: C, 61.75; H, 5.18; N, 9.82. Found: C, 62.04; H, 5.58; N, 9.72.

Example 50a

(S)-2-Amino-3-phenyl-1-thiomorpholin-4-yl-propan-1-one hydrochloride

- (S)-(1-Benzyl-2-oxo-2-thiomorpholin-4-yl-ethyl)-carbamic acid tert-butyl ester (17.8 mmol) was dissolved in 4M HCl-dioxane (67 mL) at 0 °C, the solution stirred at 25 °C for 1 hour, the reaction concentrated and the residue triturated with ether. Yield, 5.0 g, 98 %; PBMS 251 (MH+, 100 %).

Example 50b

(S)-(1-Benzyl-2-oxo-2-thiomorpholin-4-yl-ethyl)-carbamic acid tert-butyl ester

- 25 Thiomorpholine (38 mmol) and Boc-L-phenylalanine (19 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) with the following workup: the reaction mixture was concentrated, diluted with ethyl acetate, then washed first with 1 N HCl three times, then with 2 N NaOH, the organic layer dried over MgSO₄ and concentrated. The resulting foam was used without further purification. Yield 6.3 g, 30 95 %.

Example 51

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(1,1-dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-amide

(S)-2-Amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-3-phenyl-propan-1-one hydrochloride (0.8 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.8 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, washed with acid first, then base). The product was purified by chromatography on silica gel eluted with 30, 40 and 50 % ethyl acetate in hexanes followed by trituration with 1:1 ether - hexanes. Yield 266 mg, 75 %; HPLC (60/40) 5.52 minutes (>99 %); PBMS 446/448 (MH+, 100 %);
1H NMR (CDCl₃) δ 9.21 (br, 0.5H), 9.15 (br, 0.5H), 7.62 (br, 0.5H, J = 2 Hz), 7.60 (d, 0.5H, J = 2 Hz), 7.35 - 7.20 (m, 7H), 7.10 (d, 0.5H, J = 8.5 Hz), 7.02 (d, 0.5H, J = 8.5 Hz), 6.84 (d, 0.5H, J = 2 Hz), 6.81 (d, 0.5H, J = 2 Hz), 5.21 (m, 0.5H), 4.93 (m, 0.5H), 4.62 (d, 0.5H, J = 11 Hz), 4.47 (d, A of AB, 0.5H, J = 13 Hz), 4.39 (d, B of AB, 0.5H, J = 13 Hz), 4.22 (m, 0.5H), 4.03 (m, 0.5H), 3.83 (m, 0.5H), 3.44 (d, 0.5H, J = 11 Hz), 3.3-3.0 (m, 4H), 2.67 (m, 0.5H).

Example 52

5-Chloro-1H-indole-2-carboxylic acid [(1S)-(4-chloro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

(S)-2-Amino-3-(4-chloro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride (0.98 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.92 mmol) were coupled according to Procedure A and the product purified by chromatography on silica gel eluted with 50, 75 and 100 % ethyl acetate in hexanes. Yield 362 mg, 86 %; HPLC (60/40) 5.06 minutes (97 %); mp 227 - 229 °C; TSPMS 460/462 (MH+, 100%);
Anal. Calcd for C₂₃H₂₃Cl₂N₃O₃: C, 60.01; H, 5.04; N, 9.13. Found: C, 59.83; H, 5.18; N, 9.16.

Example 52a

(S)-2-Amino-3-(4-chloro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride

(S)-[1-(4-Chloro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (475 mg, 1.2 mmol) was dissolved in 4M HCl-dioxane (5 mL) at 0 °C. The mixture was stirred for 1.5 hour at 25 °C, concentrated and the residue triturated with ether. Yield, 422 mg, 105 %; TSPMS 283 (MH+, 100 %).

Example 52b

(S)-[1-(4-Chloro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-
carbamic acid tert-butyl ester

4-Hydroxypiperidine (2.6 mmol) and Boc-L-p-chlorophenylalanine (2.5 mmol) were coupled according to Procedure A and the product purified by chromatography on silica gel eluted with 1:1 and 3:1 ethyl acetate/hexanes. Yield 662 mg, 69 %.

Example 53

5-Chloro-1H-indole-2-carboxylic acid [2-(4-hydroxy-piperidin-1-yl)-(1S)-
(1H-imidazol-4-ylmethyl)-2-oxo-ethyl]-amide

(S)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-(1H-imidazol-4-yl)-propan-1-one hydrochloride (0.7 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.7 mmol) were coupled according to Procedure A (120 hour reaction time, acid wash omitted). The crude product was triturated twice with ether, with 1:1 ether-hexanes and the residue purified by chromatography on silica gel eluted with 5 - 20 % ethanol in dichloromethane containing 0.5% ammonium hydroxide. Yield 232 mg, 81 %; HPLC (40/60) 2.57 minutes (98 %); PBMS 416/418 (MH+, 100 %);
Anal. Calcd for $C_{20}H_{22}ClN_5O_3 + 0.55 H_2O$: C, 56.42; H, 5.47; N, 16.45.
Found: C, 6.07; H, 5.65; N, 16.08.

Example 53a

(S)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-(1H-imidazol-4-yl)-
propan-1-one hydrochloride

(S)-{2-(4-Hydroxy-piperidin-1-yl)-2-oxo-1-[1-(toluene-4-sulfonyl)-1H-imidazol-4-ylmethyl]-ethyl}-carbamic acid tert-butyl ester (512 mg, 1.0 mmol) was dissolved in 4 M HCl - dioxane (3 mL) at 0 °C. The mixture was stirred at 25 °C for 1.5 hours, concentrated and the residue triturated with ether. Yield, 422 mg, 105 %; TSPMS 283 (MH+, 100 %).

Example 53b

(S)-{2-(4-Hydroxy-piperidin-1-yl)-2-oxo-1-[1-(toluene-4-sulfonyl)-
1H-imidazol-4-ylmethyl]-ethyl}-carbamic acid tert-butyl ester

4-Hydroxypiperidine (303 mg, 3.0 mmol), triethylamine (394mg, 3.9 mmol) and diethyl cyanophosphonate (636 mg, 3.9 mmol) were added in that order to Boc-N_m-tosyl-L-histidine (J Med Chem 30 536 (1987); 1.32 g, 3.9 mmol) in dichloromethane (10 ml) at 25 °C. After 120 hours, the solution was diluted with ethyl acetate, washed

5 5-Chloro-1H-indole-2-carboxylic acid (2S)-[(5-chloro-1H-indole-2-carbonyl)-
amino]-3-(4-hydroxy-piperidin-1-yl)-3-oxo-propyl ester

Found: C, 56.79; H, 4.90; N, 9.65.

(S)-2-Amino-3-hydroxy-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride

Example 54b

(S)-1-Hydroxymethyl-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl-
carbamic acid tert-butyl ester

4-Hydroxypiperidine (6.7 mmol) and Boc-L-serine (6.4 mmol) were coupled according to Procedure A (60 hour reaction time) with the following workup: the reaction mixture was concentrated, the residue dissolved in chloroform and 1N NaOH (6 mL), and the resulting solution extracted repeatedly (ten or more times) with chloroform. The chloroform extracts were concentrated and the residue purified by chromatography on silica gel eluted with 1 - 16 % ethanol in dichloromethane. Yield 751 mg, 41 %; HPLC (40/60) 2.72 minutes (96 %).

Example 55

5-Chloro-1H-indole-2-carboxylic acid [(1S)-(4-hydroxy-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

- (S)-2-Amino-3-(4-hydroxy-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride (0.68 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.65 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) with the following workup: the reaction mixture was diluted with ethyl acetate, the resulting solution washed with 1N NaOH (2 ml), the aqueous layer extracted three times with ethyl acetate, the combined organic extracts washed with 1N HCl, dried and concentrated.
- 10 The residue was purified by chromatography on silica gel eluted with 1 - 16 % ethanol in dichloromethane. Yield, 150 mg, 52 %; HPLC (60/40) 3.53 minutes (99 %); PBMS 442/444 (MH+, 100 %);
- Anal. Calcd for $C_{23}H_{24}ClN_3O_4 + 0.5 H_2O$: C, 61.26; H, 5.59; N, 9.32.
- Found: C, 61.52; H, 5.89; N, 8.98.

15

Example 55a

(S)-2-Amino-3-(4-hydroxy-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride

- (S)-[1-(4-Hydroxy-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (450 mg, 1.2 mmol) was dissolved in 4M HCl-dioxane (2 mL) at 0
- 20 °C. The mixture was stirred at 25 °C for 1 hour, concentrated and the residue triturated with ether. Yield, 400 mg, 107 %; MS 265 (MH+, 100 %).

Example 55b

(S)-[1-(4-Hydroxy-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester

- 25 4-Hydroxypiperidine (3.9 mmol) and Boc-L-tyrosine (3.7 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 60 hour reaction time) with the following workup: the reaction mixture was diluted with ethyl acetate and washed once with base, the base layer was acidified with 2 N HCl and extracted three times with chloroform, and the chloroform extracts concentrated. The resulting foam was
- 30 purified by chromatography on silica gel eluted with 1 - 8 % ethanol in dichloromethane containing 0.5 % NH_4OH . Yield 550 mg; 41 %; HPLC (40/60) 5.02 minutes (87 %).

Example 56

5-Chloro-1H-indole-2-carboxylic acid [2-(4-hydroxy-piperidin-1-yl)-
2-oxo-(1S)-pyridin-3-ylmethyl-ethyl]-amide

(S)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-pyridin-3-yl-propan-1-one
5 dihydrochloride (0.8 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.7 mmol) were
coupled according to Procedure A (0 - 25 °C reaction temperature) and the product
purified by chromatography on silica gel eluted with 1-16% ethanol in dichloromethane.
Yield 26 mg, 8 %; HPLC (50/50) 5.02 minutes (99 %); PBMS 427/429 (MH+, 100 %);
Anal. Calcd for $C_{22}H_{23}ClN_4O_3 + 0.5 H_2O$: C, 60.62; H, 5.55; N, 12.85.
10 Found: C, 60.57; H, 5.74; N, 12.53.

Example 56a

(S)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-pyridin-3-yl-propan-1-one dihydrochloride

(S)-[2-(4-Hydroxy-piperidin-1-yl)-2-oxo-1-pyridin-3-ylmethyl-ethyl]-carbamic acid-
tert-butyl ester (367 mg, 1.05 mmol) was dissolved in 4M HCl-dioxane at 0 °C. The
15 resulting suspension was stirred for 1.5 hours at 25 °C, concentrated and the residue
trituated with ether. Yield, 450 mg, 100 %.

Example 56b

(S)-[2-(4-Hydroxy-piperidin-1-yl)-2-oxo-1-pyridin-3-yl-methyl-ethyl]-
carbamic acid-tert-butyl ester

20 4-Hydroxypiperidine (2.9 mmol) and N-t-Boc-L-3-(3-pyridyl)alanine (2.8 mmol)
were coupled according to Procedure A (0 - 25 °C reaction temperature, 96 hour
reaction time, acid wash omitted) and the product purified by chromatography on silica
gel eluted with 1 - 8 % ethanol in dichloromethane. Yield 454 mg, 46 %; MS 350
(MH+, 100 %).

25

Example 57

1H-Indole-2-carboxylic acid [(1R)-(4-fluoro-benzyl)-2-(4-hydroxy-
piperidin-1-yl)-2-oxo-ethyl]-amide

(R)-2-Amino-3-(4-fluoro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one
hydrochloride (0.5 mmol) and 1H-indole-2-carboxylic acid (0.5 mmol) were coupled
30 according to Procedure A (0 - 25 °C reaction temperature) and the product purified by
chromatography on silica gel eluted with 25, 30, 50, 75 and 80 % ethyl acetate in
hexanes. Yield 150 mg, 60 %; HPLC (60/40) 3.66 minutes (97 %); mp 204 - 207
°C; PBMS 410 (MH+, 100 %);

Anal. Calcd for $C_{23}H_{24}FN_3O_3$: C, 67.47; H, 5.91; N, 10.26.

Found: C, 67.18; H, 6.03; N, 10.21.

Example 57a

(R)-2-Amino-3-(4-fluoro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride

- 5 (R)-[1-(4-Fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid-tert-butyl ester (2.6 mmol) was dissolved in 4M HCl-dioxane (2 mL) at 0 °C. The solution was stirred 2 hours at 25 °C, concentrated and the residue triturated with ether. Yield, 920 mg, 124 %; HPLC (60/40) 2.23 minutes (98 %).

Example 57b

- 10 (R)-[1-(4-Fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-
carbamic acid-tert-butyl ester

- 4-Hydroxypiperidine (3.7 mmol) and (R)-N-t-Boc-p-fluoro-phenylalanine (3.5 mmol) were coupled according to Procedure A giving a foam which was used without further purification. Yield 940 mg, 73 %; HPLC (60/40) 3.64 minutes (95 %); MS 367 (MH⁺, 100 %).

Example 58

5-Chloro-1H-indole-2-carboxylic acid [(1R)-(4-fluoro-benzyl)-2-
(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

- (R)-2-Amino-3-(4-fluoro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one
20 hydrochloride (0.6 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.6 mmol) were coupled according to Procedure A and the crude product purified by chromatography on silica gel eluted with 50, 75 and 100 % ethyl acetate in hexanes. Yield 171 mg, 765 %; HPLC (60/40) 4.23 minutes (97 %); MS 444/446 (MH⁺, 100 %); TSPMS 444/446 (MH⁺, 100 %);

- 25 ¹H NMR (CDCl₃) δ 9.20 (br, 1H), 7.57 (d, 1H, J = 2 Hz), 7.33 (d, 1H, J = 8 Hz), 7.3-7.2 (m, 2H), 7.14 (m, 2H), 6.97 (m, 2H), 6.85 (m, 1H), 5.34 (m, 1H), 4.05 - 3.80 (m, 2H), 3.7 - 3.3 (m, 1.5H), 3.25 (m, 1H), 3.10 (m, 2H), 2.93 (m, 0.5H), 1.9 - 1.7 (m, 2.5H), 1.45 (m, 2H), 1.15 (m, 0.5H).

Anal. Calcd for $C_{23}H_{23}ClFN_3O_3 + 0.05 H_2O$: C, 62.11; H, 5.23; N, 9.45.

- 30 Found: C, 62.51; H, 5.66; N, 9.19.

-89-

Example 59

5-Fluoro-1H-indole-2-carboxylic acid [(1S)-(4-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

(S)-2-Amino-3-(4-fluoro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride (0.5 mmol) and 5-fluoro-1H-indole-2-carboxylic acid (0.5 mmol) were coupled according to Procedure A. The crude product was triturated once with 1:1 ether-hexanes and once with hexanes. The resulting solid was boiled in ethyl acetate, the resulting suspension filtered, and the collected solid dried. Yield 103 mg, 48 %; HPLC (60/40) 3.69 minutes (95 %); PBMS 428 (MH⁺, 100 %);

10 Anal. Calcd for C₂₃H₂₃F₂N₃O₃ + 0.25 H₂O: C, 63.95; H, 5.48; N, 9.73. Found: C, 63.93; H, 5.66; N, 9.87.

Example 59a

(S)-2-Amino-3-(4-fluoro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride

15 [(S)-1-(4-Fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (20.2 g, 55 mmol) was dissolved in 4M HCl-dioxane (25 mL) at 25 °C. After 3 hours a thick syrup had precipitated, and an additional 4M HCl-dioxanes (10 mL) was added. The mixture was stirred for 2 hours, concentrated and the solid residue suspended in 4M HCl - dioxanes. After 2 hours at 25 °C, the mixture was

20 concentrated and the residue coevaporated twice with ether. The resulting solid was stirred in a mixture of ether (75 mL) and hexanes (10 mL) at 25 °C for 18 hours, the mixture filtered, and the filtered solid washed with 1:1 ether-hexanes and dried giving a hygroscopic solid (16.3 g, 97 %).

Example 59b

25 [(S)-1-(4-Fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester

4-Hydroxypiperidine (0.29mol) and (S)-N-t-Boc-p-fluoro-phenylalanine (0.28 mol) were coupled according to Procedure A giving crude product as a foam in 84 % yield. A portion of this material (81.6 g) was dissolved in hot ethyl acetate (400 mL) and

30 hexanes (25 °C) was added to the resulting solution until slight turbidity occurred. The mixture was heated to boiling and the resulting clear solution allowed to cool to 25°C overnight. The resulting suspension was filtered and the collected solid washed with ethyl acetate-hexanes and dried (68.1 g, 67 %).

SUBSTITUTE SHEET (RULE 26)

-90-

Example 60

1-((2S)-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl)-(4R)-hydroxy-pyrrolidine-(2S)-carboxylic acid benzyl ester

- 1-((2S)-Amino-3-phenyl-propionyl)-(4R)-hydroxy-pyrrolidine-(2S)-carboxylic acid benzyl ester hydrochloride (0.56 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.53 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 60 hour reaction time) and the product purified by chromatography on silica gel eluted with 20, 30 and 50 % ethyl acetate in hexanes. Yield, 26 mg, 8 %; HPLC (60/40) 8.14 minutes (98 %); PBMS 546/548 (MH⁺, 100 %);
- 10 Anal. Calcd for C₃₀H₂₈ClN₃O₅: C, 65.99; H, 5.17; N, 7.70.
Found: C, 66.14; H, 5.37; N, 7.60.

Example 60a

1-((2S)-Amino-3-phenyl-propionyl)-(4R)-hydroxy-pyrrolidine-(2S)-carboxylic acid benzyl ester hydrochloride

- 15 1-((2S)-tert-Butoxycarbonylamino-3-phenyl-propionyl)-(4R)-hydroxy-pyrrolidine-(2S)-carboxylic acid benzyl ester (3.0 mmol) was dissolved in 4M HCl-dioxane at 0 °C. The mixture was stirred at 25 °C for 1 hour, concentrated and the residue triturated with ether. Yield 1.16 g, 96 %.

Example 60b

- 20 **1-((2S)-tert-Butoxycarbonylamino-3-phenyl-propionyl)-(4R)-hydroxy-pyrrolidine-(2S)-carboxylic acid benzyl ester**

- Trans-L-Hydroxyproline benzyl ester (3.15 mmol) and L-Boc-phenylalanine (3.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 1:1 dichloromethane / dimethylformamide) and the product used without further purification.
- 25 Yield 1.31 g, 99 %; HPLC (60/40) 6.1 minutes (95 %).

Example 61

5-Chloro-1H-indole-2-carboxylic acid [(1S)-(4-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

- (S)-2-Amino-3-(4-fluoro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride (0.051 mol) and 5-chloro-1H-indole-2-carboxylic acid (0.051 mol) were coupled according to Procedure A and the product purified by chromatography on silica gel eluted with 50 %, 75 %, 80 % and 100 % ethyl acetate-hexanes giving a foam (yield 78 %), HPLC (60/40) 4.21 minutes (99 %). A portion of this material was
- 30

-91-

recrystallized by dissolving in hot ethyl acetate (approximately 5 - 7 mL/g), and adding an approximately equal volume of hexanes at reflux, followed by slow cooling of the solution to 25 °C. The solid was filtered and washed with 1:4 ethyl acetate-hexanes and dried (70 - 90 % recovery): mp 175 - 177 °C;

- 5 ¹H NMR (CDCl₃) δ 9.41 (m, 0.5H), 9.36 (m, 0.5H), 7.59 (d, 1H, J = 2 Hz), 7.37 (d, 1H, J = 8 Hz), 7.29 (dd, 1H, J = 2, 9 Hz), 7.20 (dd, 1H, J = 2.0, 8.9 Hz), 7.14 (m, 2H), 6.95 (m, 2H), 6.86 (m, 1H), 5.34 (m, 1H), 4.05 (m, 0.5H), 3.90 (m, 1.5H), 3.65 (m, 0.5H), 3.45 (m, 1H), 3.25 (m, 1H), 3.10 (m, 2H), 2.93 (m, 0.5H), 1.88 (br, 1H, exchanges with D₂O), 1.80 (m, 1.5H), 1.45 (m, 2H), 1.12 (m, 0.5H).
- 10 PBMS 444/446 (MH⁺, 100 %);
 Anal. Calcd for C₂₃H₂₃ClFN₃O₃ + 0.2 H₂O: C, 61.73; H, 5.27; N, 9.39.
 Found: C, 61.40; H, 5.37; N, 9.11.

Example 62

5-Chloro-1H-indole-2-carboxylic acid [(1S)-(methoxy-methyl-carbamoyl)-2-pyridin-3-yl-ethyl]-amide

- 15 (2S)-Amino-N-methoxy-N-methyl-3-pyridin-3-yl-propionamide dihydrochloride (1.3 mmol) and 5-chloro-1H-indole-2-carboxylic acid (1.25 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 1:1 dichloromethane / DMF reaction solvent) and the product purified by chromatography on silica gel eluted with ethyl acetate. Yield 313 mg, 65 %; HPLC (60/40) 2.84 minutes (99 %); TSPMS 387/389 (MH⁺, 100 %);
- 20 ¹H NMR (CDCl₃) δ 9.1 (br, 1H), 8.48 (dd, 1H), 8.43 (m, 1H), 7.60 (d, 1H), 7.50 (m, 1H, J = ca. 8 Hz), 7.37 (d, 1H, J = ca. 8 Hz), 7.23 (d, 1H), 7.18 (dd, 1H, J = ca. 8 Hz), 7.10 (d, 1H, J = ca. 8 Hz), 6.82 (d, 1H), 5.42 (m, 1H), 3.78 (s, 3H), 3.25 (s, 3H), 3.32 (dd, A of AB, 1H, J = ca. 7, 14 Hz), 3.10 (dd, B of AB, 1H, J = ca. 7, 14 Hz).
- 25 Anal. Calcd for C₁₉H₁₉ClN₄O₃ + 0.4 H₂O: C, 57.91; H, 5.07; N, 14.22.
 Found: C, 58.19; H, 5.23; N, 13.82.

Example 62a

(2S)-Amino-N-methoxy-N-methyl-3-pyridin-3-yl-propionamide dihydrochloride

- 30 [(1S)-(Methoxy-methyl-carbamoyl)-2-pyridin-3-yl-ethyl]-carbamic acid tert-butyl ester (1.5 mmol) was dissolved in 4M HCl-dioxane at 0 °C. The resulting solution was stirred for 2 hours at 25 °C, concentrated and the residue triturated with ether. Yield, 390 mg, 95 %.

-92-

Example 62b

(S)-[1-(Methoxy-methyl-carbamoyl)-2-pyridin-3-yl-ethyl]-carbamic acid tert-butyl ester

N,O-Dimethylhydroxylamine hydrochloride (1.7 mmol) and Boc-3-pyridyl-L-alanine (1.6 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 2:1 dichloromethane/dimethylformamide reaction solvent, acid wash omitted, Na₂SO₄ used for drying). The residue was triturated with ether giving 428 mg (86 % yield) of a yellow solid.

Example 63

(R,S)-2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-(3-fluoro-phenyl)-propionic acid methyl ester

(R,S)-2-Amino-3-(3-fluoro-phenyl)-propionic acid methyl ester (2.05 mmol) and 5-chloro-1H-indole-2-carboxylic acid (2.03 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 1:1 dichloromethane/DMF reaction solvent) and the product purified by chromatography on silica gel eluted with 10, 20 and 40 % ethyl acetate in hexanes. The residue was triturated with 1:1 ether-hexanes, and hexanes giving an off-white solid (484 mg, 63 %): HPLC (60/40) 8.13 minutes (95 %); TSPMS 375/377 (MH⁺, 100 %);

¹H NMR (CDCl₃) δ 9.26 (br, 1H), 7.60 (d, 1H, J = ca. 1 Hz), 7.35 (d, 1H, J = 8.7 Hz), 7.25 (m, 2H), 6.95 (m, 1H), 6.91 (m, 1H), 6.84 (m, 1H), 6.77 (d, 1H, J = 1.5 Hz), 6.63 (d, 1H, J = 7.7 Hz), 5.08 (m, 1H), 3.78 (s, 3H), 3.28 (dd, 1H, A of AB, J = 5.7, 14 Hz), 3.21 (dd, 1H, B of AB, J = 5.5, 14 Hz).

Anal. Calcd for C₁₉H₁₆ClFN₂O₃: C, 60.89; H, 4.30; N, 7.47.

Found: C, 60.79; H, 4.58; N, 7.18.

Example 63a

(R,S)-2-Amino-3-(3-fluoro-phenyl)-propionic acid methyl ester hydrochloride

Trimethylsilylchloride (1.07 g, 9.9 mmol) was added to a suspension of m-fluoro-DL-phenylalanine (0.404 g, 2.2 mmol) in methanol (4 mL) at 25°C. The resulting solution was brought to reflux for 1 hour, cooled and concentrated. The residue was triturated with ether. Yield, 515 mg, 100 %; HPLC (60/40) 2.31 minutes (95 %).

-93-

Example 64**5-Chloro-1H-indole-2-carboxylic acid [(1S)-(methoxy-methyl-carbamoyl)-2-thiophen-2-yl-ethyl]-amide**

- (S)-2-Amino-N-methoxy-N-methyl-3-thiophen-2-yl-propionamide hydrochloride (1.2 mmol) and 5-chloro-1H-indole-2-carboxylic acid (1.2 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 2:1 dichloromethane / dimethylformamide reaction solvent). The crude product was purified by chromatography on silica gel eluted with 10, 20, 30 and 40 % ethyl acetate in hexanes. Yield 375 mg, 80 %; HPLC (60/40) 6.36 minutes (99 %); PBMS 392/394 (MH+, 100 %);
- 10 ¹H NMR (CDCl₃) δ 9.33 (br, 1H), 7.60 (d, 1H, J = ca. 1 Hz), 7.30 (d, 1H, J = 8.8 Hz), 7.20 (dd, 1H, J = 2.0, 8.7 Hz), 7.15 (dd, 1H, J = 1, 5.0 Hz), 6.91 (dd, 1H, J = 3.4, 5.1 Hz), 6.86 (d, 1H, J = 1.6 Hz), 6.84 (d, 1H, J = ca 2 Hz), 5.40 (m, 1H), 3.77 (s, 3H), 3.46 (dd, 1H, A of AB, J = 6.2, ca. 14 Hz), 3.37 (dd, 1H, B of AB, J = 6.2, ca. 14.2 Hz), 3.25 (s, 3H).
- 15 Anal. Calcd for C₁₈H₁₈ClN₃O₃S + 0.25 C₄H₈O₂: C, 55.14; H, 4.87; N, 10.15. Found: C, 55.41; H, 4.79; N, 10.17.

Example 64a**(S)-2-Amino-N-methoxy-N-methyl-3-thiophen-2-yl-propionamide hydrochloride**

- (S)-[1-(Methoxy-methyl-carbamoyl)-2-thiophen-2-yl-ethyl]-carbamic acid tert-butyl ester (1.3 mmol) was dissolved in 4 M HCl - dioxane (1 mL) at 0°C and the resulting solution stirred at 25°C for 2 hours. The mixture was concentrated and the residue triturated with ether giving a yellow solid (321 mg, 96 %; HPLC (60/40) 2.24 minutes (98 %); MS 215 (MH+, 100 %).
- 20

Example 64b

- 25 **(S)-[1-(Methoxy-methyl-carbamoyl)-2-thiophen-2-yl-ethyl]-carbamic acid tert-butyl ester**

- N,O-Dimethylhydroxylamine hydrochloride (1.4 mmol) and Boc-(2-thienyl)-L-alanine (1.3 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) giving the product which was used without further purification. Yield 426
- 30 mg, 104 %.

SUBSTITUTE SHEET (RULE 26)

Example 65

(R,S)-2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-(4-fluoro-phenyl)-
propionic acid methyl ester

(R,S)-2-Amino-3-(3-fluoro-phenyl)-propionic acid methyl ester (3.0 mmol) and 5-
 5 chloro-1H-indole-2-carboxylic acid (2.9 mmol) were coupled according to Procedure A
 (0 - 25 °C reaction temperature, 3:2 dichloromethane / dimethylformamide reaction
 solvent) and the resulting crude product triturated with 1:1 ether / hexanes. Yield 1.03
 g, 92 %; HPLC (60/40) 7.95 minutes (96 %); PBMS 375/377 (MH+, 100 %);
¹H NMR (CDCl₃) δ 9.30 (br, 1H), 7.60 (d, 1H, J = ca. 1 Hz), 7.35 (d, 1H, J = 8.8 Hz),
 10 7.25 (dd, 1H, J = 2.0, 8.7 Hz), 7.10 (m, 2H), 6.97 (m, 2H), 6.77 (d, 1H, J = 2 Hz), 6.62
 (d, 1H, J = 7.8 Hz), 5.06 (m, 1H), 3.78 (s, 3H), 3.27 (dd, 1H, A of AB, J = 7, 14 Hz),
 3.19 (dd, 1H, B of AB, J = 7, 14 Hz).
 Anal. Calcd for C₁₉H₁₆ClFN₂O₃: C, 60.89; H, 4.30; N, 7.47.
 Found: C, 60.74; H, 4.36; N, 7.55.

15

Example 66

5-Chloro-1H-indole-2-carboxylic acid [2-(4-amino-phenyl)-(1S)-
dimethylcarbamoyl-ethyl]-amide hydrochloride

(S)-2-Amino-3-(4-amino-phenyl)-N,N-dimethyl-propionamide dihydrochloride
 (0.7 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.7 mmol) were coupled according
 20 to Procedure A (0 - 25 °C reaction temperature, 3:1 dichloro-methane / DMF reaction
 solvent, washed with base only) and the product purified by chromatography on silica
 gel eluted with 1 - 16 % ethanol in dichloromethane with 0.5 % NH₄OH. The combined
 fractions were concentrated, dissolved in methanol at 0 °C, the resulting solution
 treated with 1.01 N HCl (1.05 eq). After 5 minutes, the reaction mixture was
 25 concentrated and the residue triturated with ether giving and orange solid (79 mg, 29
 % yield): TSPMS 385/387 (MH+, 100 %);
 Anal. Calcd for C₂₀H₂₁ClN₄O₂ + 1.5 HCl: C, 54.65; H, 5.16; N, 12.75.
 Found: C, 54.96; H, 5.53; N, 12.53.

Example 66a

30

(S)-2-Amino-3-(4-amino-phenyl)-N,N-dimethyl-propionamide dihydrochloride

(S)-[2-(4-Amino-phenyl)-1-dimethylcarbamoyl-ethyl]-carbamic acid tert-butylester
 (214 mg, 0.7 mmol) was dissolved in 4M HCl-dioxane (2 mL) at 0 °C and the solution

-95-

stirred for 2 hours at 25 °C. The mixture was concentrated and the residue triturated with ether. Yield, 294 mg, 102 %; PBMS 208 (MH+, 100 %).

Example 66b

(S)-[2-(4-Amino-phenyl)-1-dimethylcarbamoyl-ethyl]-carbamic acid tert-butyl ester

5 Dimethylamine hydrochloride (2.04 mmol) and Boc-p-amino-L-phenylalanine (1.7 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 4:1 dichloro-methane/dimethylformamide reaction solvent, washed with base only). The product was purified by chromatography on silica gel eluted with 50, 60, 70 and 100 % ethyl acetate in hexanes. Yield 226 mg, 42 %; HPLC (70/30) 2.45 minutes (100
10 %).

Example 67

5-Chloro-1H-indole-2-carboxylic acid((1S)-dimethylcarbamoyl-3-phenylpropyl)-amide

(S)-2-Amino-N,N-dimethyl-4-phenyl-butyramide hydrochloride (0.76 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.76 mmol) were coupled according to Procedure
15 A (0 - 25 °C reaction temperature, 3:1 dichloromethane / DMF reaction solvent) and the product purified by chromatography on silica gel eluted with 10, 20, 30, 40, 50 and 60 % ethyl acetate in hexanes. Yield 263 mg, 90 %; HPLC (60/40) 7.12 minutes (99 %); TSPMS 384/386 (MH+, 100 %);
Anal. Calcd for C₂₁H₂₂ClN₃O₂: C, 65.71; H, 5.78; N, 10.95,
20 Found: C, 65.34; H, 5.93; N, 10.91.

Example 67a

(S)-2-Amino-N,N-dimethyl-4-phenyl-butyramide hydrochloride

(S)-(1-Dimethylcarbamoyl-3-phenyl-propyl)-carbamic acid tert-butyl ester (235 mg, 0.8 mmol) was dissolved in 4 M HCl - dioxane (2 mL) at 0 °C. The mixture was
25 stirred at 25 °C for 1.5 hours, concentrated and the residue triturated with ether. Yield, 187 mg, 100 %; HPLC (60/40) 2.31 minutes (99 %).

Example 67b

(S)-(1-Dimethylcarbamoyl-3-phenyl-propyl)-carbamic acid tert-butyl ester

Dimethylamine hydrochloride (1.0 mmol) and (S)-N-t-butoxycarbonyl-2-amino-4-
30 phenylbutyric acid (0.84 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 3:1 dichloromethane / DMF reaction solvent) giving the product which was used without further purification. Yield 238 mg, 93 %; HPLC (60/40) 5.98 minutes (97 %).

-96-

Example 68

5-Chloro-1H-indole-2-carboxylic acid [(1S)-dimethyl-carbamoyl-
2-(4-hydroxy-phenyl)-ethyl]-amide

- (S)-2-Amino-3-(4-hydroxy-phenyl)-N,N-dimethyl-propionamide hydrochloride (1.05 mmol) and 5-chloro-1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 2:1 dichloro-methane/DMF reaction solvent, washed with acid only) and the product purified by chromatography on silica gel eluted with 20, 40, 50 and 75 % ethyl acetate in hexanes followed by trituration with ether. Yield 400 mg, 104 %; HPLC (60/40) 3.93 minutes (98 %); mp 228 - 231 °C (dec, yellowed at 210 °C); TSPMS 386/388 (MH+, 100 %);
- Anal. Calcd for $C_{20}H_{20}ClN_3O_3 + 0.9 H_2O$: C, 59.75; H, 5.47; N, 10.45.
- Found: C, 61.05; H, 5.79; N, 10.08.

Example 68a

(S)-2-Amino-3-(4-hydroxy-phenyl)-N,N-dimethyl-propionamide hydrochloride

- (S)-[1-Dimethylcarbamoyl-2-(4-hydroxy-phenyl)-ethyl]-carbamic acid tert-butyl ester (5.7 g, 18.5 mmol) was dissolved in 4M HCl-dioxane (7 mL) at 0 °C. The mixture was stirred at 25 °C for 3 hours, concentrated and the residue trituated with ether. Yield, 5.23 g; HPLC (60/40) 3.32 minutes (98 %).

Example 68b

- (S)-[1-Dimethylcarbamoyl-2-(4-hydroxy-phenyl)-ethyl]-carbamic acid tert-butyl ester

- Dimethylamine hydrochloride (79 mmol) and Boc-L-tyrosine (66 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 12:1 dichloromethane / DMF reaction solvent, 60 hour reaction time) and the product purified by chromatography on silica gel eluted with 10, 20, 30, 50 and 70 % ethyl acetate in hexanes. Yield 20.6 g, 102 %; HPLC (60/40) 3.21 minutes (96 %).

Example 69

5-Chloro-1H-indole-2-carboxylic acid ((1S)-methoxycarbamoyl-2-phenyl-ethyl)-amide

- (2S)-Amino-N-methoxy-3-phenyl-propionamide hydro-chloride (1.02 mmol) and 5-chloro-1H-indole-2-carboxylic acid (1.02 mmol) were coupled according to Procedure A. The residue was trituated with ether to give a light yellow solid. Yield, 160 mg, 36 %; mp 210 - 213 °C (dec); PBMS 372/374 (MH+, 100 %);
- Anal. Calcd for $C_{19}H_{18}ClN_3O_3 + 1.75 H_2O$: C, 56.58; H, 5.37; N, 10.42.
- Found: C, 56.88; H, 5.09; N, 10.03.

-97-

Example 69a**(2S)-Amino-N-methoxy-3-phenyl-propionamide hydrochloride**

[(1S)-(Methoxy-carbamoyl)-2-phenyl-ethyl]-carbamic acid tert-butyl ester (200 mg, 0.68 mmol) was dissolved in 4 M HCl - dioxane at 0 °C and the mixture stirred at 25 °C.
5 After 0.5 hours, the mixture was concentrated and the residue triturated with ether.

Example 69b**[(1S)-(Methoxy-carbamoyl)-2-phenyl-ethyl]-carbamic acid tert-butyl ester**

Methoxyamine hydrochloride (83.5 mmol) and Boc-L-phenylalanine (20 mmol) were coupled according to Procedure A and the product purified by chromatography
10 on silica gel eluted with 1:1 and 2:1 ethyl acetate / hexanes followed by trituration with ether. Yield 1.80 g, 31 %.

Example 70**5-Chloro-1H-indole-2-carboxylic acid ((1R)-methylcarbamoyl-2-phenyl-ethyl)-amide**

(R)-2-Amino-N-methyl-3-phenyl-propionamide hydrochloride (0.84 mmol) and 5-
15 chloro-1H-indole-2-carboxylic acid (0.84 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature). The crude product was triturated with dichloromethane and then with ether and dried. Yield 236 mg, 79 %; HPLC (60/40) 4.63 minutes (97 %); PBMS 356/358 (MH+, 100 %);
Anal. Calcd for C₁₉H₁₈ClN₃O₂ + 0.25 H₂O: C, 63.33; H, 5.18; N, 11.66.
20 Found: C, 63.37; H, 5.50; N, 12.06.

Example 70a**(R)-2-Amino-N-methyl-3-phenyl-propionamide hydrochloride**

(R)-(1-Methylcarbamoyl-2-phenyl-ethyl)-carbamic acid tert-butyl ester (722 mg, 2.6 mmol) was dissolved in 4M HCl-dioxane (10 mL) at 0 °C. The mixture was stirred
25 for 1 hour at 25 °C, concentrated and the residue triturated with ether. Yield, 517 mg, 93 %.

Example 70b**(R)-(1-Methylcarbamoyl-2-phenyl-ethyl)-carbamic acid tert-butyl ester**

Methylamine hydrochloride (3.1 mmol) and Boc-D-phenyl-alanine (2.8 mmol)
30 were coupled according to Procedure A (0 - 25 °C reaction temperature, 144 hour reaction time, washed with acid first, then base) giving the product which was used without further purification. Yield 760 mg, 96 %.

-98-

Example 71**5,6-Dichloro-1H-indole-2-carboxylic acid****((1S)-dimethylcarbamoyl-2-phenyl-ethyl)-amide**

- (S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride (0.06 mmol) and
5 5,6-dichloro-1H-indole-2-carboxylic acid (0.06 mmol) were coupled according to
Procedure A (0 - 25 °C reaction temperature, 96 hour reaction time). The crude
product was triturated with 1:1 ether-hexanes and dried. Yield 24 mg, 96 %; HPLC
(60/40) 8.05 minutes (97 %); PBMS 405/407 (MH+, 100 %);
Anal. Calcd for $C_{20}H_{19}Cl_2N_3O_2 + 0.25 H_2O$: C, 58.76; H, 4.81; N, 10.28.
10 Found: C, 58.95; H, 4.89; N, 9.90.

Example 71a**(S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride**

- (1-Dimethylcarbamoyl-2-phenyl-ethyl)-carbamic acid tert-butyl ester (8.6 g, 29
mmol) was dissolved in 4 M HCl - dioxane (110 ml) at 0 °C and the mixture stirred at
15 25°C for 1 hour. The mixture was concentrated and the solids triturated with ether.
Yield, 6.2 g, 92 %; PBMS 193 (MH+, 100 %).

Example 71b**5,6-Dichloro-1H-indole-2-carboxylic acid**

- Zinc dust (3.52 g, 54 mmol) was added slowly to a warm solution of 3,4-
20 dichloro-5-nitrophenylpyruvic acid (1.5 g, 5.4 mmol) in acetic acid (15 mL). After a few
minutes, a vigorous reaction occurred (exothermic). The resulting solution was heated
to 80 °C and the reaction appeared complete (TLC). The mixture was filtered, the
filtered solids washed with acetic acid and the filtrate concentrated. The residue was
dissolved in 2 N NaOH, the resulting solution washed with ether (3x), dichloromethane
25 (2x) and acidified to pH 1 with 6N HCl and extracted with ethyl acetate. The extracts
were dried and concentrated giving a light brown solid (458 mg, 34 %): HPLC (60/40)
5.31 (93 %).

Example 71c**3,4-dichloro-5-nitrophenylpyruvic acid**

- 30 Absolute ethanol (25 mL) was added at 3 - 15 °C to a stirred mixture of
potassium metal (2.67 g, 68 mmol) in ether (100 mL). The resulting solution was
treated at 3 °C with a solution of diethyl oxalate (10.0 g, 62 mmol) over 5 - 10 minutes,
and the resulting solution stirred 30 minutes at 3 °C and 25 °C for 18 hours. The

-99-

mixture was filtered and the resulting solid washed with ether and dried (13.7 g). This material (12.7 g) was dissolved in 400 mL hot water, the solution cooled and extracted with ether. The resulting aqueous layer was acidified to pH 2 with conc. HCl and the ether layer separated, dried and concentrated giving 7.5 g of a solid which was
5 triturated with hexanes giving the title substance as a yellow solid (7.01 g 41 %).

Example 725-Bromo-1H-indole-2-carboxylic acid((1S)-dimethylcarbamoyl-2-phenyl-ethyl)-amide

(S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride (1.0 mmol) and
10 5-bromo-1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) and the resulting foam triturated with 1:1 ether/hexanes and dried. Yield 374 mg, 90 %; HPLC (60/40) 6.17 minutes (98 %); mp 199 - 201 °C; PBMS 414/416 (MH+, 100 %);
Anal. Calcd for $C_{20}H_{20}BrN_3O_2$: C, 57.98; H, 4.82; N, 10.14.
15 Found: C, 58.07; H, 5.12; N, 10.08.

Example 735-Methyl-1H-indole-2-carboxylic acid((1S)-dimethylcarbamoyl-2-phenyl-ethyl)-amide

(S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride (1.0 mmol) and
20 5-methyl-1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature). The crude product was triturated with 1:1 ether-hexanes and dried. Yield 302 mg, 87 %; HPLC (60/40) 5.46 minutes (99 %); mp 198.5 - 200 °C; PBMS 350 (MH+, 100 %);
Anal. Calcd for $C_{21}H_{23}N_3O_2$: C, 72.18; H, 6.63; N, 12.04.
25 Found: C, 72.14; H, 6.90; N, 12.11.

Example 745-Methoxy-1H-indole-2-carboxylic acid((1S)-dimethylcarbamoyl-2-phenyl-ethyl)-amide

(S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride (1.0 mmol) and
30 5-methoxy-1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 60 hour reaction time) and the resulting foam triturated with ether. Yield 329 mg, 90 %; HPLC (60/40) 4.27 minutes (99 %); PBMS 366 (MH+, 100 %);

BEST AVAILABLE COPY

-100-

Anal. Calcd for $C_{21}H_{23}N_3O_3 + 0.125 H_2O$: C, 68.60; H, 6.37; N, 11.43.

Found: C, 68.50; H, 6.34; N, 11.45.

Example 75

5-Fluoro-1H-indole-2-carboxylic acid

5

((1S)-dimethylcarbamoyl-2-phenyl-ethyl)-amide

(S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride (1.0 mmol) and 5-fluoro-1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature, 60 hour reaction time) and the resulting solid triturated with ether. Yield 320 mg, 91 %; HPLC (60/40) 4.74 minutes (100 %); mp

10 229.5 - 232 °C; PBMS 354 (MH+, 100 %);

Anal. Calcd for $C_{20}H_{20}FN_3O_2$: C, 67.97; H, 5.70; N, 11.89. Found: C, 67.88; H, 5.74; N, 11.71.

Example 76

5-Cyano-1H-indole-2-carboxylic acid

15

((1S)-dimethylcarbamoyl-2-phenyl-ethyl)-amide

(S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride (0.16 mmol) and 5-cyano-1H-indole-2-carboxylic acid (0.16 mmol) were coupled according to Procedure A (0 - 25 °C reaction temperature) and the product purified by chromatography on silica gel eluted with 1:1 ethyl acetate / hexanes. Yield 38 mg, 66 %; HPLC (60/40)

20 4.08 minutes (97 %); PBMS 361 (MH+, 100 %);

1H NMR (DMSO- d_6) δ 12.1 (br, 1H), 9.04 (d, 1H, $J = 8.1$ Hz), 8.27 (s, 1H), 7.52 (m, 2H), 7.43 (m, 1H), 7.33 (m, 2H), 7.25 (m, 2H), 7.18 (m, 1H), 5.10 (m, 1H), 3.03 (m, 2H), 3.00 (s, 3H), 2.83 (s, 3H).

Anal. Calcd for $C_{21}H_{20}N_4O_2 + 0.5 H_2O$: C, 68.28; H, 5.73; N, 15.17.

25 Found: C, 68.51; H, 5.66; N, 14.85.

Example 76a

5-Cyano-1H-indole-2-carboxylic acid

5-Cyano-1H-indole-2-carboxylic acid ethyl ester (1.71 g, 8.0 mmol) was added to a solution of ethanol (10 mL) and potassium hydroxide (2 g) and the resulting mixture heated at reflux for 1 hour. Water was added to dissolve the precipitate, and 6N HCl was added to bring the pH to 1. The mixture was cooled in an ice bath, filtered, and the resulting colorless solid washed with cold water and dried (1.51 g). A portion (1.4 g) was suspended in hot acetic acid (40 mL) and cooled giving a solid

-101-

which was filtered, washed with cold ethyl acetate and dried: Yield 980 mg 70 %; HPLC (60/40) 3.09 minutes (97 %).

Example 76b

5-Cyano-1H-indole-2-carboxylic acid ethyl ester

5 Zinc dust (57.8 g, 887 mmol) was added to a hot suspension of 3-cyano-5-nitrophenylpyruvic acid ethyl ester (23.2 g, 88 mmol) in acetic acid (225 mL) and water (225 mL, Caution!, vigorous initial exotherm) at a rate to maintain reflux, and the reaction was held at reflux for 0.5 hours. The mixture was filtered, the filtered salts washed with hot acetic acid (150 mL), and the filtrate chilled overnight giving crystals
10 which were filtered, washed with cold 1:1 acetic acid - water, water, and dried (10.11 g, 53 %). The filtrate was concentrated, the residue dissolved in ethyl acetate, and the resulting solution washed with sat. aqueous sodium bicarbonate, brine, dried and concentrated giving a second batch (5.05 g). The major lot was used in subsequent transformations.

15

Example 76c

3-Cyano-5-nitrophenylpyruvic acid ethyl ester

A solution of sodium ethoxide in ethanol (from 2.2 g, 400 mmol sodium metal in 400 ml ethanol) was added at 0 °C to a mixture of distilled diethyl oxalate (120 g, 821 mmol) and 3-methyl-4-nitrobenzonitrile (32 g, 197 mmol). The resulting red solution
20 was heated at 40 °C for 18 hours. The cooled mixture was diluted with water (600 mL) water and acidified with conc. HCl to pH 2.1. The precipitate that formed was collected by filtration of the 13 °C mixture, dried and purified by chromatography on silica eluted with 15, 30 and 50% acetone-hexanes giving an orange solid which was used without purification (23.6 g, 31 %). A sample was recrystallized from ethyl acetate
25 for characterization.

Example 77

1H-Indole-2-carboxylic acid ((1S)-dimethylcarbamoyl-2-phenyl-ethyl)-amide

(S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride (1.0 mmol) and 1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25
30 °C reaction temperature). The resulting solid was triturated with hexanes, then with ether. Yield 272 mg, 81 %; HPLC (70/30) 3.49 minutes (99 %); mp 199 - 200 °C; PBMS 336 (MH+, 100 %);
Anal. Calcd for C₂₀H₂₁N₃O₂: C, 71.62; H, 6.31; N, 12.53.

Found: C, 71.45; H, 6.39; N, 12.50.

Example 78

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3S,4S)-
dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide

- 5 (3S,4S)-2-Amino-1-(3,4-dihydroxy-pyrrolidin-1-yl)-3-phenyl-propan-1-one hydrochloride (0.94 mmol) and 5-chloro-1H-indole-2-carboxylic acid (1.03 mmol) were coupled according to procedure A (170 hour reaction time) and the crude product purified by column chromatography on silica gel eluted with ethyl acetate. Yield 150 mg, 37 %; HPLC (60/40) 3.08 minutes (96 %);
- 10 ¹H NMR (DMSO-d₆) δ 11.73 (s, 1H), 8.90 (d, 1H, J = 8.5 Hz), 7.72 (d, 1H, J = 1.5 Hz), 7.39 (d, 1H, J = 8.7 Hz), 7.30 (m, 2H), 7.30-7.1 (m, 5H), 5.22 (m, 1H), 5.13 (m, 1H), 4.91 (m, 1H), 3.97 (m, 1H), 3.91 (m, 1H), 3.60 (m, 2H), 3.5 - 3.2 (m, 2H) 3.00 (m, 2H). Anal. Calcd for C₂₂H₂₂ClN₃O₄: C, 61.75; H, 5.18; N, 9.82. Found: C, 61.65; H, 5.45; N, 9.17.

15

Example 78a

(3S,4S)-2-Amino-1-(3,4-dihydroxy-pyrrolidin-1-yl)-
3-phenyl-propan-1-one hydrochloride

- (3S,4S)-[1-Benzyl-2-(3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (360 mg, 1.00 mmol) was dissolved in 4 M HCl - dioxane (4 ml) at
- 20 25 °C for 3 hours. The mixture was concentrated and the resulting yellow solid triturated with ether and dried. Yield 304 mg, 103 %.

Example 78b

[1-Benzyl-2-(3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-
carbamic acid tert-butyl ester

- 25 Boc-L-Phenylalanine (2.2 mmol) and (3S,4S)-dihydroxy-pyrrolidine (US Patent No. 4634775, example 1C, 206 mg, 2.0 mmol) were coupled according to procedure A (0 - 25 °C reaction temperature) giving a colorless solid which was used without further purification. Yield 431 mg, 61 %.

Example 79

30

5-Chloro-1H-indole-2-carboxylic acid

[(1S)-benzyl-2-((3RS)-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

- 2(S)-Amino-1-((3RS)-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one hydrochloride (570 mg, 2.0 mmol) and 5-chloro-1H-indole-2-carboxylic acid (429 mg, 2.2 mmol) were

-103-

coupled according to procedure A (5:2 dichloromethane - dimethylformamide solvent) and the crude product triturated with 1:1 ether - hexanes. The resulting solid was purified by column chromatography on silica gel eluted with 3:2, and 2:1 ethyl acetate / hexanes followed by trituration with 1:1 ether / hexanes. Yield 430 mg, 51 %: HPLC

5 (60/40) 3.45 minutes (95 %);

Anal. Calcd for $C_{23}H_{24}ClN_3O_3 + 0.125 C_6H_{14}$: C, 65.32; H, 5.94; N, 9.62.

Found: C, 65.01; H, 6.19; N, 9.22.

Example 79a

(2S)-Amino-1-(3-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one hydrochloride

10 [(1S)-Benzyl-2-((3RS)-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (7.13 g, 20 mmol) was dissolved in 4M HCl-dioxane (40 ml) at 25°C for 3 hours. The mixture was concentrated and the resulting oil stirred under ether for 72 hours. The resulting suspension was filtered and the solid washed with ether and dried. Yield 5.64 g, 99 %.

15

Example 79b

[(1S)-Benzyl-2-((3RS)-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester

BOC-L-Phenylalanine (8.17 g, 30.8 mmol) and 3-hydroxypiperidine hydrochloride (4.24 g, 30.8 mmol) were coupled according to procedure A, giving the title compound as an oil which was used without further purification. Yield 7.79 g, 73 %.

20

Example 80

5-Chloro-1H-indole-2-carboxylic acid

[(1S)-benzyl-2-oxo-2-(3-oxo-piperazin-1-yl)-ethyl]-amide

4-((2S)-Amino-3-phenyl-propionyl)-piperazin-2-one hydrochloride (140 mg, 0.5 mmol) and 5-chloro-1H-Indole-2-carboxylic acid (98 mg, 0.5 mmol) were coupled according to procedure A and the crude product purified by column chromatography on silica gel eluted with ethyl acetate and 2 % ethanol in ethyl acetate followed by trituration with ether. Yield 71 mg, 33 %: HPLC (60/40) 3.53 minutes (100 %); PBMS 425/427 (MH+, 100 %);

¹H NMR (DMSO-d₆) δ 11.78 (br, 0.5H), 11.76 (br, 0.5H), 9.03 (m, 0.5H), 9.02 (m, 0.5H), 8.06 (m, 0.5H), 8.04 (m, 0.5H), 7.73 (d, 1H, J = 2 Hz), 7.38 (d, 1H, J = 8.7 Hz), 7.32 (m, 2H), 7.20 (m, 2H), 7.2 - 7.1 (m, 2H), 5.15 (m, 0.5H), 5.05 (m, 0.5H), 4.20 (d, 0.5H, J = 17 Hz), 4.08 (d, 0.5H, J = 17 Hz), 3.85 (d, 0.5H, J = 17 Hz), 3.9 (m, 0.5H), 3.6 (m, 2H), 3.2-2.9 (m, 4H).

-104-

Example 80a4-((2S)-Amino-3-phenyl-propionyl)-piperazin-2-one hydrochloride

[(1S)-Benzyl-2-oxo-2-(3-oxo-piperazin-1-yl)-ethyl]-carbamic acid tert-butyl ester (400 mg, 1.2 mmol) was dissolved in 4M HCl-dioxane (10 ml) at 25 °C for 0.5 hours.

- 5 The mixture was concentrated and the residue co-evaporated with dichloromethane, triturated with ether, and dried. Yield 340 mg, 103 %.

Example 80b[(1S)-Benzyl-2-oxo-2-(3-oxo-piperazin-1-yl)-ethyl]-carbamic acid tert-butyl ester

- BOC-L-Phenylalanine (530 mg, 2 mmol) and piperazin-2-one (J. Am. Chem. Soc. 10 62 1202 (1940), 200 mg, 2 mmol) were coupled according to procedure A (2:1 dichloromethane / dimethylformamide reaction solvent, washed with 1 N NaOH after acid washes) and the product used without further purification. Yield 404 mg, 58 %.

Example 815-Chloro-1H-indole-2-carboxylic acid

- 15 [(1S)-methyl-2-morpholin-4-yl-2-oxo-ethyl]-amide

(2S)-Amino-1-morpholin-4-yl-propan-1-one hydrochloride (195 mg, 1.0 mmol) and 5-chloro-1H-indole-2-carboxylic acid (195 mg, 1.0 mmol) were coupled according to procedure A (washed with 1N NaOH after acid washes) giving crude product which was triturated with ether and dried. Yield 150 mg, 45 %; HPLC (60/40) 3.61 minutes

- 20 (100 %); PBMS 336/338 (MH+, 100 %);

Anal. Calcd for C₁₆H₁₈ClN₃O₃: C, 57.23; H, 5.40; N, 12.51.

Found: C, 57.01; H, 5.49; N, 12.24.

Example 81a(2S)-Amino-1-morpholin-4-yl-propan-1-one hydrochloride

- 25 ((1S)-Methyl-2-morpholin-4-yl-2-oxo-ethyl)-carbamic acid tert-butyl ester (3.88 g, 15 mmol) was dissolved in 4M HCl-dioxane (20 ml) at 25°C for 1.25 hours. The mixture was concentrated and the residue triturated with ether and dried: Yield 2.51 g, 86 %.

Example 81b[(1S)-Methyl-2-morpholin-4-yl-2-oxo-ethyl]-carbamic acid

- 30 tert-butyl ester

BOC-L-Alanine (3.50 mg, 20 mmol) and morpholine (1.74 g, 20 mmol) were coupled according to procedure A (washed with 1N NaOH after acid washes) giving a colorless oil which was used without further purification. Yield 3.94 g, 76 %.

-105-

Example 82

5-Chloro-1H-indole-2-carboxylic acid ((1S)-
methylcarbamoyl-2-phenyl-ethyl)-amide

(2S)-Amino-N-methyl-3-phenyl-propionamide hydrochloride (214 mg, 1.0 mmol)
5 and 5-chloro-1H-Indole-2-carboxylic acid (195 mg, 1.0 mmol) were coupled according to procedure A and the crude product triturated with ether and dried. Yield 160 mg, 45 %; HPLC (60/40) 4.60 minutes (100 %);
¹H NMR (DMSO-d₆) δ 11.70 (br, 1H), 8.73 (d, 1H, J = 8.5 Hz), 8.08 (q, 1H, J = 4.6 Hz), 7.72 (d, 1H, J = 1.9 Hz), 7.39 (d, 1H, J = 8.8 Hz), 7.32 (m, 2H), 7.25 - 7.10 (m, 10 5H), 4.68 (m, 1H), 3.10 (dd, A of AB, 1H, J = 4.2, 13 Hz), 2.96 (dd, 1H, J = 10.7, 13 Hz), 2.62 (d, 3H, J = 4.6 Hz).

Example 82a

(2S)-Amino-N-methyl-3-phenyl-propionamide hydrochloride

(((1S)-1-Methylcarbamoyl-2-phenyl-ethyl)-carbamic acid tert-butyl ester (2.35 g, 15 8.45 mmol) was dissolved in 4 M HCl - dioxane (20 ml) at 25 °C for 2 hours. The mixture was concentrated and the residue triturated with ether, and dried. Yield 1.70 g, 94 %.

Example 82b

((1S)-1-Methylcarbamoyl-2-phenyl-ethyl)-carbamic acid tert-butyl ester

20 BOC-L-phenylalanine (2.65 g, 10 mmol) and methylamine hydrochloride (675 mg, 10 mmol) were coupled according to procedure A (washed with 1N NaOH after acid washes) yielding the title compound as a colorless solid which was used without further purification. Yield 2.41 g, 87 %; HPLC (60/40) 3.83 minutes (100 %).

Example 83

25 5-Chloro-1H-indole-2-carboxylic acid

[(1S)-(methoxy-methyl-carbamoyl)-ethyl]-amide

(2S)-Amino-N-methoxy-N-methyl-propionamide hydrochloride (169 mg, 1.0 mmol) and 5-chloro-1H-indole-2-carboxylic acid (195 mg, 1.0 mmol) were coupled according to procedure A (washed with 1 N NaOH after acid washes) giving the
30 product (290 mg, 94 %): HPLC (60/40) 4.03 minutes (94 %); PBMS 310/312 (MH⁺, 100 %);

Anal. Calcd for C₁₄H₁₆ClN₃O₃: C, 54.29; H, 5.21; N, 13.57.

Found: C, 54.17; H, 5.26; N, 13.31.

-106-

Example 83a**(2S)-Amino-N-methoxy-N-methyl-propionamide hydrochloride**

5 [(1S)-(Methoxy-methyl-carbamoyl)-ethyl]-carbamic acid tert-butyl ester (3.55 g, 15.3 mmol) was dissolved in 4 M HCl - dioxane (20 ml) at 25 °C for 0.75 hours. The mixture was concentrated and the residue co-evaporated with ether and dichloromethane and dried. Yield 2.2 g (86 %).

Example 83b**[1-(Methoxy-methyl-carbamoyl)-ethyl]-carbamic acid tert-butyl ester**

10 L-BOC-Alanine (3.50 g, 20 mmol) and O,N-dimethyl-hydroxyamine hydrochloride (1.94 g, 20 mmol) were coupled according to procedure A (washed with 1N NaOH after acid washes) and the resulting colorless solid was used without further purification. Yield 3.71 g (80 %).

Example 84**5-Bromo-1H-indole-2-carboxylic acid ((1S)-carbamoyl-2-phenyl-ethyl)-amide**

15 L-Phenylalaninamide hydrochloride (835 mg, 4.17 mmol) and 5-bromo-1H-indole-2-carboxylic acid (1.0 g, 4.17 mmol) were coupled according to procedure A substituting the following workup: the reaction mixture was diluted with ethyl acetate and 2N NaOH. The resulting suspension was filtered and the collected solid washed with ethyl acetate, 2 N NaOH, 2 N HCl, ether, and dried. Yield 890 mg; PBMS
20 386/388 (MH⁺, 100 %);
Anal. Calcd for C₁₈H₁₆BrN₃O₂: C, 55.97; H, 4.18; N, 10.88.
Found: C, 55.69; H, 4.48; N, 10.48.

Example 85**5-Chloro-1H-indole-2-carboxylic acid**

25 **[(1S)-(methoxy-methyl-carbamoyl)-2-phenyl-ethyl]-amide**

(2S)-Amino-N-methoxy-N-methyl-3-phenyl-propionamide hydrochloride (317 mg, 1.3 mmol) and 5-chloro-1H-indole-2-carboxylic acid (253 mg, 1.3 mmol) were coupled according to procedure A (0 - 25 °C, washed first with acid, then base). The crude product was purified by column chromatography on silica gel eluted with 30 % and 40
30 % ethyl acetate in hexanes. The foam obtained was triturated with isopropyl ether yielding an off white solid (356 mg, 71 %): HPLC (60/40) 8.28 minutes (98 %);
Anal. Calcd for C₂₀H₂₀ClN₃O₃: C, 62.26; H, 5.22; N, 10.89.
Found: C, 62.22; H, 5.60; N, 10.73.

-107-

Example 85a

(2S)-Amino-N-methoxy-N-methyl-3-phenyl-propionamide hydrochloride

5 [(1S)-(Methoxy-methyl-carbamoyl)-2-phenyl-ethyl]-carbamic acid tert-butyl ester (2.97 g, 9.6 mmol) was dissolved in 4M HCl-dioxane (36 ml) at 0 °C. The resulting mixture was stirred at 25 °C for 1 hour, concentrated and the residue triturated with ether and dried. Yield 2.27 g, 96 %.

Example 85b

[(1S)-(Methoxy-methyl-carbamoyl)-2-phenyl-ethyl]-carbamic acid tert-butyl ester

10 BOC-L-phenylalanine (4.0 g, 15.1 mmol) and N,O-di-methylhydroxylamine hydrochloride (3.82 g, 15.1 mmol) were coupled according to Procedure A (0 - 25 °C, washed first with acid, then base). The resulting colorless oil was used without purification (3.22 g, 69 %).

Example 86

(2RS)-[(5-Chloro-1H-indole-2-carbonyl)-amino]-15 2-methyl-3-phenyl-propionic acid methyl ester

Racemic 2-amino-2-methyl-3-phenyl-propionic acid methyl ester (200 mg, 0.87 mmol) and 5-chloro-1H-indole-2-carboxylic acid (170 mg, 0.87 mmol) were coupled according to Procedure A (2:1 dichloromethane / dimethylformamide solvent) and the product purified by chromatography on silica gel eluted with 10% ethyl acetate in
20 hexanes. Yield 286 mg, 89 %; HPLC (60/40) 9.63 minutes (85 %); TSPMS 371/373 (MH+, 100 %);

¹H NMR (CDCl₃) δ 9.31 (s, 1H), 7.57 (d, 1H, J = <1 Hz), 7.37 (d, 1H, J = 8.8 Hz), 7.20 (m, 4H), 7.04 (m, 2H), 6.84 (s, 1H), 6.66 (s, 1H), 3.81 (s, 3H), 3.67 (A of AB, 1H, J = 13.5 Hz), 3.28 (B of AB, 1H, J = 13.5 Hz), 1.80 (s, 3H).

25 Example 87

(2RS)-[(5-Chloro-1H-indole-2-carbonyl)-amino]-2-methyl-3-phenyl-propionic acid

Aqueous 2N LiOH (0.10 ml, 0.50 mmol) was added to a solution of (2RS)-[(5-chloro-1H-indole-2-carbonyl)-amino]-2-methyl-3-phenyl-propionic acid methyl ester (132 mg, 0.36 mmol) in tetrahydrofuran (8 ml) at 25 °C. The resulting solution was stirred
30 for 1 hour, concentrated and the residue dissolved in ethyl acetate and water (15 ml). The pH was adjusted to 1 with 2 N HCl at 0 °C. The organic layer was separated, washed with water, brine and dried giving a foam which was used without further

-108-

purification (129 mg, 102 %): HPLC (60/40) 4.42 minutes (99 %); TSPMS 357/359 (MH+, 100 %);

¹H NMR (CDCl₃) δ 9.88 (s, 1H), 7.57 (s, 1H), 7.35 (d, 1H, J = 8.8 Hz), 7.3-7.2 (m, 5H), 7.16 (m, 2H), 6.75 (m, 1H), 6.67 (m, 1H), 3.57 (A of AB, 1H, J = 13.7 Hz), 3.42 (B of
5 AB, 1H, J = 13.7 Hz), 1.80 (s, 3H).

Anal. Calcd for C₁₉H₁₇ClN₂O₃ + 0.3 H₂O: C, 63.00; H, 4.90; N, 7.73.

Found: C, 63.38; H, 5.31; N, 7.42.

Example 88

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-oxo-2-(1-oxo-1-thiomorpholin-4-yl)-ethyl]-amide

10

m-Chloroperoxybenzoic acid (80 mg of 50 %, 0.23 mmol) was added at 25 °C to a solution of 5-chloro-1H-indole-2-carboxylic acid ((1S)-benzyl-2-oxo-2-thiomorpholin-4-yl-ethyl)-amide (100 mg, 0.23 mmol) in dichloromethane (2 mL). After 1 hour, the mixture diluted with ethyl acetate and washed three times with a 50 / 50 mixture of
15 saturated aqueous sodium bicarbonate and 10 % aqueous sodium thiosulfate, once with saturated aqueous sodium bicarbonate, brine, and dried. The crude product was purified by column chromatography on silica gel eluted with 0.5 - 8 % ethanol in dichloromethane to give the title compound. Yield 76 %; HPLC (60/40) 3.97 minutes (97 %); mp 230 - 234 °C; TSPMS 444/446 (MH+, 100 %);

20 Anal. Calcd for C₂₂H₂₂ClN₃O₃S + 0.5 H₂O: C, 58.34; H, 5.12; N, 9.28.

Found: C, 58.41; H, 5.37; N, 8.90.

Example 89

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(1,1-dioxo-1-thiomorpholin-4-yl)-2-oxo-ethyl]-amide

25

m-Chloroperoxybenzoic acid (202 mg of 50 %, 0.58 mmol) was added at 25 °C to a solution of 5-chloro-1H-indole-2-carboxylic acid ((1S)-benzyl-2-oxo-2-thiomorpholin-4-yl-ethyl)-amide (100 mg, 0.23 mmol) in dichloromethane (2 mL). After 1 hour, the mixture was diluted with ethyl acetate and the resulting solution washed three times with a 50 / 50 mixture of saturated aqueous sodium bicarbonate and 10 % aqueous
30 sodium thiosulfate, once with saturated aqueous sodium bicarbonate, brine, and dried. The crude product was purified by column chromatography on silica gel eluted with 30 %, 40 % and 50 % ethyl acetate in hexanes to give the title compound. Yield 60 %; HPLC (60/40) 5.69 minutes (98 %); PBMS 460/462 (MH+, 100 %);

-109-

Anal. Calcd for $C_{22}H_{22}ClN_3O_4S + 0.4 H_2O$: C, 56.56; H, 4.92; N, 8.99.

Found: C, 56.77; H, 5.15; N, 8.60.

Example 90

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-oxo-2-(1-oxo-1-thiazolidin-3-yl)-ethyl]-amide

5 m-Chloroperoxybenzoic acid (167 mg of 50 %, 0.48 mmol) was added at 25 °C to a solution of 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-oxo-2-thiazolidin-3-yl-ethyl]-amide (200 mg, 0.48 mmol) in dichloromethane (4 mL). After 0.5 hours, the mixture was diluted with ethyl acetate and washed three times with a 50 / 50 mixture of saturated aqueous sodium bicarbonate and 10 % aqueous sodium thiosulfate, once with saturated aqueous sodium bicarbonate, brine, and dried. The crude product was concentrated to a yellow solid and then purified by column chromatography on silica gel eluted with 1 - 8 % ethanol in dichloromethane and then triturated with ether giving the title compound. Yield 151 mg (73 %); HPLC (60/40) 3.64 minutes (98 %); PBMS 430/432 (MH⁺, 100 %);

Anal. Calcd for $C_{21}H_{20}ClN_3O_3S + 0.6 H_2O$: C, 57.23; H, 4.85; N, 9.53.

Found: C, 57.00; H, 4.85; N, 9.25.

Example 91

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxyimino-pyrrolidin-1-yl)-2-oxo-ethyl]-amide

20 Hydroxylamine hydrochloride (68 mg, 0.82 mmol) and potassium carbonate (136 mg, 0.98 mmol) were added to a solution of 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-oxo-2-(3-oxo-pyrrolidin-1-yl)-ethyl]-amide in ethanol (5 ml) and water (1 ml) at 25 °C. After 48 hours, the reaction mixture was concentrated and the residue dissolved in ethyl acetate. The resulting solution was washed two times with water and once with brine, dried over Na_2SO_4 , and concentrated. Two substances appearing to be syn/anti oxime isomers separated by chromatography on silica eluted with 2.5 %, 5 %, and 10 % ethanol in dichloromethane.

Example 91(i)

30 For the less polar isomer:

Yield 48 mg (14%); HPLC (60/40) 4.69 minutes (97%); mp 216 - 220 °C (darkened at 210 °C); PBMS 425/427 (MH⁺, 100 %).

-110-

¹H NMR (DMSO-d₆) δ 11.75 (br, 1H), 10.87 (s, 0.5H), 10.86 (s, 0.5H), 9.02 (m, 1H), 7.72 (d, 1H, J = 2.0 Hz), 7.4 - 7.1 (m, 8H), 4.95 (m, 0.5H), 4.85 (m, 0.5H), 4.40 (d, 0.5H, J = 15 Hz), 4.0 (m, 1.5H), 3.9 (m, 0.5H), 3.61 (m, 1H), 3.5 (m, 0.5H), 3.10 (m, 2H), 2.8 - 2.5 (m, 2H);

- 5 Anal. Calcd for C₂₂H₂₁ClN₄O₃: C, 62.19; H, 4.98; N, 13.19.
Found: C, 61.82; H, 5.07; N, 12.95.

Example 91(ii)

For the more polar isomer:

- Yield 69 mg (20 %); HPLC (60/40) 6.78 minutes (>99 %); mp 223 - 224 °C (dec,
10 tar); PBMS 425/427 (MH+, 100 %);

¹H NMR (DMSO-d₆) δ 11.74 (br, 1H), 10.87 (s, 1H), 10.84 (s, 1H), 9.05 (d, 0.5H, J = 8.1 Hz), 8.99 (d, 1H, J = 8.0 Hz), 7.73 (d, 1H, J = 2 Hz), 7.4 - 7.1 (m, 8H), 4.97 (m, 1H), 4.85 (m, 1H), 4.47 (d, 0.5H, J = 17 Hz), 3.95 (m, 1.5H), 3.87 (m, 0.5H), 3.65 - 3.4 (m, 1.5H), 3.10 (m, 2H), 2.7 - 2.5 (m, 2H).

- 15 Anal. Calcd for C₂₂H₂₁ClN₄O₃: C, 62.19; H, 4.98; N, 13.19.
Found: C, 61.85; H, 5.17; N, 13.16.

Example 92

5-Chloro-1H-indole-2-carboxylic acid (1-benzyl-2-oxo-2-piperidin-1-yl-ethyl)-amide

- Piperidine hydrochloride (0.34 mmol) and 2-[(5-chloro-1H-indole-2-carbonyl)-
20 amino]-3-phenyl-propionic acid (0.30 mmol) were coupled according to procedure A (0-25 °C reaction temperature). The crude product was chromatographed on silica gel eluted with 20 %, 30 %, 40 %, 50 %, 75% and 100 % ethyl acetate in hexane giving partial separation. The pure fractions were pooled giving 31 mg (25 %) of the title substance: HPLC (60/40) 9.38 minutes (94 %); PBMS 410/412 (MH+, 100 %);

- 25 Anal. Calcd for C₂₃H₂₄N₃O₂Cl + 0.5 H₂O: C, 65.94; H, 6.02; N, 10.03.
Found: C, 65.70; H, 6.19; N, 9.66.

Example 93

5-Chloro-1H-indole-2-carboxylic acid carbamoylmethyl-amide

- [(5-Chloro-1H-indole-2-carbonyl)-amino]-acetic acid methyl ester (100 mg, 0.40
30 mmol) was added to a saturated solution of ammonia in methanol (ca. 3 mL) at 25 °C. The suspension was sonicated for 1 hour and the resulting solution concentrated. The residue was triturated with ether / hexanes and dried. Yield 77 mg, 77 %; HPLC (60/40) 2.78 minutes (98 %); PBMS 252/254 (MH+, 100 %);

-111-

¹H NMR (DMSO-d₆) δ 11.82 (br, 1H), 8.80 (t, 1H), 7.71 (d, 1H, J = ca.1 Hz), 7.43 (d, 1H, J = 7 - 8 Hz), 7.42 (br, 1H), 7.18 (dd, 1H, J = 7 - 8, ca. 2 Hz), 7.14 (s, 1H), 7.08 (br, 1H), 3.82 (m, 2H).

Anal. Calcd for C₁₁H₁₀ClN₃O₂ + 0.125 H₂O: C, 52.03; H, 4.07; N, 16.55.

5 Found: C, 52.05; H, 4.08; N, 16.63.

Example 94

1-[(2S)-[(5-Bromo-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl]-pyrrolidine-(2S)-carboxylic acid

Trifluoroacetic acid was added to a solution of 1-[(2S)-[(5-bromo-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl]-pyrrolidine-(2S)-carboxylic acid tert-butyl ester (345 mg, 0.64 mmol) in dichloromethane (2 ml) at 0 °C. After 1 hour at 25 °C, the reaction mixture was concentrated, triturated with ether and dried giving a yellow solid. Yield 273 mg, 88 %; HPLC (70/30) 4.75 minutes (98 %); TSPMS 484/486 (MH⁺, 100 %);

15 Anal. Calcd for C₂₃H₂₂BrN₃O₄ + 0.25 H₂O: C, 56.51; H, 4.64; N, 8.60.

Found: C, 56.28; H, 4.78; N, 8.26.

Example 94a

1-[(2S)-[(5-Bromo-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl]-pyrrolidine-(2S)-carboxylic acid tert-butyl ester

20 L-phenylalanine-L-proline tert-butyl ester (333 mg, 1.0 mmol) and 5-bromo-1H-indole-2-carboxylic acid were coupled according to procedure A (72 hour reaction time). The product was purified by column chromatography on silica gel eluted with 15 %, 20 % and 30 % ethyl acetate giving a pale yellow foam. Yield 428 mg (79 %); HPLC (70/30) 5.84 minutes (81 %).

25

Example 95

5-Chloro-1H-indole-2-carboxylic acid [2-oxo-2-(1RS)-oxo-1-thiazolidin-3-yl]-ethyl]-amide

m-Chloroperoxybenzoic acid (426 mg of 50 %, 1.2 mmol) was added at 25 °C to a solution of 5-chloro-1H-indole-2-carboxylic acid [2-oxo-2-thiazolidin-3-yl-ethyl]-amide (400 mg, 1.2 mmol) in dichloromethane (8 mL) at 25 °C. After 1 hour, the mixture was diluted with ethyl acetate (ca 80 mL) and the resulting solution washed three times with a 1:1 mixture of saturated aqueous NaHCO₃ / 10 % aqueous Na₂S₂O₃, saturated aqueous NaHCO₃, and brine. The resulting suspension was filtered and the filtered

-112-

solid washed with water and dried giving a crystalline solid. HPLC (60/40) 2.52 minutes (98.5 %); TSPMS 340/342 (MH⁺, 70 %), 357 (100 %);

¹H NMR (DMSO-d₆) δ 11.82 (br, 1H), 8.84 (m, 1H), 7.73 (d, 1H, J = 2.0 Hz), 7.43 (d, 1H, J = 8.7 Hz), 7.19 (dd, 1H, J = 2.0, 8.7 Hz), 7.18 (s, 1H), 4.92 (dd, 0.5H, J = 12.1 Hz), 4.71 (dd, 0.5H, J = 2.2, 13 Hz), 4.47 (d, 1H, J = 12.1 Hz), 4.4 - 3.9 (m, 4.5H), 3.3 (m, 0.5H), 3.13 (m, 1H), 3.0 (m, 0.5H).

Anal. Calcd for C₁₄H₁₄ClN₃O₃S + 0.8 H₂O: C, 47.47; H, 4.44; N, 11.86.

Found: C, 47.46; H, 4.07; N, 11.83.

Example 96

10 1-[(2S)-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl]-(4R)-hydroxy-pyrrolidine-(2S)-carboxylic acid

Excess aqueous 2 M LiOH was added to a solution of 1-[(2S)-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionyl]-(4R)-hydroxy-pyrrolidine-(2S)-carboxylic acid benzyl ester (215 mg, 0.40 mmol) in tetrahydrofuran at 25 °C. After 2 hours, the mixture was diluted with ethyl acetate and ice and the mixture acidified to pH 1-2 with 6 N HCl. The acidic layer was extracted three times with ethyl acetate, and the organic layers combined and dried. The residue was triturated with ether and dried giving a colorless solid (190 mg, 106 %): HPLC (60/40) 3.43 minutes (94 %); TSPMS 456/458 (MH⁺, 100 %);

20 Anal. Calcd for C₂₃H₂₂ClN₃O₅ + 0.5 C₄H₈O₂: C, 60.06; H, 5.24; N, 8.40.

Found: C, 60.27; H, 5.33; N, 8.13.

Example 97

(S)-2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-(1H-indol-3-yl)-propionic acid methyl ester

25 L-Tryptophan methyl ester hydrochloride (1.05 mmol) and 5-chloro-1H-indole-2-carboxylic acid (1.0 mmol) were coupled according to Procedure A (0 - 25 °C, dimethyl-formamide reaction solvent) and the product purified by chromatography on silica gel eluted with 10 %, 20 %, 30 %, 40 %, 50 % and 60 % ethyl acetate-hexanes giving a yellow foam. Yield, 79 %; HPLC (60/40) 7.43 minutes (96 %);

30 ¹H NMR (CDCl₃) δ 11.78 (br, 1H), 10.85 (br, 1H), 8.93 (d, 1H, J = 7.7 Hz), 7.73 (d, 1H, J = 1.9 Hz), 7.57 (d, 1H, J = 7.7 Hz), 7.41 (d, 1H, J = 8.7 Hz), 7.32 (d, 1H, J = 8.0 Hz), 7.22 (m, 2H), 7.18 (dd, 1H, J = 2.1, 8.8 Hz), 7.06 (m, 1H), 6.99 (m, 1H), 4.74 (m, 1H), 3.65 (s, 3H), 3.35 - 3.2 (m, 2H).

Example 98

(±)-3-{[(5-Chloro-1H-indole-2-carbonyl)-amino]-acetyl}-
thiazolidine-2-carboxylic acid Methyl Ester

(±)-Thiazolidine-2-carboxylic acid methyl ester hydro-chloride (1.02 mmol) and
5 [(5-chloro-1H-indole-2-carbonyl)-amino]-acetic acid (1.02 mmol) were coupled according
to Procedure A (1:1 dichloromethane-dimethylformamide solvent) and the crude product
trituated with 1:1 ether-hexanes giving a light yellow solid. Yield 79 %; HPLC (60/40)
4.47 minutes (95 %); TSPMS 382/384 (MH⁺, 100 %).

¹H NMR (DMSO-d₆) δ 11.82 (s, 1H), 8.85 (t, 1H, J = 7 Hz), 7.73 (d, 1H, J = 2 Hz), 7.43
10 (d, 1H, J = 8.8 Hz), 7.18 (dd, 1H, J = 8.8, 2 Hz), 7.17 (s, 1H), 5.44 (s, 1H), 4.25 (m,
1H), 4.1 (m, 1H), 3.95 (m, 1H), 3.34 (s, 3H), 3.3 (m, 2H).

Anal. Calcd for C₁₆H₁₆ClN₃O₄S: C, 50.33, H 4.22; N, 11.00.

Found: C, 50.56; H, 4.46; N, 10.89.

Example 99

15 (±)-3-{[(5-Chloro-1H-indole-2-carbonyl)-amino]-acetyl}-thiazolidine-
2-carboxylic acid

A solution of 3-{[(5-chloro-1H-indole-2-carbonyl)-amino]-acetyl}-thiazolidine-2-
carboxylic acid methyl ester (196 mg, 0.5 mmol) in methanol (10 mL) was treated with
aqueous 1 N NaOH (0.5 mL) at 25 °C. After 3 hours, more 1 N NaOH (0.25 mL) was
20 added. The mixture was stirred at 25 °C overnight, concentrated, the residue stirred
with ethyl acetate (30 mL) and 1 N NaOH (5 mL), and the resulting mixture acidified to
pH 1.8 with aqueous 6 N HCl. The aqueous layer was separated and extracted with
ethyl acetate. The organic layers were combined, dried, and concentrated giving a
solid which was trituated with 1:1 ether - hexane and dried. Yield 186 mg, 99 %;
25 HPLC (60/40) 3.13 minutes (98 %); TSPMS 368/370 (MH⁺, 70 %), 339 (100 %).

¹H NMR (DMSO-d₆) δ 11.80 (s, 1H), 8.84 (br, 1H), 7.23 (s, 1H), 7.44 (d, 1H, J = 8.8
Hz), 7.18 (dd, 1H), 7.17 (s, 1H), 4.32 (s, 1H), 4.25 (m, 2H), 4.0 (m, 2H), 3.3 (m, 2H).

Example 99a

(±)-Thiazolidine-2-carboxylic Acid Methyl Ester

30 A mixture of (±)-thiazolidine-2-carboxylic acid (1.58 g, 11.9 mmol) and
chlorotrimethylsilane (5.1 g, 47 mmol) in methanol (22 mL) was heated at reflux for 5
hours, cooled, and concentrated giving a solid (2.19 g, 100 %).

Example 100**S-tert-Butyl 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionate.****Procedure B**

To a solution of 5-chloro-1H-indole-2-carboxylic acid (0.50 g, 2.6 mmol), L-phenylalanine tert-butyl ester hydro-chloride (0.66 g, 2.6 mmol), triethylamine (0.36 mL, 2.6 mmol) and 4-dimethylaminopyridine (0.16 g, 1.3 mmol) in dichloromethane (20 mL) was added 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide (0.73 g, 3.8 mmol). The mixture was stirred at room temperature overnight, diluted with chloroform, washed with 2N HCl, water and brine, dried over magnesium sulfate and concentrated. The product was purified by flash-chromatography (30 % acetone in hexanes) and obtained as a pale yellow foam (0.86 g, 85 %).

Anal. calc.: C 66.25, H 5.81, N 7.03;

Found: C 66.57, H 6.11, N 6.86.

The following examples (101 to 122) were prepared by methods analogous to Procedure B.

Example 101**R-Methyl-2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionate**

From 5-fluoro-1H-indole-2-carboxylic acid and D-phenyl-alanine methyl ester.

¹H NMR (300 MHz, CDCl₃) δ 3.22 (m, 2 H), 3.80 (s, 3 H), 5.10 (m, 1H), 6.62 (d, 6 Hz, 1H), 6.75 (d, 2 Hz, 1 H), 7.05 (dt, 2 Hz, 8 Hz, 1H), 7.10-7.15 (m, 2H), 7.25-7.40 (m, 4H), 7.73 (d, 2.1 Hz, 1H), 9.50 (br, 1H).

Example 102**R-Methyl 2-[(5-7-dichloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionate**

From 5,7-dichloro-1H-indole-2-carboxylic acid and D-phenylalanine methyl ester.

¹H NMR (300 MHz, CDCl₃) δ 3.25 (m, 2H), 3.80/3.95 (s, 3H), 5.10 (m, 1H), 6.62 (d, 6 Hz, 1H), 6.69 (d, 2 Hz, 1H), 7.10 - 7.15 (m, 2H), 7.25 - 7.35 (m, 3H), 7.50 - 7.56 (s, 1H), 9.35 (br, 1H).

Example 102a**5,7-Dichloro-1H-indole-2-carboxylic acid****30 A. Ethyl 2-oxopropionate 2,4-dichlorophenylhydrazone**

A mixture of 2,4-dichlorophenylhydrazine (1.0 g, 4.7 mmol), ethyl pyruvate (0.53 mL, 4.7 mmol), triethylamine (0.65 mL, 4.7 mmol) and ethanol (5 mL) was heated at reflux overnight. The solvent was evaporated and the residue taken up in chloroform.

-115-

The solution was washed with water and brine and dried over magnesium sulfate and concentrated, leaving an oil (1.1 g, 98 %).

B. Ethyl 5,7-dichloro-1H-indole-2-carboxylate

A solution of ethyl 2-oxopropionate 2,4-dichloro-phenylhydrazone (1.1 g, 4.6 mmol) and anhydrous zinc chloride (10 g, 74 mmol) in glacial acetic acid (12 mL) was heated at reflux for 1/2 hours. The reaction mixture was poured into water and extracted with ether twice. The combined organic layers were washed with water and brine, dried over magnesium sulfate and concentrated. The product was purified by flash-chromatography (30 % ethyl acetate in hexanes) and obtained as an oil (0.80 g, 67 %).

C. 5,7-Dichloro-1H-indole-2-carboxylic acid

A solution of ethyl 5,7-dichloro-1H-indole-2-carboxylate (0.80 g, 3.1 mmol) in 1 N NaOH (40 mL) and methanol (50 mL) was heated to reflux for 3 hours. The methanol was removed in vacuo and the aqueous residue was acidified with 1 N HCl and extracted with chloroform twice. The combined extracts were washed with water and brine, dried over magnesium sulfate and concentrated to a solid (0.58 g, 76 %).

The following indole carboxylic acids were prepared by the same sequence:

4-Chloro-5-fluoro-1H-indole-2-carboxylic acid and 6-chloro-5-fluoro-1H-indole-2-carboxylic acid (as a mixture) from 3-chloro-4-fluorophenylhydrazine.

20 5,7-Difluoro-1H-indole-2-carboxylic acid from 2,4-difluorophenylhydrazine

Example 103

(±)-Ethyl-2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionate

From 5-chloro-1H-indole-2-carboxylic acid and DLphenyl-alanine ethyl ester.

mp 146 - 147 °C.

25 Anal. Calc.: C 64.61, H 5.42, N 7.54;

Found: C 64.73, H 5.26, N 7.57.

Example 104

S-3-Bromo-5-chloro-1H-indole-2-carboxylic acid (1-dimethyl-carbamoyl-2-phenyl-ethyl)-amide

30 From 3-bromo-5-chloro-1H-indole-2-carboxylic acid and S-2-amino-N,N-dimethyl-3-phenyl-propionamide.

Anal. Calc.: C 53.53, H 4.27, N 9.36;

Found: C 53.51, H 4.46, N 9.38.

-116-

Example 104a

3-Bromo-5-chloro-1H-indole-2-carboxylic acid

To a solution of 5-chloro-1H-indole-2-carboxylic acid (2.0 g, 10.2 mmol) in acetic acid (24 mL) was added a solution of bromine (0.53 mL, 10.2 mmol) in acetic acid (16 mL). After 20 minutes, the mixture was poured into water and extracted with chloroform twice. The combined extracts were washed with water twice and brine, dried over magnesium sulfate and concentrated. The product was obtained as a solid (2.5 g, 89 %).

Example 104b

10 (S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochlorideA. (S)-(1-Dimethylcarbamoyl-2-phenyl-ethyl)-carbamic acid tert-butyl ester

To a solution of tert-Boc-phenylalanine (10 g, 38 mmol), dimethylamine hydrochloride (3.4 g, 41 mmol), triethylamine (5.8 mL, 42 mmol) and hydroxybenzo-triazole (6.6 g, 49 mmol) in dichloromethane (300 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (9.4 g, 49 mmol). The mixture was stirred overnight, then quenched with 2 N HCl and concentrated. The residue was taken up in ethyl acetate and this solution was washed with water and brine, dried over magnesium sulfate and concentrated. The residue was triturated in chloroform, the solid was filtered and the filtrate was concentrated to an oil (11 g, 100 %).

20 B. (S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride

(S)-(1-Dimethylcarbamoyl-2-phenyl-ethyl)-carbamic acid tert-butyl ester (11.0 g, 38 mmol) was dissolved in ethyl acetate (125 mL) and HCl was bubbled into the solution for 10 min. The solution was stirred for 1 hour at room temperature, then concentrated. The residue was triturated in ether, the solid was filtered and dried on high vacuum (8.6 g, 100 %).

Example 105

S-5-Chloro-4-nitro-1H-indole-2-carboxylic acid (1-dimethylcarbamoyl-2-phenyl-ethyl)-amide

From 4-nitro-5-chloro-1H-indole-2-carboxylic acid and S-2-amino-N,N-dimethyl-3-phenyl-propionamide.

¹H NMR (300 MHz, CDCl₃) δ 2.75 (s, 3H), 2.97 (s, 3H), 3.20 (m, 2H), 5.30 (m, 1H), 7.07 (d, 2 Hz, 1H), 7.24 - 7.32 (m, 5H), 7.40 (d, 7 Hz, 1H), 8.12 (br d, 7 Hz, 1H), 9.85 (br, 1H).

Example 105a

4-Nitro-5-chloro-1H-indole-2-carboxylic acidA. 2-[(4-Chloro-3-nitro-phenyl)-hydrazono]-propionic acid ethyl ester

To a solution of sodium nitrite (2.17 g, 31 mmol) in water (60 mL) and conc. HCl (12 mL) at 0 °C was added 4-chloro-3-nitroaniline (5.0 g, 29 mmol). After 5 minutes, a solution of ethyl methylacetoacetate (4.5 mL, 29 mmol) in water (60 mL), ethanol (30 mL) and 50 % potassium hydroxide (10 mL) was added and the reaction mixture was stirred overnight. The precipitate was collected (7.0 g, 91 %).

B. Ethyl 5-chloro-4-nitro-1H-indole-2-carboxylate

A mixture of 2-[(4-chloro-3-nitro-phenyl)-hydrazono]-propionic acid ethyl ester (2.0 g, 6.7 mmol) and polyphosphoric acid (7 g) was heated to 90 - 110 °C for 2 hours. The mixture was cooled, poured onto an ice/water mixture and the solid was collected. Flash-chromatography (1 % methanol in chloroform) provided the title compound (0.58 g, 32 %) and 5-chloro-6-nitro-1H-indole-2-carboxylate (0.31 g, 17 %).

C. 4-Nitro-5-chloro-1H-indole-2-carboxylic acid

The title compound was prepared by hydrolysis of ethyl 5-chloro-4-nitro-1H-indole-2-carboxylate as described for the preparation of 5,7-dichloro-1H-indole-2-carboxylic acid.

Example 106

20 S-7-Nitro-1H-indole-2-carboxylic acid (1-dimethylcarbamoyl-2-phenyl-ethyl)-amide

From 7-nitro-1H-indole-2-carboxylic acid and S-2-amino-N,N-dimethyl-3-phenyl-propionamide.

¹H NMR δ 2.8 (s, 3H), 3.0 (s, 3H), 3.1-3.3 (m, 2H), 5.35 (q, 7 Hz, 1H), 6.95 (s, 1H), 7.15 - 7.3 (m, 6H), 7.9 (d, 8 Hz, 1H), 8.2 (d, 8 Hz, 1H), 10.3 (br, 1H).

25

Example 107

(±)-Methyl 2-[(5-chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-butyrate

From 5-chloro-1H-indole-2-carboxylic acid and DL-β-methylphenylalanine methyl ester.

mp 135 - 136 °C.

30 Anal. Calc.: C 64.78, H 5.17, N 7.56;

Found: C 64.76, H 5.26, N 7.64.

Example 108

(±)-5-Chloro-1H-indole-2-carboxylic acid [1-(2-fluoro-benzyl)-
2-oxo-2-thiazolidin-3-yl-ethyl]-amide

From 5-chloro-1H-indole-2-carboxylic acid and (±)-2-amino-3-(2-fluoro-
5 phenyl)-1-thiazolidin-3-yl-propan-1-one.

mp 216 - 217 °C.

Anal. Calc.: C 58.40, H 4.43, N 9.73;

Found: C 58.45, H 4.53, N 9.71.

Example 108a

10 (±)-2-Amino-3-(2-fluorophenyl)-1-thiazolidin-3-yl-propan-1-one hydrochloride

A. (±)-2-tert-Butoxycarbonylamino-3-(2-fluoro-phenyl)-propionic acid

To a mixture of DL-3-fluoro-phenylalanine (1.0 g, 5.5 mmol) and triethylamine (1.14 mL, 8.2 mmol) in dichloro-methane (20 mL) was added di-tert-butyl dicarbonate (1.4 g, 6.55 mmol). The mixture was stirred at room temperature overnight, then
15 poured into water, acidified with 1 N HCl and extracted with chloroform. The combined extracts were washed with water and brine, dried over magnesium sulfate and concentrated. The product was purified by flash-chromatography (chloroform / methanol / acetic acid, 89:10:1) and obtained as a solid (1.28 g, 83 %, mp 118 - 119 °C).

20 B. (±)-[1-(2-Fluorobenzyl)-2-oxo-2-thiazolidin-3-yl-ethyl]-carbamic acid tert-butyl ester

To a mixture of 2-tert-butoxycarbonylamino-3-(2-fluoro-phenyl)-propionic acid (0.50 g, 1.77 mmol), thiazolidine (0.15 mL, 1.94 mmol) and 4-dimethylamino-pyridine (0.21 g, 1.77 mmol) in dichloromethane (15 mL) was added EDC (0.44 g, 2.31 mmol). The reaction mixture was stirred at room temperature overnight, diluted with chloroform,
25 washed with 2 N HCl, water and brine, dried over magnesium sulfate and concentrated. The product was purified by flash-chromatography (30 % acetone in hexanes) and obtained as a colorless solid (0.39 g, 62 %, mp 133 - 134 °C).

C. (±)-2-Amino-3-(2-fluorophenyl)-1-thiazolidin-3-yl-propan-1-one hydrochloride

HCl was bubbled into a solution of [1-(2-fluoro-benzyl)-2-oxo-2-thiazolidin-3-yl-ethyl]-carbamic acid tert-butyl ester (0.39 g, 1.1 mmol) in ethyl acetate (15 mL). The
30 solution was concentrated, the residue was triturated in ether, the solid was filtered and dried (0.27 g, 84 %, mp 217 - 218 °C).

-119-

The following amines were prepared by analogous methods in the same sequence:

- (±)-2-Amino-3-(2-chloro-phenyl)-1-thiazolidin-3-yl-propan-1-one from DL-2-chloro-phenylalanine
- 5 (±)-2-Amino-3-(3-cyano-phenyl)-1-thiazolidin-3-yl-propan-1-one from DL-3-cyano-phenylalanine
- (±)-2-Amino-3-(3-chloro-phenyl)-1-thiazolidin-3-yl-propan-1-one from DL-3-chloro-phenylalanine
- (±)-2-Amino-3-(3-trifluoromethyl-phenyl)-1-thiazolidin-3-yl-propan-1-one from DL-3-
- 10 trifluoromethyl-phenylalanine
- (S)-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-(4-methoxy-phenyl)-propan-1-one from L-4-methoxy-phenylalanine

Example 109

(±)-5-Chloro-1H-indole-2-carboxylic acid [1-(2-chloro-benzyl)-

15 2-oxo-2-thiazolidin-3-yl-ethyl]-amide

From 5-chloro-1H-indole-2-carboxylic acid and (±)-2-amino-3-(2-chloro-phenyl)-1-thiazolidin-3-yl-propan-1-one.

mp 214 - 216 °C.

Anal. Calc.: C 56.26, H 4.58, N 9.37;

20 Found: C 56.27, H 4.54, N 9.36.

Example 110

(±)-5-Chloro-1H-indole-2-carboxylic acid [2-(3-cyano-phenyl)-

1-(thiazolidine-3-carbonyl)-ethyl]-amide

From 5-chloro-1H-indole-2-carboxylic acid and (±)-2-amino-3-(3-cyano-

25 phenyl)-1-thiazolidin-3-yl-propan-1-one.

mp 183 - 184 °C.

Anal. Calc.: C 60.20, H 4.36, N 12.77;

Found: C 60.11, H 4.84, N 12.43.

Example 111

30 (±)-5-Chloro-1H-indole-2-carboxylic acid [1-(3-chloro-benzyl)-

2-oxo-2-thiazolidin-3-yl-ethyl]-amide

From 5-chloro-1H-indole-2-carboxylic acid and (±)-2-amino-3-(3-chloro-phenyl)-1-thiazolidin-3-yl-propan-1-one.

-120-

mp 188 - 190 °C.

Anal. Calc.: C 56.26, H 4.27, N 9.37;

Found: C 56.38, H 5.04, N 9.04.

Example 112

5 (±)-5-Chloro-1H-indole-2-carboxylic acid [2-oxo-2-thiazolidin-3-yl-1-(3-trifluoromethyl-benzyl)-ethyl]-amide

From 5-chloro-1H-indole-2-carboxylic acid and (±)-2-amino-3-(3-trifluoromethyl-phenyl)-1-thiazolidin-3-yl-propan-1-one.

mp 205 - 207 °C.

10 Anal. Calc.: C 54.83, H 3.97, N 8.72;

Found: C 54.44, H 4.14, N 8.88.

Example 113

S-5-Chloro-1H-indole-2-carboxylic acid [1-(4-methoxy-benzyl)-2-oxo-2-thiazolidin-3-yl-ethyl]-amide

15 From 5-chloro-1H-indole-2-carboxylic acid and S-2-amino-3-(4-methoxy-phenyl)-1-thiazolidin-3-yl-propan-1-one.

Anal. Calc.: C 59.52, H 5.00, N 9.47;

Found: C 60.00, H 5.55, N 8.90.

Mass Spec. m/e 444 (M⁺+1).

20

Example 114

(±)-5-Chloro-1H-indole-2-carboxylic acid [1-(3-chloro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

From 5-chloro-1H-indole-2-carboxylic acid and (±)-2-amino-1-(4-hydroxy-piperidin-1-yl)-3-(3-chloro-phenyl)-propan-1-one.

25 mp 98 °C dec.

Example 115

S-5-Chloro-4-fluoro-1H-indole-2-carboxylic acid (1-benzyl-2-oxo-2-thiazolidin-3-yl-ethyl)-amide and S-6-chloro-4-fluoro-1H-indole-2-carboxylic acid (1-benzyl-2-oxo-2-thiazolidin-3-yl-ethyl)-amide

30 From a mixture of 5-chloro-4-fluoro-1H-indole-2-carboxylic acid and 6-chloro-4-fluoro-1H-indole-2-carboxylic acid, and S-2-amino-3-phenyl-1-thiazolidin-3-yl-propan-1-one.

mp 105 - 125 °C dec.

-121-

Anal. Calcd.: C, 58.40; H, 4.43; N, 9.73;

Found: C, 58.54; H, 4.59; N, 9.58.

Example 116

(+)-2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-propionic acid methyl ester

5 From 5-chloro-1H-indole-2-carboxylic acid and DL-alanine methyl ester hydrochloride.

mp 199 - 201 °C.

Anal. Calc.: C 55.63, H 4.67, N 9.98;

Found: C 55.70, H 4.75, N 10.06.

10

Example 117

(±)-2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-[4-(4,5-dihydro-1H-imidazol-2-yl)-phenyl]-propionic acid methyl ester

From 5-chloro-1H-indole-2-carboxylic acid and (±)-2-amino-3-[4-(4,5-dihydro-1H-imidazol-2-yl)-phenyl]-propionic acid methyl ester.

15 ¹H NMR (300 MHz, CDCl₃) δ 3.1 - 3.3 (m, 2H), 3.70 (s, 3H), 3.95 (s, 4H), 4.85 (m, 1H), 7.15 (s, 1H), 7.17 (d, 8 Hz, 1H), 7.40 (d, 8 Hz, 1H), 7.65 (d, 7 Hz, 1H), 7.75 (s, 1H), 7.88 (d, 8 Hz, 1H), 9.10 (br d, 9 Hz, 1H), 10.5 (s, 1H), 11.8 (br s, 1H).

Example 117a

(±)-2-Amino-2-[4-(4,5-dihydro-1H-imidazol-2-yl)-phenyl]-propionic acid methyl ester

20 A. 2-Acetylamino-2-[4-(4,5-dihydro-1H-imidazol-2-yl)-benzyl]-malonic acid diethyl ester

A solution of 2-acetylamino-2-[(4-methoxycarbon-imidoyl)-benzyl]-malonic acid, diethyl ester (G. Wagner et al. *Pharmazie* 1974, 29, 12) (5.3 g, 13 mmol) and ethylenediamine (4.8 g, 80 mmol) in ethanol (100 mL) was stirred at 60 °C for 5 hours. After cooling, the solvent was evaporated, water was added to the residue and the solid
25 was filtered and dissolved in hot 1 N HCl. After cooling, the precipitate was filtered and dried (3.1 g).

B. (±)-2-Amino-3-[4-(4,5-dihydro-1H-imidazol-2-yl)-phenyl]-propionic acid dihydrochloride

To 2-acetylamino-2-[4-(4,5-dihydro-1H-imidazol-2-yl)-benzyl]-malonic acid, diethyl
30 ester (3.0 g, 7.3 mmol) was added glacial acetic acid (50 mL) and 3N HCl (100 mL). The solution was heated to reflux for 3 hours, cooled and concentrated to a white solid which was recrystallized from methanol/ether (2.0 g, mp 270 - 272 °C dec.).

C. (±)-2-Amino-3-[4-(4,5-dihydro-1H-imidazol-2-yl)-phenyl]-propionic acid methyl ester

-122-

(±)-2-Amino-3-[4-(4,5-dihydro-1H-imidazol-2-yl)-phenyl]-propionic acid dihydrochloride (0.50 g, 1.6 mmol) was placed in thionyl chloride (1 mL) and methanol (25 mL). The mixture was heated to reflux for 30 minutes, at which time more thionyl chloride (3 mL) and methanol (75 mL) were added. After another 3 hours at reflux, the solution was concentrated, the residue was dissolved in a small amount of methanol and ethyl acetate was added to induce precipitation. The solid was collected and dried (0.40 g, mp 230 °C dec.).

Example 118

10 (S)-5,7-Difluoro-1H-indole-2-carboxylic acid [1-benzyl-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

From 5,7-difluoro-1H-indole-2-carboxylic acid and (S)-2-amino-1-(4-hydroxy-piperidin-1-yl)-3-phenyl-propan-1-one.

mp 95 - 110 °C.

Example 119

15 S-4-chloro-5-fluoro-1H-indole-2-carboxylic acid (1-dimethylcarbamoyl-2-phenyl-ethyl)-amide and S-6-chloro-5-fluoro-1H-indole-2-carboxylic acid (1-dimethylcarbamoyl-2-phenyl-ethyl)-amide

From a mixture of 5-chloro-4-fluoro-1H-indole-2-carboxylic acid and 6-chloro-4-fluoro-1H-indole-2-carboxylic acid, and (S)-2-amino-N,N-dimethyl-3-phenyl-propionamide.

mp 200 - 210 °C.

Anal. Calc.: C 61.94, H 4.94, N 10.83;

Found: C 62.21, H 4.99, N 10.84.

Example 120

25 (S)-5,7-Difluoro-1H-indole-2-carboxylic acid (1-benzyl-2-oxo-2-thiazolidin-3-yl-ethyl)-amide

From 5,7-difluoro-1H-indole-2-carboxylic acid and (S)-2-amino-3-phenyl-1-thiazolidin-3-yl-propan-1-one.

mp 175 - 185 °C.

Anal. Calc.: C 60.71, H 4.61, N 10.11;

30 Found: C 60.79, H 4.66, N 9.93.

-123-

Example 121

(S)-5,7-Difluoro-1H-indole-2-carboxylic acid [1-benzyl-2-(1,1-dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-amide

From 5,7-difluoro-1H-indole-2-carboxylic acid and (S)-2-amino-1-(1,1-dioxo-1-thiazolidin-3-yl)-3-phenyl-propan-1-one hydrochloride.

mp 95 - 110 °C.

MS 448 (MH⁺).

Example 122

S-5-Chloro-1H-indole-2-carboxylic acid [1-(2-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

Procedure C

To a solution of 5-chloro-1H-indole-2-carboxylic acid (0.49 g, 2.5 mmol), S-2-amino-3-(2-fluoro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one (0.76 g, 2.5 mmol), triethylamine (0.35 mL, 2.5 mmol) and hydroxybenzotriazole (0.34 g, 2.5 mmol) in dichloromethane (6 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.53 g, 2.8 mmol). The mixture was stirred at room temperature overnight, diluted with dichloromethane, washed with water, 1 N HCl and saturated sodium bicarbonate, dried over magnesium sulfate and concentrated. The product was purified by flash-chromatography (chloroform / methanol, 8:1) and obtained as an off-white solid (0.82 g, 73 %).

mp 120 - 122 °C.

Example 122a

(S)-2-Amino-3-(2-fluoro-phenyl)-1-

(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride

A. (S)-2-tert-Butoxycarbonylamino-3-(2-fluorophenyl)-1-thiazolidin-3-yl-propan-1-one

From L-Boc-2-fluorophenylalanine and 4-hydroxypiperidine by a method analogous to Procedure C.

B. (S)-2-Amino-3-(2-fluoro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one hydrochloride

The title compound was prepared by reaction of L-2-tert-butoxycarbonylamino-3-(2-fluorophenyl)-1-thiazolidin-3-yl-propan-1-one with HCl according to the analogous method described in Example 108a, step C.

-124-

The following amines were prepared by analogous methods in the same sequence:

S-2-Amino-3-(4-methoxy-phenyl)-1-thiazolidin-3-yl-propan-1-one

S-2-Amino-3-(2-fluoro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one

5 S-2-Amino-1-(4-hydroxy-piperidin-1-yl)-3-(4-methoxy-phenyl)-propan-1-one

S-2-Amino-3-(2-chloro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one

S-2-Amino-3-(4-methoxy-phenyl)-1-morpholin-4-yl-propan-1-one

S-2-amino-3-(4-methoxy-phenyl)-1-(4-acetyl-piperazinyl)-propan-1-one

By an analogous process to that of (Procedure C) were prepared the following
10 examples (122-138).

Example 123

(2SR), (3RS)-2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-hydroxy-3-phenyl-propionic acid methyl ester

From 5-chloro-1H-indole-2-carboxylic acid and (+)-threo b-phenylserine methyl
15 ester.
mp 196 - 197 °C.

Example 124

S-5-Fluoro-1H-indole-2-carboxylic acid [1-(4-methoxy-benzyl)-2-oxo-2-thiazolidin-3-yl-ethyl]-amide

20 From 5-fluoro-1H-indole-2-carboxylic acid and S-2-amino-3-(4-methoxy-phenyl)-1-thiazolidin-3-yl-propan-1-one.
mp 90 - 115 °C.

Anal. calc.: C 61.81, H 5.19, N 9.83;

Found: C 60.94, H 5.33, N 10.01.

25

Example 125

S-5-Chloro-1H-indole-2-carboxylic acid [1-(2-chloro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide

From 5-chloro-1H-indole-2-carboxylic acid and S-2-amino-3-(2-chloro-phenyl)-1-(4-hydroxy-piperidin-1-yl)-propan-1-one.

30 mp 127 - 129 °C

-125-

Example 126

S-5-Chloro-1H-indole-2-carboxylic acid [1-(4-methoxy-benzyl)-2-morpholin-4-yl-2-oxo-ethyl]-amide

From 5-chloro-1H-indole-2-carboxylic acid S-2-amino-3-(4-methoxy-phenyl)-1-morpholin-4-yl-propan-1-one.

mp 95 - 105 °C.

Anal. calc.: C 62.51, H 5.47, N 9.51;

Found: C 61.82, H 6.05, N 8.97.

Example 127

10 S-5-Chloro-1H-indole-2-carboxylic acid [2-(4-acetyl-piperazin-1-yl)-1-(4-methoxy-benzyl)-2-oxo-ethyl]-amide

From 5-chloro-1H-indole-2-carboxylic acid and S-2-amino-3-(4-methoxy-phenyl)-1-(4-acetyl-piperazinyl)-propan-1-one.

mp 120 - 135 °C.

15 Anal. calc.: C 62.17, H 5.64, N 11.60;

Found: C 62.76, H 6.20, N 10.44.

Example 128

S-5-Fluoro-1H-indole-2-carboxylic acid [1-(benzothiazol-2-ylcarbamoyl)-2-phenyl-ethyl]-amide

20 From S-2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and 2-amino-1,3-benzothiazole.

mp 139 - 141 °C.

Example 129

S-5-Fluoro-1H-indole-2-carboxylic acid (1-benzyl-2-morpholin-4-yl-2-oxo-ethyl)-amide

25 From S-2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and morpholine.

mp 234 - 236 °C.

Example 130

5-Fluoro-1H-indole-2-carboxylic acid [1S-benzyl-2-oxo-2-(3,3,5-trimethyl-azepan-1-yl)-ethyl]-amide

30 From 2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and (+)-3,3,5-trimethylazepane.

mp 125 - 127 °C.

-126-

Anal. calc.: C 72.14, H 7.18, N 9.35;

Found: C 72.00, H 7.58, N 9.10.

Example 131

5 5-Fluoro-1H-indole-2-carboxylic acid [1S-benzyl-2-(3RS-carbamoyl-piperidin-1-yl)-2-oxo-ethyl]-amide

From S-2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and 3-carbamoyl-piperidine.

mp 234 - 236 °C.

Example 132

10 5-Fluoro-1H-indole-2-carboxylic acid [2-phenyl-1S-(thiochroman-4RS-ylcarbamoyl)-ethyl]-amide

From S-2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and (+)thiochroman-4-ylamine.

mp 225 - 226 °C.

15 Anal. calc.: C 68.48, H 5.11, N 8.88;

Found: C 68.40, H 5.64, N 8.61.

Example 133

20 S-5-Fluoro-1H-indole-2-carboxylic acid [1-(5-methyl-isoxazol-3-ylcarbamoyl)-2-phenyl-ethyl]-amide

From S-2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and 5-methyl-isoxazol-3-ylamine.

mp 219 - 221 °C.

Example 134

25 S-5-Fluoro-1H-indole-2-carboxylic acid [2-phenyl-1-(4,5,6,7-tetrahydro-benzothiazol-2-ylcarbamoyl)-ethyl]-amide

From S-2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and 4,5,6,7-tetrahydro-benzothiazol-2-yl-amine.

mp 162 - 165 °C.

Example 135

30 S-5-Fluoro-1H-indole-2-carboxylic acid [1-(5-methyl-thiazol-2-ylcarbamoyl)-2-phenyl-ethyl]-amide

From S-2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and 4-methyl-thiazol-2-ylamine.

-127-

mp 211 - 213 °C.

Example 136

S-5-Methyl-1H-indole-2-carboxylic acid [1-(5-methyl-isoxazol-3-ylcarbamoyl)-2-phenyl-ethyl]-amide

5 From S-2-[(5-methyl-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and 5-methyl-isoxazol-3-ylamine.

mp 243 - 245 °C.

Anal. calc.: C 68.64, H 5.51, N 13.93;

Found: C 68.29, H 5.81, N 14.05.

10

Example 137

S-5-Methyl-1H-indole-2-carboxylic acid [2-(4-acetyl-piperazin-1-yl)-1-benzyl-2-oxo-ethyl]-amide

From S-2-[(5-methyl-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and 1-piperazin-1-yl-ethanone.

15 mp 221 - 223 °C.

Example 138

S-5-Chloro-1H-indole-2-carboxylic acid (1-carbamoyl-2-phenyl-ethyl)-amide

From 5-chloro-1H-indolecarboxylic acid and S-2-amino-3-phenyl-propionamide.

mp 257 - 258 °C.

20

Examples 139 and 140

(2RS)-2,3-Dihydro-1H-indole-2-carboxylic acid(R-1-dimethylcarbamoyl-2-phenyl-ethyl)-amide

To a mixture of DL-indoline-2-carboxylic acid (0.38 g, 2.3 mmol), (R)-2-amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride (0.53 g, 2.3 mmol), hydroxybenzotriazole (0.66 g, 4.2 mmol) and triethylamine (0.32 mL, 2.3 mmol) in dichloromethane (5 mL) was added EDC (0.64 g, 2.7 mmol). The solution was stirred overnight, diluted with dichloro-methane, washed with water and brine, dried over magnesium sulfate and concentrated. The two isomeric products were separated by flash-chromatography (EtOAc, then EtOAc / MeOH, 20:1).

30

Example 139

Less polar isomer (oil, 0.23 g, 30%):

-128-

¹H NMR (300 MHz, CDCl₃) δ 2.68 (s, 3H), 2.87 (s, 3H), 3.02-3.09 (m, 3H), 3.55 (dd, J = 10 Hz, 6 Hz, 1H), 4.61 (m, 1H), 5.10 (q, J = 8 Hz, 1H), 6.95 (d, J = 8 Hz, 1H), 7.11-7.30 (m, 8H), 8.12 (br, 1H).

MS (Cl, NH₃) 394 (M⁺ + 17).

5

Example 140

More polar isomer (0.11 g, 14 %): mp 136 - 140 °C.

Examples 141 and 142**(2RS)-5-Chloro-2,3-dihydro-1H-indole-2-carboxylic acid****(1-S-dimethylcarbamoyl-2-phenyl-ethyl)-amide.**

- 10 To a solution of S-5-chloro-1H-indole-2-carboxylic acid (1-dimethylcarbamoyl-2-phenyl-ethyl)-amide (2.60 g, 7.0 mmol) in THF (20 mL) and methanol (20 mL) was added magnesium (1.75 g, 73 mmol) by portions, at such a rate as to maintain the reaction going without excess heat. After the reaction had ceased, the reaction was concentrated to a low volume, the residue was partitioned between 1 N HCl and ethyl
- 15 acetate, the combined ethyl acetate layers were washed with water and brine, dried over magnesium sulfate and concentrated. The products were separated by flash-chromatography (1 % methanol in chloroform).

Example 141

Less polar isomer:

- 20 ¹H NMR (300 MHz, CDCl₃) δ 2.61 (s, 3H), 2.83 (s, 3H), 3.00 - 3.02 (m, 3H), 3.47 (dd, J = 9.9 Hz, 6.4 Hz, 1H), 4.43 (m, 1H), 5.10 (q, J = 7.5 Hz, 1H), 6.64 (d, J = 8.8 Hz, 1H), 7.00 (s, 1H) 7.16-7.29 (m, 7H), 7.70 (br, 1H).
- MS (Cl, NH₃) 372 (M⁺ + 1).

Example 142

- 25 More polar isomer: mp 125 °C dec.

Examples 143 and 144**2RS 5-chloro-2,3-dihydro-1H-indole-2-carboxylic acid (1R-dimethylcarbamoyl-2-phenyl-ethyl)-amide**

- By an analogous method to that of Example 141 and 142, using R-5-chloro-1H-
- 30 indole-2-carboxylic acid (1-dimethylcarbamoyl-2-phenyl-ethyl)-amide, the two diastereomers were prepared.

Example 143

Less polar isomer: mp 122 - 124 °C dec.

Example 144

More polar isomer:

¹H NMR (300 MHz, CDCl₃) δ 2.70 (s, 3H), 2.83 (s, 3H), 2.77-2.97 (m, 3H), 3.40 (dd, J = 16.6 Hz, 10.8 Hz, 1H), 4.28 (m, 1H), 4.40 (d, J = 5.2 Hz, 1H), 5.12 (q, J = 7.8 Hz, 1H), 6.56 (d, J = 9.0 Hz, 1H), 6.96 (d, J = 7.8 Hz, 1H), 6.99 (s, 1H) 7.03 - 7.07 (m, 2H), 7.11 - 7.18 (m, 3H), 7.74 (d, J = 8.8 Hz, 1H).

Example 145

3-Chloro-1H-indole-2-carboxylic acid (1R-dimethylcarbamoyl-2-phenyl-ethyl)-amide

To a solution of 2,3-dihydro-1H-indole-2-carboxylic acid (1R-dimethylcarbamoyl-2-phenyl-ethyl)-amide (less polar isomer, 0.50 g, 1.42 mmol) in DMF (7.5 mL) was added N-chlorosuccinimide (0.55 g, 1.42 mmol). After overnight stirring, the solvent was evaporated and the product purified by flash-chromatography (hexanes / ethyl acetate, 1:1).

¹H NMR (300 MHz, CDCl₃) δ 2.55 (s, 3H), 2.80 (s, 3H), 3.05-3.20 (m, 2H), 5.32 (m, 1H), 7.10-7.25 (m, 6H), 7.30 (d, 7 Hz, 1H), 7.58 (d, 7 Hz, 1H), 8.11 (br d, 7 Hz, 1H), 10.20 (br, 1H). MS m/e 370 (M⁺ + 1).

Example 146

3-Chloro-1H-indole-2-carboxylic acid (1S-dimethylcarbamoyl-2-phenyl-ethyl)-amide

The title compound was prepared by an analogous method to that of Example 145 from the more polar isomer of 2,3-Dihydro-1H-indole-2-carboxylic acid (1R-dimethylcarbamoyl-2-phenyl-ethyl)-amide.

¹H NMR (300 MHz, CDCl₃) δ 2.55 (s, 3H), 2.85 (s, 3H), 3.05 - 3.20 (m, 2H), 5.32 (m, 1H), 7.10 - 7.25 (m, 6H), 7.35 (d, 7 Hz, 1H), 7.58 (d, 7 Hz, 1H), 8.11 (br d, 7 Hz, 1H), 10.30 (br, 1H). MS m/e 370 (M⁺ + 1).

HPLC conditions for Examples 147-165: Detector wavelength 215 nm.

HPLC retention time (in minutes) from a Waters Novapac C18 3.9 X 150 mm column.

Eluent A = 50 mM KH₂PO₃, pH 3; Eluent B = Acetonitrile; Flow rate 1.5 mL/minute;

Gradient 90 % A / 10 % B (5 minutes) to 40 % A / 60 % B (5 minutes hold). HPLC retention times (RT) are in minutes. The percent value given is the percent of total

integration due to the specified peak.

By HPLC, the starting acid was present in an amount less than 5% of the total integration, unless specified otherwise.

-130-

Example 147

(S) 2-[(5-Fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid

To a solution of 5-fluoroindole-2-carboxylic acid (5.0 g, 28 mmol) and methylene chloride (250 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (5.53 g, 27.9 mmol), L-phenylalanine t-butyl ester hydrochloride (6.54 g, 27.9 mmol) and triethyl amine (7.1 mL, 51.3 g, 51 mmol). After stirring for 40 hours at room temperature, the reaction mixture was washed with an equal volume of water and then an equal volume of 1 N HCl. The aqueous acid was extracted with methylene chloride and the combined organic layers were sequentially washed with equal volumes of water (twice) and brine. The organic solution was dried (MgSO_4), filtered and concentrated to give S-t-butyl 2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionate (2.97 g, 31 %). This was then diluted with methylene chloride (75 mL) and cooled to 0 °C. Trifluoro-acetic acid (8 mL) was added and the reaction was then stirred at room temperature for 2 days and then heated to reflux for 6.5 hours. After allowing to come to room temperature over night, the solution was concentrated to dryness to yield a brown solid. This was then dissolved in a small amount of ether and pentanes, filtered to remove particulates and concentrated to give the title compound as a brown foam (2.65 g, quantitative yield): mp 125 - 127 °C; HPLC RT 5.72; TSPMS ion (expected) 327(326);

^1H NMR (CDCl_3) δ 9.0 (br s, 2H), 7.4-7.2 (m, 6H), 7.02 (dt, $J = 2.4, 9.1$ Hz, 1H), 6.80 (d, $J = 7.7$ Hz, 1H), 6.75 (d, $J = 1.6$ Hz, 1H), 5.09 (q, $J = 7.6$ Hz, 1H), 3.35 (dd, $J = 5.8, 7.6$ Hz, 1H), 3.26 (dd, $J = 5.8, 7.6$ Hz, 1H).

Example 148

(S)- 2-[(5-Methyl-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid

A repeat of the above procedure with 5-methylindole-2-carboxylic acid (3.0 g, 17 mmol), methylene chloride (185 mL), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (3.28g, 17.1 mmol), L-phenylalanine t-butyl ester hydrochloride (4.01 g, 15.6 mmol) and triethyl amine (4.5 mL, 3.31 g, 32.7 mmol) afforded the analogous t-butyl ester (2.42, 41 %). After dilution with methylene chloride (60 mL) and trifluoroacetic acid (6.6 mL), the reaction was heated to reflux for 3 hours, allowed to come to room temperature over night and concentrated. The crude product was slurried in ethyl acetate, filtered to remove insoluble material and concentrated (twice) to give the title compound as a brown foam (2.54 g, quantitative yield).

-131-

HPLC RT 5.98; TSPMS ion (expected) 323 (322);

¹H NMR (CDCl₃) δ 9.9 (br s, 1H), 8.5 (br s, 2H), 7.38 (s, 1H), 7.3-7.1 (m, 6H), 6.77 (m, 2H), 5.09 (q, J = 7.6 Hz, 1H), 3.35 (dd, J = 5.6, 7.6 Hz, 1H), 3.26 (dd, J = 5.6, 7.6 Hz, 1H), 2.43 (s, 3H).

5

Example 149

5-Fluoro-1H-indole-2-carboxylic acid {1-[2-(5-methoxy-1H-indol-3-yl)-ethylcarbamoyl]-2-phenyl-ethyl}-amide

To 5.0 μmol of 2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid (50 μL of a 0.1 mM solution in dimethylformamide) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (50 μL of a 0.11 mM solution in DMF, 5.5 μmol) followed by 5-methoxytryptamine (50 μL of a 0.11 mM solution in DMF, 5.5 μmol). The reaction was agitated for 3 days and then concentrated to dryness. The crude product was partitioned between chloroform (0.5 mL) and water (0.25 mL) and the organic layer was then concentrated to give the title compound.

15 TSPMS ion (expected) 499 (499); HPLC RT 6.78 (25%).

For the following examples, prepared analogously to Example 149, Examples 150 - 156 utilize 2-[(5-fluoro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid and Example 157-163 utilize 2-[(5-methyl-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid.

20

Example 150

5-Fluoro-1H-indole-2-carboxylic acid {1-[2-(1H-indol-3-yl)-1-methyl-ethylcarbamoyl]-2-phenyl-ethyl}-amide

TSPMS ion (expected) 482 (483); HPLC RT unknown, several small peaks noted; est. purity <10 %; % SM (HPLC) ND.

25

Example 151

5-Fluoro-1H-indole-2-carboxylic acid [1-benzyl-2-(2-ethyl-piperidin-1-yl)-2-oxo-ethyl]-amide

TSPMS ion (expected) 420 (421); HPLC RT 6.61 (40 %).

Example 152

30

5-Fluoro-1H-indole-2-carboxylic acid (1-cyclohexylcarbamoyl-2-phenyl)-amide

TSPMS ion (expected) 408 (407); HPLC RT 6.60 / 7.11 (The two largest peaks are of roughly equal concentration.); est. purity (25 %).

Example 153

5-Fluoro-1H-indole-2-carboxylic acid {2-phenyl-1-
[(thiophen-2-ylmethyl)-carbamoyl]-ethyl}-amide

TSPMS ion (expected) 422 (421); HPLC RT 7.50 (50 %).

5

Example 154

5-Fluoro-1H-indole-2-carboxylic acid [1-benzyl-2-(3,4-dihydro-1H-
isoquinolin-2-yl)-2-oxo-ethyl]-amide

TSPMS ion (expected) 442 (441); HPLC RT 6.78 (35 %), 5% SM.

Example 155

10

5-Fluoro-1H-indole-2-carboxylic acid [1-(2-cyclohexen-1-yl-
ethylcarbamoyl)-2-phenyl-ethyl]-amide

TSPMS ion (expected) 434 (433); HPLC RT 6.27 / 6.60 (The two largest peaks are of roughly equal concentration.); est. purity (35 %), 5% SM.

Example 156

15

5-Fluoro-1H-indole-2-carboxylic acid [1-(5-cyano-pentyl-carbamoyl)-
2-phenyl-ethyl]-amide

TSPMS ion (expected) 421 (420); HPLC RT 6.61 / 7.71 (The two largest peaks are of roughly equal concentration.) (40 %).

Example 157

20

5-Methyl-1H-indole-2-carboxylic acid [2-phenyl-1-
(thiochroman-4-ylcarbamoyl)-ethyl]-amide

TSPMS ion (expected) 470 (470).

Example 158

5-Methyl-1H-indole-2-carboxylic acid (1-cyclohexylcarbamoyl-2-phenyl-ethyl)-amide

25 TSPMS ion (expected) 404 (404); HPLC RT 6.21 (70 %).

Example 159

5-Methyl-1H-indole-2-carboxylic acid (1-benzyl-2-morpholin-4-yl-2-oxo-ethyl)-amide

TSPMS ion (expected) 392 (391); HPLC RT 6.86 (50 %).

Example 160

30 5-Methyl-1H-indole-2-carboxylic acid (1-benzyl-2-oxo-2-pyrrolidin-1-yl-ethyl)-amide

TSPMS ion (expected) 376 (375); HPLC RT 6.50 (40 %).

-133-

Example 161

5-Methyl-1H-indole-2-carboxylic acid {2-phenyl-1-
[(thiophen-2-ylmethyl)-carbamoyl]-ethyl}-amide

TSPMS ion (expected) 418 (417); HPLC RT 7.89 (70 %).

5

Example 162

5-Methyl-1H-indole-2-carboxylic acid
[1-(5-cyano-pentylcarbamoyl)-2-phenyl-ethyl]-amide

TSPMS ion (expected) 417 (417); HPLC RT 6.49 / 6.88 (The two largest peaks are of roughly equal concentration.); (40 %).

10

Example 163

5-Methyl-1H-indole-2-carboxylic acid (1-cyclopentylcarbamoyl-2-phenyl-ethyl)-amide

TSPMS ion (expected) 390 (389); HPLC RT 6.96 (55 %).

Example 164

{2-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-
phenyl-propionylamino}-acetic acid methyl ester

15

To 5.0 mmol of 5-chloro-1H-indole-2-carboxylic acid (50 mL of a 0.1 M solution in acetonitrile) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (50 mL of a 0.10 M solution in acetonitrile, 5.0 mmol), 1-hydroxybenzotriazole (50 mL of a 0.10 M solution in acetonitrile, 5.0 mmol), followed by (2-amino-3-phenyl-propionylamino)-acetic acid methyl ester (50 mL of a 0.10 M solution in acetonitrile, 5.0 mmol). The reaction was agitated overnight at 80 °C and then concentrated to dryness to give the title compound.

HPLC RT 8.15 (65 %).

Example 165

25 2-(S)-[(5-Chloro-1H-indole-2-carbonyl)-amino]-3-phenyl-propionic acid benzyl ester

The title compound was prepared by substituting L-phenylalanine benzyl ester for (2-amino-3-phenyl-propionyl-amino)-acetic acid methyl ester in a method analogous to that of Example 164 procedure.

HPLC RT 8.13 (40 %).

-134-

Example 166

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-
azetidin-1-yl)-2-oxo-ethyl]-amide

2-Amino-1-(3-hydroxy-azetidin-1-yl)-3-phenyl-propan-1-one hydrochloride (1.18 mmol)
5 and 5-chloro-1H-indole-2-carboxylic acid (1.18 mmol) were coupled according to
Procedure A (4:1 dichloromethane-dimethylformamide reaction solvent) and the
product purified by chromatography on silica gel eluted with 25%, 50%, 75% and
100% ethyl acetate-hexanes giving the title substance as a colorless foam (104 mg,
22%). A mixture (180 mg) of less polar products was also isolated. Title substance:
10 HPLC (60/40) 4.18 minutes (97%); TSPMS 398/400 (MH⁺, 100%);

Example 166a

(2S)-Amino-1-(3-hydroxy-azetidin-1-yl)-3-phenyl-propan-1-one hydrochloride

[(1S)-Benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester
(515 mg, 1.6 mmol) was dissolved in cold 4N HCl-dioxane, the mixture stirred 2 h at
15 25°C, concentrated, and the residue coevaporated with ether giving a colorless
solid (415 mg, 100%).

Example 167

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxyimino-
azetidin-1-yl)-2-oxo-ethyl]-amide

20 A solution of 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-oxo-azetidin-1-
yl)-2-oxo-ethyl]-amide (product of Example 170, 50 mg, 0.13 mmol), sodium acetate
trihydrate (43 mg, 0.32 mmol) and hydroxylamine hydrochloride (18 mg, 0.25 mmol)
in methanol (2 mL) was heated at reflux for 8 h and concentrated. The residue was
partitioned between dichloromethane and saturated aqueous NaHCO₃. The organic
25 layer was separated and dried giving a colorless solid which was triturated with
ether-hexanes and dried (yield 36 mg, 69%): HPLC (50/50) 6.74 min (99%); TSPMS
411/413 (MH⁺, 10%), 180 (100%); ¹H NMR (DMSO-d₆) δ 11.75 (br, 1H), 11.10 (s,
0.5H), 11.08 (s, 0.5H), 8.99 (d, 1H, J = 9 Hz), 7.73 (d, 1H, J = 2 Hz), 7.4-7.1 (m,
8H), 5.0 (m, 1H), 4.8-4.5 (m, 4H), 3.1 (m, 2H).

Example 168

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(4-hydroxyimino-piperidin-1-yl)-2-oxo-ethyl]-amide

A mixture of 5-chloro-1H-indole-2-carboxylic acid [1(S)-benzyl-2-oxo-2-(4-oxo-
5 piperidin-1-yl)-ethyl]-amide (406 mg, 0.96 mmol), hydroxylamine hydrochloride (80 mg, 1.15 mmol), and potassium carbonate, (159 mg, 1.15 mmol) in ethanol (6 mL) and water (1 mL) was stirred at 25 °C for 18 h and concentrated. The residue was dissolved in ethyl acetate and the resulting solution washed with water and dried (411 mg, 98%): HPLC (60/40) 5.13 minutes (97%); TSPMS 439/441 (MH⁺, 100%);
10 ¹H NMR (DMSO-d₆) δ 11.75 (br, 1H), 10.45 (s, 0.5H), 10.44 (s, 0.5H), 9.00 (m, 1H), 7.72 (d, 1H, J = 2 Hz), 7.40 (d, 1H, J = 8.8 Hz), 7.35-7.15 (m, 7H), 5.17 (m, 1H), 3.8-3.5 (m, 4H), 3.1 (m, 2H), 2.45 (m, 2H), 2.25 (m, 2H).

Example 168a

5-Chloro-1H-indole-2-carboxylic acid [1(S)-benzyl-2-oxo-2-(4-oxo-piperidin-1-yl)-ethyl]-amide

15 5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide (Example 46, 669 mg) was added in one portion at 0 °C to a mixture of 1-(3-dimethylaminopropyl)3-ethylcarbodiimide hydrochloride (DEC, 1.80 g, 9.4 mmol) and dichloroacetic acid (307 mg, 1.5 mmol) in anhydrous toluene (e mL) and
20 anhydrous dimethylsulfoxide (e mL). The mixture was stirred at 0-20 °C for 2h, diluted with ethyl acetate, the resulting solution washed twice with 1N HCl, twice with saturated aqueous NaHCO₃, dried, concentrated and the residue purified by chromatography on silica gel eluted with 25%, 50%, and 75% ethyl acetate-hexanes giving a foam (424 mg, 64%).

25

Example 169

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(1,3-dihydro-isoindol-2-yl)-2-oxo-ethyl]-amide

2-Amino-1-(1,3-dihydro-isoindol-2-yl)-3-phenyl-propan-1-one hydrochloride (0.20 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.20 mmol) were coupled according
30 to Procedure A and the product purified by chromatography on silica gel eluted with 5%, 10%, 20%, and 50% ethyl acetate-hexanes (55mg, 62% yield): HPLC (70/30) 6.58 minutes (90%); TSPMS 444/446 (MH⁺, 50%), 180 (100%).

-136-

¹H NMR (CDCl₃) δ 9.25 (br, 1H), 7.60 (s, 1H), 7.45 (m, ca. 1H), 7.3-7.1 (m, ca. 11H), 6.90 (5.25 (m, 1H), 5.0 (d, 1H, ca. 16 Hz), 4.85 (d, 1H, J = ca. 16 Hz), 4.70 (d, 1H, J = ca. 16 Hz), 4.20 (d, 1H, J = 16 Hz).

Example 169a

5 (2S)-Amino-1-(1,3-dihydro-isoindol-2-yl)-3-phenyl-
 propan-1-one hydrochloride

[(1S)-Benzyl-2-(1,3-dihydro-isoindol-2-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (88 mg) was dissolved in cold 4N HCl-dioxane (1.5 mL), stirred 2 h at 25 °C, and the mixture concentrated. The residue was triturated with ether and dried (65 mg,
10 91%). TSPMS 267 (MH⁺, 100%).

Example 169b

[(1S)-Benzyl-2-(1,3-dihydro-isoindol-2-yl)-2-oxo-ethyl]-carbamic
 acid tert-butyl ester

N-t-Boc-L-phenylalanine (1 mmol) and isoindoline (J. Org. Chem. 1988, 53, p5382,
15 70-80% purity, 1 mmol) were coupled according to Procedure A and the product purified by chromatography on silica gel eluted with 20% and 50% ethyl acetate-hexanes giving an amber oil (88 mg, 23%): TSPMS 367 (MH⁺, 100%).

Example 170

20 5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-oxo-
 azetidin-1-yl)-2-oxo-ethyl]-amide

1-((2S)-Amino-3-phenyl-propionyl)-azetidin-3-one hydrochloride (3.2 mmol) and 5-chloro-1H-indole-2-carboxylic acid (3.2 mmol) were coupled according to Procedure A (0-25 °C reaction temperature) and the resulting yellow foam purified by chromatography on silica gel eluted with 20%, 30%, 40% and 50% ethyl acetate in
25 hexane giving the title substance as a colorless foam (600 mg, 47%): HPLC (60/40) 5.09 minutes (98%); TSP-MS 396 (MH⁺, 100%); ¹H NMR (CDCl₃) δ 9.14 (br, 1H), 7.62 (d, 1H, J = 3 Hz), 7.4-7.2 (m, 7H), 7.11 (d, 1H, J = 8.0 Hz), 6.85 (m, 1H), 4.90 (m, 1H), 4.78 (m, 2H), 4.63 (m, 1H), 3.65 (m, 1H), 3.25 (dd, 1H, A of AB, J = 5.1, 12.9 Hz), 3.10 (dd, 1H, B of AB, J = 10, 12.9 Hz).

30

Example 170A

1-((2S)-Amino-3-phenyl-propionyl)-azetidin-3-one hydrochloride

[(1S)-Benzyl-2-oxo-2-(3-oxo-azetidin-1-yl)-ethyl]-carbamic acid tert-butyl ester (297 mg, 0.9 mmol) was dissolved in 4N HCl-dioxane (3 mL). The resulting solution was

-137-

stirred at 25 °C for 2 h, concentrated, and the residue triturated with ether and dried (196 mg, 82%).

Example 170B

[(1S)-Benzyl-2-oxo-2-(3-oxo-azetidin-1-yl)-ethyl]-carbamic

5

acid tert-butyl ester

[(1S)-Benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-carbamic acid tert-butyl ester (320 mg, 1 mmol) was added in one portion to a mixture of 1-(3-dimethylaminopropyl)3-ethylcarbodiimide hydrochloride (DEC, 575 mg, 3 mmol) and dichloroacetic acid (192 mg, 1.5 mmol) in anhydrous toluene (2 mL) and anhydrous
10 dimethylsulfoxide (2 mL). The mixture was stirred at 0-20°C for 1h, diluted with ethyl acetate, the resulting solution washed twice with 1N HCl, twice with saturated aqueous NaHCO₃, dried and concentrated giving a colorless solid (304 mg, 96%).

Example 170C

[(1S)-Benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-carbamic

15

acid tert-butyl ester

3-Hydroxyazetidine hydrochloride (J. Chem. Soc., Chem. Commun. 1968, p93, 27 mmol) and N-t-Boc-L-phenylalanine (27 mmol) were coupled according to Procedure A giving the title substance as a colorless foam (8.15 g, 93%).

Example 171

20

5-Chloro-1H-benzoimidazole-2-carboxylic acid

(1-dimethylcarbamoyl-2-phenyl-ethyl)-amide

(S)-2-Amino-N,N-dimethyl-3-phenyl-propionamide hydrochloride (2.0 mmol) and 5-chloro-1H-benzoimidazole-2-carboxylic acid (Crowther et al., J. Chem. Soc. 1949, p.1268, 2.0 mmol) were coupled according to Procedure A and the product purified
25 by chromatography on silica gel eluted with 1:1 ethyl acetate-hexanes (235 mg, 63%): HPLC (60/40) 4.92 min (91%); PBMS 371/373 (MH⁺, 100%); ¹H NMR (CDCl₃) δ 11.25 (br, 0.6H), 10.9 (br, 0.4H), 8.36 (m, 1H), 7.78 (d, 0.4H, J = 7.72 (d, 0.6H, J = 8.8 Hz), 7.52 (d, 0.6H, J = 2 Hz), 7.41 (d, 0.4H, J = 8.4 Hz), 7.35-7.1 (m, 6H), 7.35 (m, 1H), 3.16 (m, 2H), 2.90 (s, 3H), 2.68 (s, ca. 2H), 2.67 (s, ca. 1H).

-138-

Example 172

5-Chloro-1H-indole-2-carboxylic acid [1-benzyl-2-oxo-2-(2-oxo-oxazolidin-3-yl)-ethyl]-amide

3-((2S)-Amino-3-phenyl-propionyl)-oxazolidin-2-one hydrochloride (0.50 mmol) and 5-chloro-1H-indole-2-carboxylic acid (0.50 mmol) were coupled according to Procedure A (3:1 dimethylformamide-dichloromethane reaction solvent) and the product triturated with 2:1 ether-hexanes and dried (130 mg, 63%): HPLC (60/40) 6.22 minutes (95%); TSPMS 429/431 (45%, MH+NH₃), 412/414 (30%, MH+), 325/327 (100%). ¹H NMR (DMSO-d₆) δ 11.68 (br, 1H), 8.92 (d, 1H, J = 8.5 Hz), 7.75 (s, 1H), 7.42 (m, 3H), 7.26 (m, 3H), 7.18 (m, 2H), 5.83 (m, 1H), 4.50 (m, 2H), 4.0 (m, 1H), 3.25 (m, 1H), 2.95 (m, 1H).

Example 172a

3-((2S)-Amino-3-phenyl-propionyl)-oxazolidin-2-one hydrochloride

[(1S)-Benzyl-2-oxo-2-(2-oxo-oxazolidin-3-yl)-ethyl]-carbamic acid tert-butyl ester (2.29 g, 6.68 mmol) was dissolved in 4N HCl-dioxane (10 mL) at 0 °C. The resulting solution was stirred at 25 °C for 2h, concentrated, and the residue triturated with ether and dried (1.98 g, 107%).

Example 172b

[(1S)-Benzyl-2-oxo-2-(2-oxo-oxazolidin-3-yl)-ethyl]-carbamic acid tert-butyl ester


N-Butyllithium (2.35 M in hexanes, 11.5 mL) was added at -78 °C to a solution of 2-oxazolidinone (2.04 g, 23.4 mmol) in tetrahydrofuran (25 mL). After 30 minutes at -78 °C the solution was treated with N-t-Boc-L-phenylalanine N-hydroxysuccinimide ester (9.31 g, 25.7 mmol) in tetrahydrofuran (10 mL), and the stirred mixture was allowed to warm to 25 °C overnight. Water (10 mL) was added, and the resulting mixture concentrated, the residue dissolved in ethyl acetate, and the resulting solution washed twice with 1N NaOH, once with water, once with brine, dried, and concentrated. The residue was chromatographed on silica gel eluted with 25% and 50% ethyl acetate in hexanes giving a colorless solid (3.42 g, 44%).

It should be understood that the invention is not limited to the particular embodiments described herein, but that various changes and modifications may be made without departing from the spirit and scope of this novel concept as defined by the following claims.

-139-

CLAIMS

1. A compound of Formula I



Formula I

and the pharmaceutically acceptable salts and prodrugs thereof
wherein

the dotted line (---) is an optional bond;

A is $-C(H)=$, $-C((C_1-C_4)alkyl)=$, $-C(halo)=$ or $-N=$, when the dotted line (---) is
a bond, or A is methylene or $-CH((C_1-C_4)alkyl)-$, when the dotted line (---) is not a
bond;

R_1 , R_{10} or R_{11} are each independently H, halo, cyano, 4-, 6-, or 7-nitro, $(C_1-$
 $C_4)alkyl$, $(C_1-C_4)alkoxy$, fluoromethyl, difluoromethyl or trifluoromethyl;

R_2 is H;

R_3 is H or $(C_1-C_5)alkyl$;

R_4 is H, methyl, ethyl, n-propyl, hydroxy $(C_1-C_3)alkyl$, $(C_1-C_3)alkoxy(C_1-$
 $C_3)alkyl$, phenyl $(C_1-C_4)alkyl$, phenylhydroxy $(C_1-C_4)alkyl$, (phenyl)((C_1-C_4)-alkoxy) $(C_1-$
 $C_4)alkyl$, thien-2- or -3-yl $(C_1-C_4)alkyl$ or fur-2- or -3-yl $(C_1-C_4)alkyl$ wherein said R_4
rings are mono-, di- or tri-substituted independently on carbon with H, halo, $(C_1-$
 $C_4)alkyl$, $(C_1-C_4)alkoxy$, trifluoromethyl, hydroxy, amino, cyano or 4,5-dihydro-1H-
imidazol-2-yl; or

R_4 is pyrid-2-, -3- or -4-yl $(C_1-C_4)alkyl$, thiazol-2-, -4- or -5-yl $(C_1-C_4)alkyl$,
imidazol-2-, -4- or -5-yl $(C_1-C_4)alkyl$, pyrrol-2- or -3-yl $(C_1-C_4)alkyl$, oxazol-2-, -4- or -5-
yl $(C_1-C_4)alkyl$, pyrazol-3-, -4- or -5-yl $(C_1-C_4)alkyl$, isoxazol-3-, -4- or -5-yl $(C_1-C_4)alkyl$,
isothiazol-3-, -4- or -5-yl $(C_1-C_4)alkyl$, pyridazin-3- or -4-yl $(C_1-C_4)alkyl$, pyrimidin-2-, -4-,
-5- or -6-yl $(C_1-C_4)alkyl$, pyrazin-2- or -3-yl $(C_1-C_4)alkyl$, 1,3,5-triazin-2-yl $(C_1-C_4)alkyl$ or
indol-2- $(C_1-C_4)alkyl$, wherein said preceding R_4 heterocycles are optionally mono- or

-140-

di-substituted independently with halo, trifluoromethyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, amino, hydroxy or cyano and said substituents are bonded to carbon; or

- R₄ is R₁₅-carbonyloxymethyl, wherein said R₁₅ is phenyl, thiazolyl, imidazolyl, 1H-indolyl, furyl, pyrrolyl, oxazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl or 1,3,5-triazinyl and wherein said preceding R₁₅ rings are optionally mono- or di-substituted independently with halo, amino, hydroxy, (C₁-C₄)alkyl, (C₁-C₄)alkoxy or trifluoromethyl and said mono- or di-substituents are bonded to carbon;

R₅ is H, methyl, ethyl, n-propyl, hydroxymethyl or hydroxyethyl;

- 10 R₆ is carboxy, (C₁-C₈)alkoxycarbonyl, benzyloxycarbonyl, C(O)NR₈R₉ or C(O)R₁₂ wherein

R₈ is H, (C₁-C₆)alkyl, cyclo(C₃-C₆)alkyl, cyclo(C₃-C₆)alkyl(C₁-C₅)alkyl, hydroxy or (C₁-C₈)alkoxy; and

- 15 R₉ is H, cyclo(C₃-C₈)alkyl, cyclo(C₃-C₈)alkyl(C₁-C₅)alkyl, cyclo(C₄-C₇)alkenyl, cyclo(C₃-C₇)alkyl(C₁-C₅)alkoxy, cyclo(C₃-C₇)alkyloxy, hydroxy, methylene-perfluorinated(C₁-C₈)alkyl, phenyl, or a heterocycle wherein said heterocycle is pyridyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, pyridinyl, piperidinyl, morpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, thiochromanyl or tetrahydrobenzothiazolyl wherein said heterocycle rings are carbon-nitrogen linked; or

- R₉ is (C₁-C₆)alkyl or (C₁-C₈)alkoxy wherein said (C₁-C₆)alkyl or (C₁-C₈)alkoxy is optionally monosubstituted with cyclo(C₄-C₇)alken-1-yl, phenyl, thienyl, pyridyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, piperidinyl, morpholinyl, thiomorpholinyl, 1-oxothiomorpholinyl, 1,1-dioxothiomorpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl or indolyl and wherein said (C₁-C₆)alkyl or (C₁-C₈)alkoxy are optionally additionally independently mono- or di-substituted with halo, hydroxy, (C₁-C₅)alkoxy, amino, mono-N- or di-N,N-(C₁-C₅)alkylamino, cyano, carboxy, or (C₁-C₄)alkoxycarbonyl; and

-141-

wherein the R_9 rings are optionally mono- or di-substituted independently on carbon with halo, (C_1-C_4) alkyl, (C_1-C_4) alkoxy, hydroxy, hydroxy (C_1-C_4) alkyl, amino (C_1-C_4) alkyl, mono-N- or di-N,N- (C_1-C_4) alkylamino (C_1-C_4) alkyl, (C_1-C_4) alkoxy (C_1-C_4) alkyl, amino, mono-N- or di-N,N- (C_1-C_4) alkylamino, cyano, carboxy, 5 (C_1-C_5) alkoxycarbonyl, carbamoyl, formyl or trifluoromethyl and said R_9 rings may optionally be additionally mono- or di-substituted independently with (C_1-C_5) alkyl or halo;

with the proviso that no quaternized nitrogen on any R_9 heterocycle is included;

10 R_{12} is morpholino, thiomorpholino, 1-oxothiomorpholino, 1,1-dioxothiomorpholino, thiazolidin-3-yl, 1-oxothiazolidin-3-yl, 1,1-dioxothiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, piperazin-4-yl, azetidin-1-yl, 1,2-oxazinan-2-yl, pyrazolidin-1-yl, isoxazolidin-2-yl, isothiazolidin-2-yl, 1,2-oxazetidin-2-yl, oxazolidin-3-yl, 3,4-dihydroisoquinolin-2-yl, 1,3-dihydroisoindol-2-yl, 3,4-dihydro-2H- 15 quinol-1-yl, 2,3-dihydro-benzo[1,4]oxazin-4-yl, 2,3-dihydro-benzo[1,4]thiazine-4-yl, 3,4-dihydro-2H-quinoxalin-1-yl, 3,4-dihydro-benzo[c][1,2]oxazin-1-yl, 1,4-dihydro-benzo[d][1,2]oxazin-3-yl, 3,4-dihydro-benzo[e][1,2]-oxazin-2-yl, 3H-benzo[d]isoxazol-2-yl, 3H-benzo[c]isoxazol-1-yl or azepan-1-yl,

wherein said R_{12} rings are optionally mono-, di- or tri-substituted 20 independently with halo, (C_1-C_5) alkyl, (C_1-C_5) alkoxy, hydroxy, amino, mono-N- or di-N,N- (C_1-C_5) alkylamino, formyl, carboxy, carbamoyl, mono-N- or di-N,N- (C_1-C_5) alkylcarbamoyl, (C_1-C_6) alkoxy (C_1-C_3) alkoxy, (C_1-C_5) alkoxycarbonyl, benzyloxycarbonyl, (C_1-C_5) alkoxycarbonyl (C_1-C_5) alkyl, (C_1-C_4) alkoxycarbonylamino, carboxy (C_1-C_5) alkyl, carbamoyl (C_1-C_5) alkyl, mono-N- or di-N,N- (C_1-C_5) alkylcarbamoyl (C_1-C_5) alkyl, hydroxy (C_1-C_5) alkyl, (C_1-C_4) alkoxy (C_1-C_4) alkyl, 25 amino (C_1-C_4) alkyl, mono-N- or di-N,N- (C_1-C_4) alkylamino (C_1-C_4) alkyl, oxo, hydroxyimino or (C_1-C_6) alkoxyimino and wherein no more than two substituents are selected from oxo, hydroxyimino or (C_1-C_6) alkoxyimino and oxo, hydroxyimino or (C_1-C_6) alkoxyimino are on nonaromatic carbon; and

30 wherein said R_{12} rings are optionally additionally mono- or di-substituted independently with (C_1-C_5) alkyl or halo;

with the proviso that when R_6 is (C_1-C_5) alkoxycarbonyl or benzyloxycarbonyl then R_1 is 5-halo, 5- (C_1-C_4) alkyl or 5-cyano and R_4 is (phenyl)(hydroxy) (C_1-C_4) alkyl,

(phenyl)((C₁-C₄)alkoxy)(C₁-C₄)alkyl, hydroxymethyl or Ar(C₁-C₂)alkyl, wherein Ar is thien-2- or -3-yl, fur-2- or -3-yl or phenyl wherein said Ar is optionally mono- or di-substituted independently with halo; with the provisos that when R₄ is benzyl and R₅ is methyl, R₁₂ is not 4-hydroxy-piperidin-1-yl or when R₄ is benzyl and R₅ is methyl R₆

5 is not C(O)N(CH₃)₂;

with the proviso that when R₁ and R₁₀ and R₁₁ are H, R₄ is not imidazol-4-ylmethyl, 2-phenylethyl or 2-hydroxy-2-phenylethyl;

with the proviso that when both R₈ and R₉ are n-pentyl, none of R₁ is 5-chloro, 5-bromo, 5-cyano, 5(C₁-C₅)alkyl, 5(C₁-C₅)alkoxy or trifluoromethyl;

10 with the proviso that when R₁₂ is 3,4-dihydroisoquinol-2-yl, said 3,4-dihydroisoquinol-2-yl is not substituted with carboxy((C₁-C₄)alkyl);

with the proviso that when R₈ is H and R₉ is (C₁-C₆)alkyl, R₉ is not substituted with carboxy or (C₁-C₄)alkoxycarbonyl on the carbon which is attached to the nitrogen atom N of NHR₉; and

15 with the proviso that when R₆ is carboxy and R₁, R₁₀, R₁₁ and R₅ are all H, then R₄ is not benzyl, H, (phenyl)(hydroxy)methyl, methyl, ethyl or n-propyl.

2. A compound as recited in claim 1 wherein

R₁ is 5-H, 5-halo, 5-methyl, 5-cyano or 5-trifluoromethyl;

R₁₀ and R₁₁ are each independently H or halo;

20 A is -C(H)=;

R₂ and R₃ are H;

R₄ is H, methyl, phenyl(C₁-C₂)alkyl, wherein said phenyl groups are mono- or di-substituted independently with H, halo, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, trifluoromethyl, hydroxy, amino or cyano and wherein said R₄ groups are optionally additionally

25 mono-substituted with halo; or

R₄ is thien-2- or -3-yl(C₁-C₂)alkyl, pyrid-2-, -3- or -4-yl(C₁-C₂)alkyl, thiazol-2-, -4- or -5-yl(C₁-C₂)alkyl, imidazol-2-, -4- or -5-yl(C₁-C₂)alkyl, fur-2- or -3-yl(C₁-C₂)alkyl, pyrrol-2- or -3-yl(C₁-C₂)alkyl, oxazol-2-, -4- or -5-yl(C₁-C₂)alkyl, pyrazol-3-, -4- or -5-yl(C₁-C₂)alkyl, isoxazol-3-, -4- or -5-yl(C₁-C₂)alkyl, isothiazol-3-, -4- or -5-yl(C₁-C₂)alkyl, pyridazin-3- or -4-yl(C₁-C₂)alkyl, pyrimidin-2-, -4-, -5- or -6-yl(C₁-C₂)alkyl, pyrazin-2- or -3-yl(C₁-C₂)alkyl or 1,3,5-triazin-2-yl(C₁-C₂)alkyl wherein said preceding R₄ heterocycles are optionally mono- or di-substituted independently with halo,

30

-143-

trifluoromethyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, amino or hydroxy and said mono- or di-substituents are bonded to carbon;

R₅ is H; and

R₆ is C(O)NR₈R₉ or C(O)R₁₂.

5 3. A compound as recited in claim 2 wherein

R₄ is H, phenyl(C₁-C₂)alkyl, thien-2- or -3-yl(C₁-C₂)alkyl, fur-2- or -3-yl(C₁-C₂)alkyl wherein said R₄ rings are mono- or di-substituted independently with H or fluoro;

R₆ is C(O)R₁₂; and

10 R₁₂ is morpholino, thiomorpholino, 1-oxothiomorpholino, 1,1-dioxothiomorpholino, thiazolidin-3-yl, 1-oxothiazolidin-3-yl, 1,1-dioxothiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, piperazin-4-yl, azetidin-1-yl, 1,2-oxazinan-2-yl, isoxazolidin-2-yl, isothiazolidin-2-yl, 1,2-oxazetidin-2-yl, oxazolidin-3-yl, 1,3-dihydroisoindol-2-yl, or azepan-1-yl,

15 wherein said R₁₂ rings are optionally mono- or di-substituted independently with halo, (C₁-C₅)alkyl, (C₁-C₅)alkoxy, hydroxy, amino, mono-N- or di-N,N-(C₁-C₅)alkylamino, formyl, carboxy, carbamoyl, mono-N- or di-N,N-(C₁-C₅)alkylcarbamoyl, (C₁-C₅)alkoxycarbonyl, hydroxy(C₁-C₅)alkyl, amino(C₁-C₄)alkyl, mono-N- or di-N,N-(C₁-C₄)alkylamino(C₁-C₄)alkyl, oxo, hydroxyimino or (C₁-

20 C₆)alkoxyimino with the proviso that only the R₁₂ heterocycles thiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, piperazin-4-yl, azetidin-1-yl, 1,2-oxazinan-2-yl, isoxazolidin-2-yl, or oxazolidin-3-yl are optionally mono- or di-substituted with oxo, hydroxyimino, or (C₁-C₆)alkoxyimino; and

wherein said R₁₂ rings are optionally additionally mono- or di-substituted

25 independently with (C₁-C₅)alkyl.

4. A compound as recited in claim 3 wherein

R₄ is H; and

R₁₂ is thiazolidin-3-yl, 1-oxo-thiazolidin-3-yl, 1,1-dioxo-thiazolidin-3-yl or oxazolidin-3-yl or said R₁₂ substituents optionally mono- or di-substituted

30 independently with carboxy, (C₁-C₅)alkoxycarbonyl, hydroxy(C₁-C₃)alkyl, amino(C₁-C₃)alkyl, mono-N- or di-N,N-(C₁-C₃)alkylamino(C₁-C₃)alkyl or

R₁₂ is mono- or di-substituted pyrrolidin-1-yl wherein said substituents are independently carboxy, (C₁-C₅)alkoxycarbonyl, (C₁-C₅)alkoxy, hydroxy, hydroxy(C₁-

-144-

C₃)alkyl, amino, amino(C₁-C₃)alkyl, mono-N- or di-N,N-(C₁-C₃)alkylamino(C₁-C₃)alkyl or mono-N- or di-N,N-(C₁-C₄)alkylamino; and

the R_{1,2} rings are optionally additionally independently disubstituted with (C₁-C₅)alkyl.

5 5. A compound as recited in claim 3 wherein

R₄ is phenylmethyl, thien-2- or -3-ylmethyl wherein said R₄ rings are optionally mono- or di-substituted with fluoro; and

R_{1,2} is thiazolidin-3-yl, 1-oxo-thiazolidin-3-yl, 1,1-dioxo-thiazolidin-3-yl or oxazolidin-3-yl or said R_{1,2} substituents optionally mono- or di-substituted

10 independently with carboxy or (C₁-C₅)alkoxycarbonyl, hydroxy(C₁-C₃)alkyl, amino(C₁-C₃)alkyl or mono-N- or di-N,N-(C₁-C₃)alkylamino(C₁-C₃)alkyl

or R_{1,2} is mono- or di-substituted azetidin-1-yl or mono- or di-substituted pyrrolidin-1-yl or mono- or di-substituted piperidin-1-yl wherein said substituents are independently carboxy, (C₁-C₅)alkoxycarbonyl, hydroxy(C₁-C₃)alkyl, amino(C₁-

15 C₃)alkyl, mono-N- or di-N,N-(C₁-C₃)alkylamino(C₁-C₃)alkyl, hydroxy, (C₁-C₅)alkoxy, amino, mono-N- or di-N,N-(C₁-C₅)alkylamino, oxo, hydroxyimino or (C₁-C₅)alkoxyimino; and

the R_{1,2} rings are optionally additionally mono- or di-substituted independently with (C₁-C₅)alkyl.

20 6. A compound as recited in claim 3 selected from

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxyimino-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [2-(cis-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

25 5-Chloro-1H-indole-2-carboxylic acid [2-((3S,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(cis-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

30 5-Chloro-1H-indole-2-carboxylic acid [2-(1,1-dioxo-thiazolidin-3-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid (2-oxo-2-thiazolidin-3-yl-ethyl)-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-(4-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3RS)-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [2-oxo-2-((1RS)-oxo-1-thiazolidin-3-yl)-ethyl]-amide,

5 5-Chloro-1H-indole-2-carboxylic acid [(1S)-(2-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3S,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide,

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxyimino-azetidin-1-yl)-2-oxo-ethyl]-amide or

5-Chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(4-hydroxyimino-piperidin-1-yl)-2-oxo-ethyl]-amide.

15 7. The compound as recited in claim 4 wherein

R₁ is 5-chloro;

R₁₀ and R₁₁ are H; and

R₁₂ is cis-3,4-dihydroxy-pyrrolidin-1-yl.

8. The compound as recited in claim 4 wherein

20 R₁ is 5-chloro;

R₁₀ and R₁₁ are H; and

R₁₂ is (3S,4S)-dihydroxy-pyrrolidin-1-yl.

9. The compound as recited in claim 4 wherein

R₁ is 5-chloro;

25 R₁₀ and R₁₁ are H; and

R₁₂ is 1,1-dioxo-thiazolidin-3-yl.

10. The compound as recited in claim 4 wherein

R₁ is 5-chloro;

R₁₀ and R₁₁ are H; and

30 R₁₂ is thiazolidin-3-yl.

11. A compound as recited in claim 4 wherein

R₁ is 5-chloro;

R₁₀ and R₁₁ are H; and

R_{12} is 1-oxo-thiazolidin-3-yl.

12. The compound as recited in claim 5 wherein

R_1 is 5-chloro;

R_{10} and R_{11} are H;

5 R_4 is 4-fluorobenzyl;

R_{12} is 4-hydroxypiperidin-1-yl; and

the stereochemistry of carbon (a) is (S).

13. A compound as recited in claim 5 wherein

R_1 is 5-chloro;

10 R_{10} and R_{11} are H;

R_4 is benzyl;

R_{12} is 3-hydroxypiperidin-1-yl; and

the stereochemistry of carbon (a) is (S).

14. The compound as recited in claim 5 wherein

15 R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

R_{12} is cis-3,4-dihydroxy-pyrrolidin-1-yl; and

the stereochemistry of carbon (a) is S.

20 15. A compound as recited in claim 5 wherein

R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

R_{12} is 3-hydroxyimino-pyrrolidin-1-yl; and

25 the stereochemistry of carbon (a) is (S).

16. The compound as recited in claim 5 wherein

R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is 2-fluorobenzyl;

30 R_{12} is 4-hydroxypiperidin-1-yl; and

the stereochemistry of carbon (a) is (S).

17. The compound as recited in Claim 5 wherein

R_1 is 5-chloro;

-147-

R_{10} and R_{11} are H;

R_4 is benzyl;

R_{12} is (3S,4S)-dihydroxy-pyrrolidin-1-yl; and

the stereochemistry of carbon (a) is (S).

5 18. The compound as recited in Claim 5 wherein

R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

R_{12} is 3-hydroxy-azetidin-1-yl; and

10 the stereochemistry of carbon (a) is (S).

19. The compound as recited in Claim 5 wherein

R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

15 R_{12} is 3-hydroxyimino-azetidin-1-yl; and

the stereochemistry of carbon (a) is (S).

20. The compound as recited in Claim 5 wherein

R_1 is 5-chloro;

R_{10} and R_{11} are H;

20 R_4 is benzyl;

R_{12} is 4-hydroxyimino-piperidin-1-yl; and

the stereochemistry of carbon (a) is (S).

21. A compound as recited in claim 2 wherein

R_4 is H, phenyl(C_1 - C_2)alkyl, thien-2- or -3-yl(C_1 - C_2)alkyl, fur-2- or -3-yl(C_1 -

25 C_2)alkyl wherein said R_4 rings are mono- or di-substituted independently with H or fluoro;

R_6 is $C(O)NR_8R_9$; and

R_8 is H, (C_1 - C_5)alkyl, hydroxy or (C_1 - C_4)alkoxy; and

R_9 is H, cyclo(C_4 - C_6)alkyl, cyclo(C_3 - C_6)alkyl(C_1 - C_5)alkyl, methylene-

30 perfluorinated(C_1 - C_3)alkyl, pyridyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, piperidinyl, benzothiazolyl or thiochromanyl; or

R_9 is (C_1 - C_5)alkyl wherein said (C_1 - C_5)alkyl is optionally substituted with cyclo(C_4 - C_6)alkenyl, phenyl, thienyl, pyridyl, pyrrolidinyl, oxazolyl, thiazolyl,

-148-

imidazolyl, pyrazolyl, piperidinyl, morpholinyl, thiomorpholinyl, 1-oxothiomorpholinyl, or 1,1-dioxothiomorpholinyl and wherein said (C₁-C₅)alkyl or (C₁-C₄)alkoxy is optionally additionally independently mono- or di-substituted with halo, hydroxy, (C₁-C₅)alkoxy, amino, mono-N- or di-N,N-(C₁-C₅)alkylamino, cyano, carboxy, or (C₁-

5 C₄)alkoxycarbonyl; and

wherein the R₅ rings are optionally mono- or di-substituted independently on carbon with halo, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, hydroxy, amino, mono-N- or di-N,N-(C₁-C₄)alkylamino, carbamoyl, (C₁-C₅)alkoxycarbonyl or carbamoyl.

22. A compound as recited in claim 2 wherein

10 R₄ is H, phenyl(C₁-C₂)alkyl, thien-2- or -3-yl(C₁-C₂)alkyl, fur-2- or -3-yl(C₁-C₂)alkyl wherein said R₄ rings are mono- or di-substituted independently with H or fluoro;

R₆ is C(O)NR₈R₉; and

R₈ is H, (C₁-C₅)alkyl, hydroxy or (C₁-C₄)alkoxy; and

15 R₉ is (C₁-C₄)alkoxy wherein said (C₁-C₄)alkoxy is optionally substituted with cyclo(C₄-C₆)alkenyl, phenyl, thienyl, pyridyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, piperidinyl, morpholinyl, thiomorpholinyl, 1-oxothiomorpholinyl, or 1,1-dioxothiomorpholinyl and wherein said (C₁-C₅)alkyl or (C₁-C₄)alkoxy is optionally additionally independently mono- or di-substituted with halo, hydroxy, (C₁-C₅)alkoxy, amino, mono-N- or di-N,N-(C₁-C₅)alkylamino, cyano, carboxy, or (C₁-C₄)alkoxycarbonyl; and

wherein the R₉ rings are optionally mono- or di-substituted independently on carbon with halo, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, hydroxy, amino, mono-N- or di-N,N-(C₁-C₄)alkylamino, carbamoyl, (C₁-C₅)alkoxycarbonyl or carbamoyl.

25 23. A compound as recited in claim 21 wherein

R₁ is 5-chloro;

R₁₀ and R₁₁ are H;

R₄ is benzyl;

R₈ is methyl; and

30 R₉ is 3-(dimethylamino)propyl.

24. The compound as recited in claim 21 wherein

the stereochemistry of carbon (a) is (S);

R₁ is 5-chloro;

-149-

R_{10} and R_{11} are H;

R_4 is benzyl;

R_8 is methyl; and

R_9 is 3-pyridyl.

- 5 25. The compound as recited in claim 21 wherein
the stereochemistry of carbon (a) is (S);

R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

10 R_8 is methyl; and

R_9 is 2-hydroxyethyl.

26. The compound as recited in claim 21 wherein
the stereochemistry of carbon (a) is (S);

R_1 is 5-fluoro;

15 R_{10} and R_{11} are H;

R_4 is 4-fluorophenylmethyl;

R_8 is methyl; and

R_9 is 2-morpholinoethyl.

27. A compound as recited in claim 22 wherein

20 R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

R_8 is methyl; and

R_9 is 2-hydroxyethoxy.

- 25 28. The compound as recited in claim 22 wherein
the stereochemistry of carbon (a) is (S);

R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is 4-fluorophenylmethyl;

30 R_8 is methyl; and

R_9 is methoxy.

29. The compound as recited in claim 22 wherein
the stereochemistry of carbon (a) is (S);

-150-

R_1 is 5-chloro;

R_{10} and R_{11} are H;

R_4 is benzyl;

R_8 is methyl; and

5 R_9 is methoxy.

30. A compound as recited in claim 1 wherein

R_1 is 5-halo, 5-methyl, 5-cyano or trifluoromethyl;

R_{10} and R_{11} are each independently H or halo;

A is $-C(H)=$;

10 R_2 and R_3 are H;

R_4 is H, phenyl(C_1 - C_2)alkyl, thien-2- or -3-yl(C_1 - C_2)alkyl, fur-2- or -3-yl(C_1 - C_2)alkyl wherein said rings are mono- or di-substituted independently with H or fluoro;

R_5 is H; and

15 R_6 is (C_1 - C_5)alkoxycarbonyl.

31. A compound as recited in claim 1 wherein

R_1 is 5-halo, 5-methyl, 5-cyano or trifluoromethyl;

R_{10} and R_{11} are each independently H or halo;

A is $-C(H)=$;

20 R_2 and R_3 are H;

R_4 is H, methyl or phenyl(C_1 - C_2)alkyl, wherein said phenyl groups are mono- or di-substituted independently with H, halo, (C_1 - C_4)alkyl, (C_1 - C_4)alkoxy, trifluoromethyl, hydroxy, amino or cyano and wherein said phenyl groups are additionally mono- or di-substituted independently H or halo; or

25 R_4 is thien-2- or -3-yl(C_1 - C_2)alkyl, pyrid-2-, -3- or -4-yl(C_1 - C_2)alkyl, thiazol-2-, -4- or -5-yl(C_1 - C_2)alkyl, imidazol-2-, -4- or -5-yl(C_1 - C_2)alkyl, fur-2- or -3-yl(C_1 - C_2)alkyl, pyrrol-2- or -3-yl(C_1 - C_2)alkyl, oxazol-2-, -4- or -5-yl(C_1 - C_2)alkyl, pyrazol-3-, -4- or -5-yl(C_1 - C_2)alkyl, isoxazol-3-, -4- or -5-yl(C_1 - C_2)alkyl, isothiazol-3-, -4- or -5-yl(C_1 - C_2)alkyl, pyridazin-3- or -4-yl(C_1 - C_2)alkyl, pyrimidin-2-, -4-, -5- or -6-yl(C_1 - C_2)alkyl, pyrazin-2-
30 or -3-yl(C_1 - C_2)alkyl or 1,3,5-triazin-2-yl(C_1 - C_2)alkyl wherein said preceding R_4 heterocycles are optionally mono- or di-substituted independently with halo, trifluoromethyl, (C_1 - C_4)alkyl, (C_1 - C_4)alkoxy, amino or hydroxy and said mono- or di-substituents are bonded to carbon;

-151-

R_5 is H; and

R_6 is carboxy.

32. A compound as recited in claim 31 wherein

R_{10} and R_{11} are H; and


5 R_4 is H.

33. The compound as recited in claim 32 wherein

R_1 is 5-chloro.

34. An intermediate of Formula QZ

10



Formula QZ

15

wherein

R_5 is H;

R_4 is H, phenylmethyl, thien-2- or -3-ylmethyl, fur-2- or -3-ylmethyl wherein said rings are optionally mono- or di-substituted with fluoro; and

20 R_{12} is thiazolidin-3-yl, 1-oxothiazolidin-3-yl, 1,1-dioxothiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, azetidin-1-yl, 1,2-oxazinan-2-yl, isoxazolidin-2-yl, isothiazolidin-2-yl, 1,2-oxazetidin-2-yl or oxazolidin-3-yl,

wherein said R_{12} rings are optionally mono- or di-substituted independently with halo, (C₁-C₅)alkyl, (C₁-C₅)alkoxy, hydroxy, amino, mono-N- or di-N,N-(C₁-C₅)alkylamino, formyl, carboxy, carbamoyl, mono-N- or di-N,N-(C₁-

25 C₅)alkylcarbamoyl, (C₁-C₅)alkoxycarbonyl, hydroxy(C₁-C₅)alkyl, amino(C₁-C₄)alkyl, mono-N- or di-N,N-(C₁-C₄)alkylamino(C₁-C₄)alkyl, oxo, hydroxyimino or (C₁-C₆)alkoxyimino with the proviso that only the R_{12} heterocycles thiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, azetidin-1-yl, 1,2-oxazinan-2-yl, isoxazolidin-2-yl, or oxazolidin-3-yl are optionally mono- or di-substituted independently with oxo,

30 hydroxyimino, or (C₁-C₆)alkoxyimino; and

wherein said R_{12} rings are optionally additionally mono- or di-substituted independently with (C₁-C₅)alkyl and

-152-

with the proviso that R_{12} is not 2-carboxy-4-hydroxy-pyrrolidin-1-yl, 2-((C_1 - C_5)alkoxycarbonyl)-4-hydroxy-pyrrolidin-1-yl, 2-carboxy-piperidin-1-yl or 2-((C_1 - C_5)alkoxycarbonyl)-piperidin-1-yl.

35. A compound as recited in claim 34 wherein

- 5 R_4 is phenylmethyl, said phenyl optionally mono- or di-substituted with fluoro;
and

R_{12} is 3-mono-substituted azetidin-1-yl, 3-mono- or 3,4-disubstituted pyrrolidin-1-yl, 3-, 4-, or 5- mono- or di-substituted piperidin-1-yl, thiazolidin-3-yl, 1-oxo-thiazolidin-3-yl or 1,1-dioxothiazolidin-3-yl wherein said pyrrolidin-1-yl or
10 piperidin-1-yl are mono- or di-substituted independently with hydroxy, oxo, hydroxyimino, amino, mono-N- or di-N,N-(C_1 - C_4)alkylamino, (C_1 - C_5)alkoxycarbonyl or carboxy

and said R_{12} rings are optionally additionally mono- or di-substituted independently with (C_1 - C_4)alkyl.

- 15 36. The compound as recited in claim 34 wherein

R_4 is H; and

R_{12} is thiazolidin-3-yl.

37. The compound as recited in claim 34 wherein

R_4 is H; and

- 20 R_{12} is 1,1-dioxo-thiazolidin-3-yl.

38. A compound as recited in claim 34 wherein

R_4 is H; and

R_{12} is 1-oxo-thiazolidin-3-yl.

39. A compound as recited in claim 35 wherein

- 25 R_4 is benzyl;

R_{12} is 3-hydroxypyrrolidin-3-yl; and

the stereochemistry of carbon (a) is (S).

40. The compound as recited in claim 35 wherein

R_4 is benzyl;

- 30 R_{12} is 3-hydroxyazetidin-1-yl; and

the stereochemistry of carbon (a) is (S).

41. A compound as recited in claim 35 wherein

R_4 is benzyl;

-153-

R_{1,2} is 3,4-dihydroxypyrrolidin-1-yl; and
the stereochemistry of carbon (a) is (S).

42. The compound as recited in claim 35 wherein

R₄ is benzyl;

5 R_{1,2} is 4-hydroxypiperidin-1-yl; and
the stereochemistry of carbon (a) is (S).

43. The compound as recited in claim 35 wherein

R₄ is 4-fluorophenylmethyl;

R_{1,2} is 4-hydroxypiperidin-1-yl; and

10 the stereochemistry of carbon (a) is (S).

44. The compound as recited in claim 35 wherein

R₄ is benzyl;

R_{1,2} is 4-hydroxyiminoazetidin-1-yl; and

the stereochemistry of carbon (a) is (S).

15 45. A method for treating a glycogen phosphorylase dependent disease or
condition in a mammal which comprises administering to a mammal suffering from a
glycogen phosphorylase dependent disease or condition a glycogen phosphorylase
dependent disease or condition treating amount of a compound of claim 1.

46. The method as recited in claim 45 for treating hyperglycemia in a mammal by
20 administering to a mammal suffering from hyperglycemia a hyperglycemia treating
amount of a compound of claim 1.

47. The method as recited in claim 45 for treating diabetes in a mammal by
administering to a mammal suffering from diabetes a diabetes treating amount of a
compound of claim 1.

25 48. The method as recited in claim 45 for treating hypercholesterolemia in a
mammal by administering to a mammal suffering from hypercholesterolemia a
hypercholesterolemia treating amount of a compound of claim 1.

49. The method as recited in claim 45 for treating atherosclerosis in a mammal by
administering to a mammal suffering from atherosclerosis an atherosclerosis treating
30 amount of a compound of claim 1.

50. The method as recited in claim 45 for treating hyperinsulinemia in a mammal
by administering to a mammal suffering from hyperinsulinemia a hyperinsulinemia
treating amount of a compound of claim 1.

-154-

51. The method as recited in claim 45 for treating hypertension in a mammal by administering to a mammal suffering from hypertension a hypertension treating amount of a compound of claim 1.
52. The method as recited in claim 45 for treating hyperlipidemia in a mammal by administering to a mammal suffering from hyperlipidemia a hyperlipidemia treating amount of a compound of claim 1.
53. The method as recited in claim 45 for preventing a myocardial ischemic injury in a mammal by administering to a mammal at risk for perioperative myocardial ischemic injury a perioperative myocardial ischemic injury preventing amount of a compound of claim 1.
54. A pharmaceutical composition which comprises a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.
55. The pharmaceutical composition as recited in claim 54 for the treatment of glycogen phosphorylase dependent diseases or conditions in mammals which comprises a glycogen phosphorylase dependent disease or condition treating amount of a compound of claim 1 and a pharmaceutically acceptable carrier.
56. A pharmaceutical composition which comprises a therapeutically effective amount of
- a) a glycogen phosphorylase inhibitor;
 - b) an antidiabetic agent selected from insulin and insulin analogs; insulinotropin; Sulfonylureas and analogs; Biguanides; α 2-Antagonists and Imidazolines; insulin secretagogues; Glitazones; Fatty Acid Oxidation inhibitors; α -Glucosidase inhibitors; β -Agonists; Phosphodiesterase Inhibitors; Lipid-lowering Agents; Antiobesity Agents; Vanadate and vanadium complexes and peroxovanadium complexes; Amylin Antagonists; Glucagon Antagonists; Gluconeogenesis Inhibitors; Somatostatin Analogs; Antilipotic Agents; and
 - c) optionally a pharmaceutically acceptable carrier.
57. The pharmaceutical composition as recited in claim 56 wherein the glycogen phosphorylase inhibitor is a compound of claim 1.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IB 95/00442

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D209/42 A61K31/40 C07D403/12 C07D401/12 C07D417/12
C07D401/14 C07D409/12 C07D413/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	No relevant documents disclosed -----	

☐ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

17 January 1996

Date of mailing of the international search report

29 -01- 1996

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

11. national application No.

PCT/IB 95/00442

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark : Although claims 45-53 are directed to a method of treatment of (diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.